

PM 22 400-398

1984

Proposed New Label,
marked up to indicate
changes.

HARVADE®-25F

HARVEST GROWTH REGULANT FOR COTTON

Net contents:

COMPOSITION

Active Ingredient: (% by weight)	
dimethipin* (2,3-dihydro-5,6-dimethyl-1,4-dithiin-1,1,4,4-tetraoxide)	22.4%**
Inert Ingredients:	77.6%
Total:	100.0%

*U.S. Patent No. 3,920,438

**Contains 2 pounds of active ingredient per U. S. gallon.

***Contains 240 grams of active ingredient per liter.

KEEP OUT OF REACH OF CHILDREN

DANGER • PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN: Wash thoroughly with soap and water. Call a physician immediately.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. Note to Physicians: Probable mucosal damage may contraindicate use of gastric lavage.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Contact a physician.

IF INHALED: Remove to fresh air; if abnormal symptoms develop, contact a physician.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive, causes burns. Do not get in eyes, on skin or on clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapors.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear: Coveralls over long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; chemical-resistant footwear plus socks; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(c)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not apply when weather conditions favor drift from the treated area.

ACCEPTED

DEC 31 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for this pesticide registered under EPA Reg. No. 400-398



Uniroyal Chemical Company, Inc., Middlebury, CT 06749

EPA Reg. No. 400-398

EPA Est. No.

007WPS

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; chemical-resistant footwear plus socks; chemical-resistant headgear for overhead exposure.

GENERAL INFORMATION

HARVADE®-25F is a plant growth regulator that affects certain plant processes which lead to defoliation in cotton. Since HARVADE-25F must enter the plant through the leaves for optimum activity, it must be applied in a manner that will allow maximum penetration through the leaf surface. Complete coverage of the entire plant is essential for optimum effectiveness. See APPLICATION NOTES for additional important information.

HARVADE-25F also may improve the harvest operation by desiccating most mature morning glory species. Desiccation/defoliation of other weed species such as sicklepod, croton and taaweed has been observed. Good coverage of the above weeds is necessary for best results.

MIXING INSTRUCTIONS

ALWAYS SHAKE OR STIR WELL BEFORE USE. Fill the spray tank with 2/3 of the required amount of water. Add the desired amount of HARVADE-25F and remaining required amount of water. The HARVADE-25F will suspend easily with minimum agitation. To assure that container is emptied - **TRIPLE RINSE (OR EQUIVALENT) INTO SPRAY TANK. PRESSURE RINSE IS PREFERRED.**

COMPATIBILITY

HARVADE-25F is compatible with most commonly used cotton desiccants, defoliant, and insecticides. HARVADE-25F has been shown to be incompatible with EPN when used as a tank mix. Consult your supplier or Uniroyal Chemical for details.

USE PRECAUTIONS

1. Do not plant rotational crops within 8 months after application.
2. Keep HARVADE-25F drift from non-target areas. Do not apply when weather conditions favor drift from treated areas.
- 3a. Do not apply HARVADE-25F directly to, or permit mist to drift onto, citrus, broccoli, almonds, grapes, peppers, tomatoes, lettuce, alfalfa, sugarbeets or wheat. These crops may sustain significant injury if contacted with HARVADE-25F.
- b. Do not apply HARVADE-25F to, or permit mist to drift onto, soil that will subsequently be seeded to peppers, tomatoes, lettuce, broccoli, alfalfa or sugarbeets. Significant crop injury may occur from such an application.

- c. Do not apply HARVADE-25F to, or permit mist to drift onto, irrigation water that will subsequently be applied to peppers, tomatoes or lettuce.

DIRECTIONS FOR USE ON RAIN-BELT COTTON

For use in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, South Carolina, Tennessee, Virginia, eastern Texas (including the Rio Grande Valley) and eastern Oklahoma.

APPLICATION EQUIPMENT

Nearly all spray application equipment is suitable for applying HARVADE-25F if fine spray droplets can be deposited. If aerial equipment is used, it should be fitted with a fixed boom set at 45 to 90° relative to the trailing edge of the wing and flown 10 to 12 feet above the crop canopy.

TIMING OF APPLICATION ON RAIN-BELT COTTON

Apply 7 to 14 days prior to anticipated harvest. Optimum defoliation will be obtained when HARVADE-25F is applied to mature cotton plants; i.e. when 70% or more of the bolls are open, when the last boll to be harvested is not more than four nodes above the last cracked boll showing fiber (4NACB), when the last boll to be harvested is hard to cut through and seed coat is light brown, or when there is no active growth and when lower leaves have purple tinge.

APPLICATION RATE ON RAIN-BELT COTTON

One gallon of HARVADE-25F will treat 8 acres of cotton, when applied at the 1 pint (16 fl. oz.) per acre rate.

Aerial Application: Mix 1 to 1.25 pint (16 to 20 fl. oz.) of HARVADE-25F in a minimum of 5 gallons of spray solution per acre. Use the higher rate when the plants are more vigorous and robust. If cotton is rank or conditions favor regrowth, see APPLICATION NOTES.

Ground Application: Mix 1 to 1.25 pint (16 to 20 fl. oz.) of HARVADE-25F in 10 to 20 gallons of spray solution per acre. Use the higher rate when the plants are more vigorous and robust. If cotton is rank or conditions favor regrowth, see APPLICATION NOTES.

APPLICATION NOTES FOR RAIN-BELT COTTON

Partial defoliation from the first application of HARVADE-25F may result from one or more of the following situations:

- Rank cotton foliage is dense enough to prevent spray droplets from reaching the bottom leaves.
- Plants are actively growing, with new growth and fruiting evident at the top.
- Less than 70% of the bolls are open.
- Cotton is drought stressed. (Under this condition, applications should be made before mid morning or at late afternoon).
- Night temperatures are below 60°F for 3 to 4 nights either before or after application.
- Rainfall occurs within 6 hours after application.

Under these conditions a second application of HARVADE-25F of 1 pint (16 fl. oz./A) may be required 5 to 7 days after the initial application. Do not use more than a total of 2.25 pints (36 fl. oz.) of HARVADE-25F per acre per year.

TANK MIX COMBINATIONS FOR RAIN-BELT COTTON

Before using a tank mix combination, read each product label carefully and follow the Precautionary Statements on each label:

1. HARVADE-25F plus DROPP® 50WP

Under conditions that promote excessive regrowth, a tank mix of HARVADE-25F at 1 to 1.25 pint (16 to 20 fl. oz./A) plus DROPP 50WP at 0.1 to 0.2 pounds/A provides excellent regrowth control. Use the lower rate of both products when the night temperature is above 65°F; use the higher rates when the night temperature falls below 65°F.

DROPP® is a registered trademark of AgrEvo USA Co.

2. HARVADE-25F plus DEF® (or other phosphoric defoliants)

If cotton is rank or actively growing, HARVADE-25F at 1 to 1.25 pint (16 to 20 fl. oz./A) may be applied with 8 to 16 fl. oz./A of DEF to enhance defoliation. Use the lower rate of both products when the

BEST AVAILABLE COPY

night temperature is above 70°F, use the higher rates when the night temperature falls below 70°F.

DEF[®] is a registered trademark of Bayer Corp.

3. **HARVADE-5F plus an ethephon product** (such as PREP[®], SUPER BOLL[™], PLUCK[®], or ETHEPHON 6)
To enhance boll opening HARVADE-5F may be applied at the labeled rate with ethephon at the rate of 1 to 2 pounds active ingredient/A. Apply this tank mixture when the bolls are 60 to 70% open.

PREP[®] is a registered trademark of Rhone-Poulenc Ag. Company

SUPER BOLL[™] is a trademark of Griffin Corp.

PLUCK[®] is a registered trademark of Cedar Chemical Corp.

ETHEPHON 6 is distributed and sold by Micro-Flo Company

4. **HARVADE-25F plus STARFIRE[®]**
On rain-belt cotton if a high percentage of cotton leaves are still actively growing, 4 to 6 fl. oz./A of STARFIRE may be added to the labeled rate of HARVADE-25F for more complete defoliation.

STARFIRE[®] is a registered trademark of a Zeneca Group company

DIRECTIONS FOR USE ON STRIPPER AND OTHER IRRIGATED COTTON

For use in New Mexico and the High Plains of western Texas and Oklahoma.

APPLICATION EQUIPMENT

Nearly all spray application equipment is suitable for applying HARVADE-25F if fine spray droplets can be deposited. If aerial equipment is used, it should be fitted with a fixed boom set at 45 to 90° relative to the trailing edge of the wing and flown 10 to 12 feet above the crop canopy.

TIMING OF APPLICATION ON STRIPPER AND OTHER IRRIGATED COTTON

Optimum defoliation will be obtained when HARVADE-25F is applied to mature cotton. It may be used alone or tank mixed with other registered cotton harvest aids.

Under growing conditions for stripper cotton, HARVADE-25F and HARVADE-25F tank mixes will initiate the defoliation process if applied when the last bolls to be harvested are mature. This normally occurs 14 to 21 days before harvest. Application of CYCLONE[®] brand desiccant 7 to 10 days after HARVADE-25F is necessary to complete the defoliation process.

*CYCLONE is a registered trademark of a Zeneca Group company

RATE OF APPLICATION ON STRIPPER AND OTHER IRRIGATED COTTON

One gallon of HARVADE-25F will treat 8 acres of cotton, when applied at the 1 pint (16 fl. oz.) per acre rate.

Aerial Application: Mix 1.25 pint (20 fl. oz.) of HARVADE-25F in a minimum of 2 gallons of spray solution per acre. Use CYCLONE at 16 fl. oz./A at 7 to 10 days after HARVADE-25F to complete the defoliation process.

Ground Application: Mix 1.25 pint (20 fl. oz.) of HARVADE-25F in 10 to 20 gallons of spray solution per acre. Use CYCLONE at 16 fl. oz./A at 7 to 10 days after HARVADE-5F to complete the defoliation process.

TANK MIX COMBINATIONS FOR STRIPPER AND OTHER IRRIGATED COTTON

Before using a tank mix combination, read each product label carefully and follow the Precautionary Statements on each label.

1. **HARVADE-25F plus DEF (or other phosphate defoliants)**
To initiate defoliation on cotton that may have been water-stressed at some time during the growing season, HARVADE-25F may be applied at 1.25 pint (20 fl. oz.)/A with 8 fl. oz./A of DEF. Use CYCLONE at 16 fl. oz./A at 7 to 10 days after the HARVADE-25F/phosphate defoliant application to complete the defoliation process.
2. **HARVADE-25F plus a formulated sodium chlorate product**
To provide good defoliant initiation if cotton has been stressed over

an extended period of time or has any new growth, HARVADE-25F at 0.5 pint (8 fl. oz.)/A may be applied with 3 lb. a.i./A of sodium chlorate (example: 5 pints/A of a 5 lb/gal formulated product or 4 pints/A of a 6 lb/gal product). Use CYCLONE at 16 fl. oz./A at 7 to 10 days after the HARVADE-25F/sodium chlorate application to complete the defoliation process.

3. **HARVADE-25F plus an ethephon product** (such as PREP, SUPER BOLL, PLUCK, or ETHEPHON 6)
To enhance harvest initiation and boll opening, HARVADE-25F at 1.25 pint (20 fl. oz.)/A may be applied with an ethephon product at the rate of 1 to 2 pounds active ingredient/A. Use CYCLONE at 16 fl. oz./A at 7 to 10 days after the HARVADE/ethephon application to complete the defoliation process.

4. **HARVADE-25F plus DROPP 50WP**
Under conditions that promote excessive regrowth, HARVADE-25F at 1.25 pint (20 fl. oz.)/A may be used with DROPP 50WP at 0.1 to 0.2 lb./A. Use CYCLONE at 16 fl. oz./A at 7 to 10 days after the HARVADE-25F/DROPP 50W application to complete the defoliation process.

5. **HARVADE-25F plus ROUNDUP[®]**
If perennial weeds such as johnsongrass are prevalent, HARVADE-25F is compatible with ROUNDUP. Refer to the ROUNDUP label for further application information.

ROUNDUP[®] is a registered trademark of Monsanto Company

DIRECTIONS FOR USE ON COTTON IN CALIFORNIA

GENERAL INFORMATION

HARVADE-25F is an excellent cotton defoliant to use in tank mix combination with other harvest aid products. Application to Acala SJ-2 and related varieties is not recommended.

APPLICATION EQUIPMENT

Nearly all spray application equipment is suitable for applying HARVADE-25F if fine spray droplets can be deposited. If aerial equipment is used, it should be fitted with a fixed boom set at 45 to 90° relative to the trailing edge of the wing and flown 10 to 12 feet above the crop canopy.

TIMING OF APPLICATION IN CALIFORNIA

Optimum defoliation will be obtained when HARVADE-25F is applied to mature cotton. In California, it is most frequently tank mixed with other registered cotton harvest aids.

Apply to mature plants, when 70% or more of the bolls are open (4NACB, when the last boll to be harvested is no more than four nodes above the last cracked boll, showing fiber).

Under California conditions, if harvest is desired within 14 to 21 days of the HARVADE-25F application, a follow-up treatment with an approved crop desiccant may be necessary. Apply the desiccant according to its label directions.

RATE OF APPLICATION IN CALIFORNIA

One gallon of HARVADE-25F will treat 8 acres of cotton, when applied at the 1 pint (16 fl. oz.) per acre rate.

Aerial Application: Mix 1 pint (16 fl. oz.) of HARVADE-25F with the desired amount of tank mix partner product, in a minimum of 10 gallons of spray solution per acre.

Ground Application: Mix 1 pint (16 fl. oz.) of HARVADE-25F with the desired amount of tank mix partner product, in a minimum of 10 to 20 gallons of spray solution per acre.

TANK MIX COMBINATIONS FOR USE IN CALIFORNIA

Before using a tank mix combination, read each product label carefully and follow the Precautionary Statements on each label.

Use each product in a tank mix in accordance with its label Directions for Use under California conditions.

To achieve optimum defoliation under local conditions, HARVADE-25F may be tank mixed with one of the following tank mix partners, registered

BEST AVAILABLE COPY

for use on cotton in California.

- DEF (or other phosphate defoliant product),
- sodium chlorate products,
- GINSTAR[®], or
- DROPP.

*GINSTAR is a registered trademark of AgrEvo USA Co.

When additional crop desiccation or weed control is desired, other registered harvest aid products such as paraquat or ROUNDUP may be tank mixed with HARVADE-25F.

Do not mix HARVADE-25F with cacodylic acid products, because they are antagonistic to HARVADE-25F activity.

ALWAYS SHAKE OR STIR THIS PRODUCT WELL BEFORE USE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate and dry. Keep from freezing. Do not store below 40°F. ALWAYS SHAKE OR STIR WELL BEFORE USE.

PESTICIDE DISPOSAL: Pesticides wastes are acutely hazardous. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic Containers: Triple rinse (or equivalent). Pressure rinse is preferred. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

® HARVADE is a Registered Trademark of Uniroyal Chemical Company, Inc.

©Copyright 1996, Uniroyal Chemical Company, Inc.

BEST AVAILABLE COPY