

MAY 21 1992

Willard F. Cummings  
 U.S. Registration Manager  
 Crop Protection Research & Development  
 Uniroyal Chemical Company, Inc.  
 74 Amity Road  
 Bethany, Connecticut 06525

Subject: Terraclor 75 WP  
 EPA Reg. No. 400-399  
 RE: Label Amendment (Rate limitations)  
 Your Letter Dated February 22, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

\* The 7.0 lb ai/A transplant treatment for tomatoes is inconsistent with other instructions on label. Please make changes as specified in 5b and 5c on page 14 of attached review.

In addition, please be advised that while the revised maximum seasonal rate should result in lower residue levels of PCNB in or on beans, available field trial data indicate that residue levels could still exceed the interim tolerance of 0.1 ppm. Please indicate to the Agency within 30 days of the date stamped on this letter how you intend to address this issue. Failure to respond could result in the suspension of this product. Please see attached review for details.

Sincerely yours,

Robert J. Taylor  
 Product Manager (25)  
 Fungicide-Herbicide Branch  
 Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C						
SURNAME	moll						
DATE	5/19/92						

# TERRACLOR

## 75% WETTABLE POWDER

### VEGETABLE, TURF & ORNAMENTAL

### SOIL FUNGICIDE

**ACCEPTED**  
with **COMMENTS**  
in EPA Letter Dated:

MAY 21 1991

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

400-399

**UNIROYAL**  
**CHEMICAL**

#### COMPOSITION

Active Ingredients: (% by weight)	
Pentachloronitrobenzene (PCNB) .....	75%
Inert Ingredients .....	25%
Total: .....	100%

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749  
EPA REG. NO. 400-399  
EPA EST. NO.

TERRACLOR is a Registered Trademark of Uniroyal Chemical Company, Inc.  
©Copyright 1991, Uniroyal Chemical Company, Inc.

009

#### NET WEIGHT

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.  
**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.  
**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.  
**IF INHALED:** Remove to fresh air. See a physician if an abnormal reaction occurs.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

**BEST AVAILABLE COPY**

Harmful if swallowed. Avoid inhaling dust. May cause skin irritation. Wash hands and face thoroughly after using. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes. Wash thoroughly all contaminated clothing before reuse.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT PLANT ROOT CROPS IN PCNB TREATED FIELDS WITHIN 12 MONTHS OF BROADCAST AND BANDING APPLICATIONS UNLESS PCNB IS REGISTERED FOR USE ON THOSE CROPS.

### GENERAL INFORMATION

TERRACLOR 75%WP is a protective fungicide recommended for control of certain soil borne diseases of vegetable and ornamental crops. Best results are achieved when used in a preventative program following recommended rates and application directions. Carefully read, understand and follow label directions.

### VEGETABLES

TERRACLOR 75%WP should be applied as a preventative fungicide to vegetable crops. It is important when using TERRACLOR 75%WP as a protective fungicide that material be placed into the soil. This can be accomplished as a soil drench, mechanical incorporation or as an

infurrow application. Under certain crop use (snap & dry beans), repeated applications may be necessary. See use and rate recommendations for specific crop use recommendations and diseases controlled by TERRACLOR 75%WP.

When rate range is shown the lower rate should be used on lighter soils or disease pressure is expected to be low. The higher rates should be used on heavy soils, in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

### CROP

BEANS (snap & dry)

PEPPERS, TOMATOES

GARLIC

BROCCOLI

BRUSSELS SPROUTS

CABBAGE

CAULIFLOWER

### DISEASES CONTROLLED

Root & Stem Rot (*Rhizoctonia solani*)  
White Mold (*Sclerotium sclerotiorum*)

Southern Blight (*Sclerotium rolfsii*)

White Rot (*Sclerotium capworum*)

Wire Stem or Black Root (*Corticium solani*)

Club Root (*Plasmodiophora brassicae*)

**BEST AVAILABLE COPY**

CROP	DISEASE	RAT. TERRACLOR 75%WP		RECOMMENDED APPLICATION
		POUNDS/ACRE	OUNCE/1000 FEET OF ROW	

COTTON	Damping off (Rhizoctonia solani)	1 1/3 to 2 3/4 Apply in at least 15 gals. of water. (40 inch row spacings)	1.6 to 3.4	Apply specified dosage as a directed in-furrow spray to the seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.
	Apply only at planting time. Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.			

PEANUTS	Root and Stem Rot (Rhizoctonia solani)	1 1/3 to 2 3/4 Apply in 8 to 10 gallons of water. (36 inch row spacing)	1.5 to 3.0	AT PLANTING: Apply specified dosage as a directed in-furrow spray to the seed and surrounding soil at planting time. Do not use in California.
	Southern stem and root rot (Sclerotium rolfsii)	13 1/3 Apply in 8 to 10 gallons of water. (36 inch row spacing)	14.7	AT-PEGGING BAND: Apply specified dosage as a spray in a 12 inch band centered on the row so as to cover the crown and pegging zones of the plant. Do not move any untreated soil into the treated area.
		4 1/3 Apply in 8 to 10 gallons of water. (36 inch row spacing)	4.8	AT CULTIVATION: Apply specified dosage at cultivation. Do not apply more than 3 times per crop season. Attach two spray nozzles per row of inside cultivator sweeps (one on each side) and adjust so as to treat a total band of 12 inches centered on the row. Do not pile untreated soil on crown of plant.
Do not exceed 10 lbs. PCNB active ingredient per acre in any one season, regardless of the method of application or formulation used. Do not apply within 45 days of harvest. Do not feed treated vines to livestock. Do not allow hogging down of peanuts.				

BEANS Dry, Succulent & Snap	Root and Stem Rot (Rhizoctonia solani)	1 1/3 to 2 Apply in 8 to 10 gallons of water. (36 inch row spacing)	1.4 to 2.2	Apply as a directed spray in the seed furrow and to the covering soil at planting time.
	White Mold (Sclerotinia sclerotiorum)	2 3/4 Apply in 8 to 10 gallons of water. (36 inch row spacing)	3	Apply as an 8 inch band centered on the row at or immediately after planting. If the disease problem is severe, applications may be repeated at 2 to 3 week intervals using directed spray application with one or two nozzles per row aimed at the soil base of the plants. Soil should remain undisturbed after application.
Do not exceed 8 lbs. PCNB active ingredient per acre in any one season, regardless of the method of application or formulation used. Do not apply after first bloom. Do not feed treated vines or hay to livestock. PREHARVEST INTERVALS: Dry beans - Do not apply within 45 days of harvest; Succulent beans - Do not apply within 35 days of harvest; Snap beans - Do not apply within 14 days of harvest.				

BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS* KALE* MUSTARD GREENS*	Clubroot (plasmodiophora brassicae)	Mix 2 pounds in 100 gallons of water. Use 1/2 pint of this solution per plant.		Transplant Solution: The applied dosage of TERRACLOR should be based on the severity of the disease infection. Maintain agitation to hold the powder in suspension
		- OR -		
		30 Apply in 25 gallons of water. (40 inch row spacings)	37	Band Application: Spray as a 12 to 15 inch band centered on the row and incorporate to a depth of 4 to 6 inches immediately prior to planting. May be used on cole crops which are direct seeded
	- OR -			
	40 Apply in 30 gallons of water.	-	Broadcast Application: Apply as a preplant broadcast prior to planting. Thoroughly incorporate to a depth of 4 to 6 inches using a disc or other suitable equipment. May be used on cole crops which are direct seeded	
	Note: for effective control of clubroot, it is extremely important to thoroughly mix TERRACLOR with the soil.			
	- OR -			
	Wire Stem or Black Root (Corticium solani)	15 to 20 Apply in 50 gallons of water	-	Broadcast Drench Application: Apply as a soil drench at the time of or immediately after seeding. For smaller areas use 1 level tablespoon per gallon of water as a soil drench per 50 sq ft of seed bed. Apply with water can or similar equipment
- OR -				
	10 to 15 Apply in 35 gallons of water (40 inch row spacings)	12.2 to 18.4	Row Drench Treatment: Spray as an 8 inch band centered on the row at the time of or immediately after seeding	
*Based on the available residue data, the use of TERRACLOR on collards, kale and mustard greens is restricted to the state of Georgia. Apply only at planting time. Do not exceed 30 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.				

**BEST AVAILABLE COPY**

CROP	DISEASE	RATE OF TERRACLOR 75%WP		RECOMMENDED APPLICATION
		POUNDS/ACRE	DOUNCE/1000 FEET OF ROW	
GARLIC	White Rot (Sclerotium capivorum)	27½ Apply in 100 gallons of water. (24 inch row spacings)	20	In-Furrow Application: Apply as an in-furrow spray at planting time. Attach a spray rig to the planter and spray 8 - 12 inch bands using 2 nozzles per furrow. Direct front nozzle to spray the "seed" as it is dropped; direct the rear nozzle to spray the covering soil. Use good mechanical agitation in the spray tank to maintain a uniform suspension.
		- OR -		
		13½ pounds per 1,000 lbs. of "seed" cloves.		Clove Treatment: Mix in sufficient water to apply dosage as a uniform mist-spray for complete coverage of treated cloves. For optimum results add a commercial sticker to the spray solution.
Apply only at planting time. Do not exceed 21 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.				

PEPPERS TOMATOES	Southern Blight (Sclerotium rolfsii)	Mix 3 pounds in 100 gallons of water. Use ½ pint of this solution per plant.		Transplant Solution: Apply at the time of transplanting. The solution should be agitated often to maintain a uniform suspension and assure proper dosage. Apply only at planting time. TOMATOES - Do not exceed 7.0 lbs. PCNB active ingredient per acre in any one season. PEPPERS - Do not exceed 20 lbs. PCNB active ingredient per acre in any one season.
		- OR -		
		6½ to 10 Apply in 100 gallons of water. (36 inch row spacings peppers; 72 inch row spacings flat tomatoes; 48 inch row spacings staked tomatoes)	Peppers: 11 Tomatoes: 14.6	In-Furrow: Apply as an in-furrow spray to the open "V" trench, just prior to planting. When cultivating, set plows as flat as possible to avoid getting untreated soil against stems of plants. Apply only at planting time. Do not exceed 7.5 lbs. of PCNB active ingredient per acre in any one season.

**ROTATION RESTRICTION:** Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.

**GENERAL INFORMATION**

TERRACLOR 75%WP is a preventative fungicide recommended for control of certain soil borne diseases of turf and certain ornamental crops. Best results are achieved when used in preventative program

following recommended rates and application directions. Carefully read, understand and follow label directions.

**TURF**

TERRACLOR 75%WP is a fungicide for the prevention and control of certain soil borne diseases on golf courses, sod farms, home lawns and institutional areas where turf is grown.

Best results are obtained when TERRACLOR 75%WP is used as a preventative application before diseases appear; however, if you do not apply as a preventative treatment, TERRACLOR 75%WP should be applied at first sign of the disease.

**DISEASES CONTROLLED:**

- Brown Patch
- Helminthosporium Leaf Spot\*
- Dollar Spot\*
- Gray Snow Mold
- Pink Snow Mold

**Fungi**

- Rhizoctonia solani
- Helminthosporium spp.
- Sclerotinia homoeocarpa
- Typhula spp.
- Fusarium nivale

Areas treated with TERRACLOR 75%WP should be watered following application to move material down to the soil level. If treated area is subjected to unusually heavy rainfall, or flooded, or if disease is severe or reappears, the area should be treated again. See use and rate recommendations for intervals for retreating.

**USE & RATE RECOMMENDATION FOR TURF:**

DISEASE/CROP	RATE/1,000 SQ. FT.	REMARKS
Brown Patch (Rhizoctonia solani) St. Augustine, Bermuda, and similar warm climate grasses	16 oz. in 10 - 15 gallons water	1. Apply one application in the spring or fall or at the 1 <sup>st</sup> indication of disease. 2. If disease is severe or reappears, retreat 3 - 4 weeks later.
Dichondra	16 oz. in 40 gallons water	1. Treat as above.
Bluegrass, Rye Grass, Fescue, Bent and similar cool climate grasses	3 - 4 oz. in 3 - 6 gallons water	1. Apply at first sign of disease. 2. Repeat every 7 - 10 days during warm damp weather. 3. For control of existing infections, apply at higher rates as soon as symptoms appear.
Helminthosporium Leaf Spot* (Helminthosporium spp.) Dollar Spot* (Sclerotinia homoeocarpa)	7 - 10 oz. in 5 - 10 gallons water	1. Preventative applications should be in spring or fall of each year. 2. For control of existing infections, apply at higher rate as soon as symptoms appear. 3. If disease is severe or reappears, area should be retreated 3 - 4 weeks later.
Gray Snow Mold (Typhula spp.) Pink Snow Mold (Fusarium nivale)	8 oz. in 5 - 10 gallons water	1. Make one application prior to first snowfall

**\*DO NOT USE IN CALIFORNIA**

Treated area should be watered lightly as soon as possible following treatment to wash chemical down to the soil level  
Do not graze treated areas or feed clippings to livestock

Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time

**ORNAMENTALS**

TERRACLOR 75%WP can be used as soil treatment for protection against a variety of soil borne diseases in nursery, greenhouse and field grown ornamentals.

It is important that TERRACLOR 75%WP be placed or mixed in the soil for best results.

See use and rate recommendations for specific plant and disease recommendations.

When rate range is shown the lower rate should be used on lighter soils or when disease pressure is expected to be low. The higher rates should be used on heavy soils in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

**DISEASES CONTROLLED**

- Root and Stem Rot
- Storage Rot
- Neck Dry Rot
- Bulb Dry Rot
- Needle Blight
- Petal Blight
- Bulb and Stem Rot
- Flower Blight
- Crown Rot
- Black Rot

**Fungi**

- Rhizoctonia spp.
- Botrytis cinerea
- Stromatinia gladioli
- Sclerotium gladioli
- Dothistromia pini
- Ovulinia azaleae
- Sclerotium rolfsii
- Sclerotium camelliae
- Pellicularia rolfsii
- Sclerotinia sclerotiorum

**PLANT TOLERANCE** With specific plants not listed below we recommend a preliminary trial to determine plant tolerance.

**USE AND RATE RECOMMENDATION:**

TERRACLOR 75%WP as Soil Drench at Seeding or Transplanting in Nursery and Greenhouse Grown Ornamentals		
<b>Foliage Plants</b> Aglaonema      Ficus      Pothos Ajuga            Fittonia    Schefflera Anthurum       Gynura     Schlumbergera Aphelandra     Hoya       Spathiphyllum Caladium        Maranta    Syngonium Cissus            Palms       Tradescantia Dffenbachia    Peperomia   Similar foliage plants Dracena          Pilea		<b>For control of Rhizoctonia diseases.</b> <b>Application Rates:</b> Use 4 to 8 oz. in 100 gallons of water and apply to 400 - 800 sq. ft. of flat, bench or bed area.  <b>Coverage:</b> For soil depths of 4 inches or less, apply 100 gallons to 800 sq. ft. For soil depths of more than 4 inches, apply 100 gallons to 400 sq. ft.  One repeat application can be made 4 to 6 weeks later, if necessary.
<b>Do not apply to Philodendron or Pilea or injury may occur.</b>		
<b>Bedding and Flowering Plants</b> Abutilon        Dianthus    Petunia Alyssum         Foxglove    Pointsetta Calceolaria     Fuchsia     Portulaca Cannna           Gardenia    Salvia Carnation       Gazania     Scabiosa Chrysanthemum Geranium   Sedum Cineraria       Gloxinia    Sempervivum Columbine       Impatiens   Verbena Coral Bells      Lily          Vinca Cyclamen        Lupine       Zinnia Dahlia            Mangold     Similar plants Delphinium      Monarda		
<b>Woody Ornamentals</b> Arborvitae      English Ivy   Podocarpus Aucuba           Holly          Privet Azalea           Juniper       Rhododendron Boxwood         Maple         Viburnum Carissa          Pine           Similar woody plants		

**BEST AVAILABLE COPY**

Solutions of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application.

**USE AND RATE RECOMMENDATIONS:**

TERRACLOR 75%WP as a Broadcast Incorporation Prior to Planting Field Grown Crops			
Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 sq. ft. area)	Application
Easter Lilies	Root Rot (Rhizoctonia spp.)	3 1/4 to 6 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 6 - 7 inches and see note below.
Ins Hyacinth Narcissus Tulips	Crown Rot (Pellicularia rolfsii) Black Rot (Sclerotinia sclerotiorum)		
		Use lower rates for light soils and higher rates for heavy soils.	
Dutch Iris	Bulb Rot Stem Rot (Sclerotium rolfsii)	3 1/4 to 4 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 2 - 3 inches
Gladiolus	Neck or Bulb Dry Rot (Stromatinia or Sclerotinia gladioli)	3 2/3	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 1 - 6 inches Bulbs may be planted immediately after incorporation.
Calendula Larkspur Snapdragon Sweet Pea	Stem Rot Root Rot (Rhizoctonia spp.)	3 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 1 inches
Southern Pine Seedlings	Root/Stem Rot (Rhizoctonia spp.) Needle Blight (Dothistromia pini)	1 1/4	Apply specified rate in sufficient water to insure uniform ground coverage. An additional 1 1/2 inches of water per acre should be applied by irrigation prior to planting. Seed may be planted immediately after irrigation or within one week of this date

Solution of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application of the fungicide.

TERRACLOR		%WP as a Ground Spray on Established Woody C		rentals
Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 150 sq. ft. area)	Application	
Azaleas	Azalea petal blight (Ovulinia Azaleae)	1	Apply specified rate per 150 sq. ft. in sufficient water to assure uniform ground coverage. Apply to the ground beneath the bushes and immediate surrounding area. Begin application prior to opening of buds and repeat every 3 - 4 weeks during bloom season. Solution of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application of fungicide.	
Camellia	Camellia flower blight (Sclerotinia camelliae)	1		
TERRACLOR 75%WP as a Dip Application				
Crop	Disease	Rate of TERRACLOR 75%WP (Lbs.)	Application	
Iris Hyacinth Narcissus Tulips	Crown Rot (Pellicularia rolfsii) Black Rot (Sclerotinia sclerotiorum)	2	Dip bulbs for 5 minutes in 7.5% concentration which is obtained by adding 2 lbs. of TERRACLOR 75%WP to 3.2 gallons of water. Add 1% sticker to solution and maintain good agitation.	
Easter Lilies	Root Rot (Rhizoctonia spp.)	4 - 5	Mix TERRACLOR 75%WP in 100 gallons of water and dip bulbs for 5 - 15 minutes. A sticking agent may be added. Maintain good agitation in dip tanks.	
Roses	Storage Rot (Botrytis cinerea)	2	Mix TERRACLOR 75%WP in 100 gallons of water and dip dormant roses prior to storage. A spray application of the same solution may be substituted for dipping. Maintain good agitation.	
TERRACLOR 75%WP as Band Application Prior to Planting Field Grown Crops				
Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 linear ft. of row)	Application	
Gladiolus	Neck or Bulb Dry Rot (Stomatium or Sclerotinia gladioli)	5 1/2	Apply as a 12 inch band centered on the row at planting time. Incorporate to depth of 4 inches. Bulb may be planted immediately after incorporation.	
Dutch Iris	Bulb Rot Stem Rot (Sclerotium rolfsii)	1/2	Apply as a 12 inch band centered on the row at planting time. Follow with 3 additional applications at the same rate at 15 day intervals.	
TERRACLOR 75%WP as In Furrow Application at Planting				
Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 linear ft. of row)	Application	
Easter Lilies	Root Rot (Rhizoctonia spp.)	4 1/2 to 6	Apply in 5 - 10 gallons of water. Direct spray into open furrow to thoroughly cover entire furrow and bulbs. Close furrow in normal manner. See notes below.  Use lower rates for lighter soils and higher rates for heavier soils. For Black rot control, use high rates. Solution of TERRACLOR 75%WP and water should be agitated to assure uniform application of the fungicide.	
Iris Hyacinth Narcissus Tulips	Crown Rot (Pellicularia rolfsii) Black Rot (Sclerotinia sclerotiorum)			

### STORAGE AND DISPOSAL

**Do not contaminate water, food or feed by storage or disposal.**

**STORAGE:** Store in a dry location.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:**

**Paper Bags:** Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Cardboard Canisters: (Commercial)** Completely empty canisters into application equipment. Then dispose of container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Household:** Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**BEST AVAILABLE COPY**