


TERRACLOR®

75% WETTABLE POWDER
VEGETABLE, TURF & ORNAMENTAL
SOIL FUNGICIDE



COMPOSITION

Active Ingredients: (% by weight)	
Pentachloronitrobenzene (PCNB)	75%
Inert Ingredients	25%
Total:	100%

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
EPA REG. NO. 400-399
EPA EST. NO.

TERRACLOR is a Registered Trademark of Uniroyal Chemical Company, Inc.
©Copyright 1991, Uniroyal Chemical Company, Inc.

NET WEIGHT

PA 25 2 3 999
19 185

ACCEPTED
SEP 16 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-399

oil

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air. See a physician if an abnormal reaction occurs.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid inhaling dust. May cause skin irritation. Wash hands and face thoroughly after using. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes. Wash thoroughly all contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT PLANT ROOT CROPS IN PCNB TREATED FIELDS WITHIN 12 MONTHS OF BROADCAST AND BANDING APPLICATIONS UNLESS PCNB IS REGISTERED FOR USE ON THOSE CROPS.

GENERAL INFORMATION

TERRACLOR 75%WP is a protective fungicide recommended for control of certain soil borne diseases of vegetable and ornamental crops. Best results are achieved when used in a preventative program following recommended rates and application directions. Carefully read, understand and follow label directions.

VEGETABLES

TERRACLOR 75%WP should be applied as a preventative fungicide to vegetable crops. It is important when using TERRACLOR 75%WP as a protective fungicide that material be placed into the soil. This can be accomplished as a soil drench, mechanical incorporation or as an

in-furrow application. Under certain crop use (snap & dry beans), repeated applications may be necessary. See use and rate recommendations for specific crop use recommendations and diseases controlled by TERRACLOR 75%WP.

When rate range is shown the lower rate should be used on lighter soils or disease pressure is expected to be low. The higher rates should be used on heavy soils, in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

CROP

BEANS (snap & dry)

PEPPERS, TOMATOES

GARLIC

BROCCOLI

BRUSSELS SPROUTS

CABBAGE

CAULIFLOWER

DISEASES CONTROLLED

Root & Stem Rot (*Rhizoctonia solani*)
White Mold (*Sclerotium sclerotiorum*)

Southern Blight (*Sclerotium rolfsii*)
White Rot (*Sclerotium capivorum*)

Wire Stem or Black Root (*Corticium solani*)

Club Root (*Plasmodiophora brassicae*)

CROP	DISEASE	RATE OF TERRACLOR 75%WP		RECOMMENDED APPLICATION
		POUNDS/ACRE	OUNCE/1000 FEET OF ROW	

COTTON	Damping off (<i>Rhizoctonia solani</i>)	1 1/3 to 2 3/4 Apply in at least 15 gals. of water. (40 inch row spacings)	1.6 to 3.4	Apply specified dosage as a directed in-furrow spray to the seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.
	Apply only at planting time. Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.			

PEANUTS	Root and Stem Rot (<i>Rhizoctonia solani</i>)	1 1/3 to 2 3/4 Apply in 8 to 10 gallons of water. (36 inch row spacing)	1.5 to 3.0	AT PLANTING: Apply specified dosage as a directed in-furrow spray to the seed and surrounding soil at planting time. Do not use in California.
	Southern stem and root rot (<i>Sclerotium rolfsii</i>)	13 1/3 Apply in 8 to 10 gallons of water. (36 inch row spacing)	14.7	AT-PEGGING BAND: Apply specified dosage as a spray in a 12 inch band centered on the row so as to cover the crown and pegging zones of the plant. Do not move any untreated soil into the treated area.
		4 1/3 Apply in 8 to 10 gallons of water. (36 inch row spacing)	4.8	AT CULTIVATION: Apply specified dosage at cultivation. Do not apply more than 3 times per crop season. Attach two spray nozzles per row of inside cultivator sweeps (one on each side) and adjust so as to treat a total band of 12 inches centered on the row. Do not pile untreated soil on crown of plant.
Do not exceed 10 lbs. PCNB active ingredient per acre in any one season, regardless of the method of application or formulation used. Do not apply within 45 days of harvest. Do not feed treated vines to livestock. Do not allow hogging down of peanuts.				

BEANS Dry, Succulent & Snap	Root and Stem Rot (<i>Rhizoctonia solani</i>)	1 1/3 to 2 Apply in 8 to 10 gallons of water. (36 inch row spacing)	1.4 to 2	Apply as a directed spray in the seed furrow and to the covering soil at planting time.
	White Mold (<i>Sclerotinia sclerotiorum</i>)	2 3/4 Apply in 8 to 10 gallons of water. (36 inch row spacing)	3	Apply as an 8 inch band centered on the row at or immediately after planting. If the disease problem is severe, applications may be repeated at 2 to 3 week intervals using directed spray application with one or two nozzles per row, aimed at the soil base of the plants. Soil should remain undisturbed after application.
Do not exceed 6 lbs. PCNB active ingredient per acre in any one season, regardless of the method of application or formulation used. Do not apply after first bloom. Do not feed treated vines or hay to livestock. PREHARVEST INTERVALS: Dry beans - Do not apply within 45 days of harvest; Succulent beans - Do not apply within 35 days of harvest; Snap beans - Do not apply within 14 days of harvest.				

BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS* KALE* MUSTARD GREENS*	Clubroot (<i>plasmiodiophora brassicae</i>)	Mix 2 pounds in 100 gallons of water. Use 1/2 pint of this solution per plant.		Transplant Solution: The applied dosage of TERRACLOR should be based on the severity of the disease infection. Maintain agitation to hold the powder in suspension.
	- OR -			
		30 Apply in 25 gallons of water. (40 inch row spacings)	37	Band Application: Spray as a 12 to 15 inch band centered on the row and incorporate to a depth of 4 to 6 inches immediately prior to planting. May be used on cole crops which are direct seeded.
	- OR -			
	40 Apply in 30 gallons of water.	-	Broadcast Application: Apply as a preplant broadcast prior to planting. Thoroughly incorporate to a depth of 4 to 6 inches using a disc or other suitable equipment. May be used on cole crops which are direct seeded.	
Note: for effective control of clubroot, it is extremely important to thoroughly mix TERRACLOR with the soil.				
- OR -				
	Wire Stem or Black Root (<i>Corticium solani</i>)	15 to 20 Apply in 50 gallons of water.	-	Broadcast Drench Application: Apply as a soil drench at the time of or immediately after seeding. For smaller areas use 1 level tablespoon per gallon of water as a soil drench per 50 sq. ft. of seed bed. Apply with water can or similar equipment.
- OR -				
		10 to 15 Apply in 35 gallons of water. (40 inch row spacings)	12.2 to 18.4	Row Drench Treatment: Spray as an 8 inch band centered on the row at the time of or immediately after seeding.
*Based on the available residue data, the use of TERRACLOR on collards, kale and mustard greens is restricted to the state of Georgia. Apply only at planting time. Do not exceed 30 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.				

CROP	DISEASE	POUNDS/ACRE	OUNCE/1000 FEET OF ROW	RECOMMENDED APPLICATION
GARLIC	White Rot (Sclerotium cepivorum)	27½ Apply in 100 gallons of water. (24 inch row spacings)	20	In-Furrow Application: Apply as an in-furrow spray at planting time. Attach a spray rig to the planter and spray 8 - 12 inch bands using 2 nozzles per furrow. Direct front nozzle to spray the "seed" as it is dropped; direct the rear nozzle to spray the covering soil. Use good mechanical agitation in the spray tank to maintain a uniform suspension.
		- OR -		Clive Treatment: Mix in sufficient water to apply dosage as a uniform mist-spray for complete coverage of treated cloves. For optimum results add a commercial sticker to the spray solution.
Apply only at planting time. Do not exceed 21 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.				

PEPPERS TOMATOES	Southern Blight (Sclerotium rolfsii)	Mix 3 pounds in 100 gallons of water. Use ½ pint of this solution per plant.		Transplant Solution: Apply at the time of transplanting. The solution should be agitated often to maintain a uniform suspension and assure proper dosage. Apply only at planting time. TOMATOES - Do not exceed 7.0 lbs. PCNB active ingredient per acre in any one season (based on 5000 plants / acre) or 16.5 lbs. PCNB active ingredient per acre in any one season (based on 11,733 plants / acre) PEPPERS - Do not exceed 20 lbs. PCNB active ingredient per acre in any one season.
		- OR -		
		6½ to 10 Apply in 100 gallons of water. (36 inch row spacings peppers; 72 inch row spacings flat tomatoes; 48 inch row spacings staked tomatoes)	Peppers: 11 Tomatoes: 14.6	In-Furrow: Apply as an in-furrow spray to the open "V" trench, just prior to planting. When cultivating, set plows as flat as possible to avoid getting untreated soil against stems of plants. Apply only at planting time. Do not exceed 7.5 lbs. of PCNB active ingredient per acre in any one season.

ROTATION RESTRICTION: Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.

GENERAL INFORMATION

TERRACLOR 75%WP is a preventative fungicide recommended for control of certain soil borne diseases of turf and certain ornamental crops. Best results are achieved when used in preventative program

following recommended rates and application directions. Carefully read, understand and follow label directions.

TURF

TERRACLOR 75%WP is a fungicide for the prevention and control of certain soil borne diseases on golf courses, sod farms, home lawns and institutional areas where turf is grown.

Best results are obtained when TERRACLOR 75%WP is used as a preventative application before diseases appear, however, if you do not apply as a preventative treatment, TERRACLOR 75%WP should be applied at first sign of the disease.

DISEASES CONTROLLED:

DISEASES CONTROLLED:	Fungi
Brown Patch	Rhizoctonia solani
Helminthosporium Leaf Spot*	Helminthosporium spp.
Dollar Spot*	Sclerotinia homoeocarpa
Gray Snow Mold	Typhula spp.
Pink Snow Mold	Fusarium nivale

Areas treated with TERRACLOR 75%WP should be watered following application to move material down to the soil level. If treated areas are subjected to unusually heavy rainfall, or flooded, or if disease is severe or reappears, the area should be treated again. See use and rate recommendations for intervals for retreating.

USE & RATE RECOMMENDATION FOR TURF:

DISEASE/CROP	RATE/1,000 SQ. FT.	REMARKS
Brown Patch (Rhizoctonia solani) St. Augustine, Bermuda, and similar warm climate grasses	16 oz. in 10 - 15 gallons water	1. Apply one application in the spring or fall or at the 1 st indication of disease. 2. If disease is severe or reappears, retreat 3 - 4 weeks later.
Dichondra	16 oz. in 40 gallons water	1. Treat as above.
Bluegrass, Rye Grass, Fescue, Bent and similar cool climate grasses	3 - 4 oz. in 3 - 6 gallons water	1. Apply at first sign of disease. 2. Repeat every 7 - 10 days during warm damp weather. 3. For control of existing infections, apply at higher rates as soon as symptoms appear.
Helminthosporium Leaf Spot* (Helminthosporium spp.) Dollar Spot* (Sclerotinia homoeocarpa)	7 - 10 oz. in 5 - 10 gallons water	1. Preventative applications should be in spring or fall of each year. 2. For control of existing infections, apply at higher rate as soon as symptoms appear. 3. If disease is severe or reappears, area should be retreated 3 - 4 weeks later.
Gray Snow Mold (Typhula spp.) Pink Snow Mold (Fusarium nivale)	8 oz. in 5 - 10 gallons water	1. Make one application prior to first snowfall.

*DO NOT USE IN CALIFORNIA

Treated area should be watered lightly as soon as possible following treatment to wash chemical down to the soil level.
Do not graze treated areas or feed clippings to livestock.

Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

ORNAMENTALS

TERRACLOR 75%WP can be used as soil treatment for protection against a variety of soil borne diseases in nursery, greenhouse and field grown ornamentals.

It is important that TERRACLOR 75%WP be placed or mixed in the soil for best results.

See use and rate recommendations for specific plant and disease recommendations.

When rate range is shown, the lower rate should be used on lighter soils or when disease pressure is expected to be low. The higher rates should be used on heavy soils in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

DISEASES CONTROLLED

Root and Stem Rot
Storage Rot
Neck Dry Rot
Bulb Dry Rot
Needle Blight
Petiole Blight
Bulb and Stem Rot
Flower Blight
Crown Rot
Black Rot

Fungi

Rhizoctonia spp.
Botrytis cinerea
Stromatinia gladioli
Sclerotium gladioli
Dothistromia pini
Ovulinia azaleae
Sclerotium rolfsii
Sclerotium cameliae
Peltularia rolfsii
Sclerotinia sclerotiorum

PLANT TOLERANCE With specific plants not listed below we recommend a preliminary trial to determine plant tolerance.

USE AND RATE RECOMMENDATION:

TERRACLOR 75%WP as Soil Drench at Seeding or Transplanting in Nursery and Greenhouse Grown Ornamentals

<p>Foliage Plants</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;">Aglaonema</td> <td style="width: 33%;">Ficus</td> <td style="width: 33%;">Pothos</td> </tr> <tr> <td>Ajuga</td> <td>Fittonia</td> <td>Schefflera</td> </tr> <tr> <td>Anthurium</td> <td>Gynura</td> <td>Schubertgera</td> </tr> <tr> <td>Aphelandra</td> <td>Hoya</td> <td>Spathiphyllum</td> </tr> <tr> <td>Caladium</td> <td>Maranta</td> <td>Syngonium</td> </tr> <tr> <td>Cissus</td> <td>Palms</td> <td>Tradescantia</td> </tr> <tr> <td>Differbachia</td> <td>Peperomia</td> <td>Similar foliage plants</td> </tr> <tr> <td>Dracena</td> <td>Pilea</td> <td></td> </tr> </table> <p>Do not apply to Philodendron or Pilea or injury may occur.</p>	Aglaonema	Ficus	Pothos	Ajuga	Fittonia	Schefflera	Anthurium	Gynura	Schubertgera	Aphelandra	Hoya	Spathiphyllum	Caladium	Maranta	Syngonium	Cissus	Palms	Tradescantia	Differbachia	Peperomia	Similar foliage plants	Dracena	Pilea		<p>For control of Rhizoctonia diseases. Application Rates: Use 4 to 8 oz. in 100 gallons of water and apply to 400 - 800 sq. ft. of flat, bench or bed area.</p> <p>Coverage: For soil depths of 4 inches or less, apply 100 gallons to 800 sq. ft. For soil depths of more than 4 inches, apply 100 gallons to 400 sq. ft.</p> <p>One repeat application can be made 4 to 6 weeks later, if necessary.</p>												
Aglaonema	Ficus	Pothos																																			
Ajuga	Fittonia	Schefflera																																			
Anthurium	Gynura	Schubertgera																																			
Aphelandra	Hoya	Spathiphyllum																																			
Caladium	Maranta	Syngonium																																			
Cissus	Palms	Tradescantia																																			
Differbachia	Peperomia	Similar foliage plants																																			
Dracena	Pilea																																				
<p>Bedding and Flowering Plants</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;">Abutilon</td> <td style="width: 33%;">Dianthus</td> <td style="width: 33%;">Petunia</td> </tr> <tr> <td>Alyssum</td> <td>Foxglove</td> <td>Pointsettia</td> </tr> <tr> <td>Calceolaria</td> <td>Fuchsia</td> <td>Portulaca</td> </tr> <tr> <td>Canna</td> <td>Gardenia</td> <td>Salvia</td> </tr> <tr> <td>Carnation</td> <td>Gazania</td> <td>Scabiosa</td> </tr> <tr> <td>Chrysanthemum</td> <td>Geranium</td> <td>Sedum</td> </tr> <tr> <td>Cineraria</td> <td>Gloxinia</td> <td>Sempervivum</td> </tr> <tr> <td>Columbine</td> <td>Impatiens</td> <td>Verbena</td> </tr> <tr> <td>Coral Bells</td> <td>Lily</td> <td>Vinca</td> </tr> <tr> <td>Cyclamen</td> <td>Lupine</td> <td>Zinnia</td> </tr> <tr> <td>Dahlia</td> <td>Marigold</td> <td>Similar plants</td> </tr> <tr> <td>Delphinium</td> <td>Monarda</td> <td></td> </tr> </table>	Abutilon	Dianthus	Petunia	Alyssum	Foxglove	Pointsettia	Calceolaria	Fuchsia	Portulaca	Canna	Gardenia	Salvia	Carnation	Gazania	Scabiosa	Chrysanthemum	Geranium	Sedum	Cineraria	Gloxinia	Sempervivum	Columbine	Impatiens	Verbena	Coral Bells	Lily	Vinca	Cyclamen	Lupine	Zinnia	Dahlia	Marigold	Similar plants	Delphinium	Monarda		
Abutilon	Dianthus	Petunia																																			
Alyssum	Foxglove	Pointsettia																																			
Calceolaria	Fuchsia	Portulaca																																			
Canna	Gardenia	Salvia																																			
Carnation	Gazania	Scabiosa																																			
Chrysanthemum	Geranium	Sedum																																			
Cineraria	Gloxinia	Sempervivum																																			
Columbine	Impatiens	Verbena																																			
Coral Bells	Lily	Vinca																																			
Cyclamen	Lupine	Zinnia																																			
Dahlia	Marigold	Similar plants																																			
Delphinium	Monarda																																				
<p>Woody Ornamentals</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;">Arbutus</td> <td style="width: 33%;">English Ivy</td> <td style="width: 33%;">Podocarpus</td> </tr> <tr> <td>Aucuba</td> <td>Holly</td> <td>Privet</td> </tr> <tr> <td>Azalea</td> <td>Juniper</td> <td>Rhododendron</td> </tr> <tr> <td>Baccharis</td> <td>Maple</td> <td>Viburnum</td> </tr> <tr> <td>Carissa</td> <td>Pine</td> <td>Similar woody plants</td> </tr> </table>	Arbutus	English Ivy	Podocarpus	Aucuba	Holly	Privet	Azalea	Juniper	Rhododendron	Baccharis	Maple	Viburnum	Carissa	Pine	Similar woody plants																						
Arbutus	English Ivy	Podocarpus																																			
Aucuba	Holly	Privet																																			
Azalea	Juniper	Rhododendron																																			
Baccharis	Maple	Viburnum																																			
Carissa	Pine	Similar woody plants																																			

Solutions of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application.

USE AND RATE RECOMMENDATIONS:

TERRACLOR 75%WP as a Broadcast Incorporation Prior to Planting Field Grown Crops

Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 sq. ft. area)	Application
Easter Lilies	Root Rot (Rhizoctonia spp.)	3 1/4 to 6 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 6 - 7 inches and see note below.
Iris Hyacinth Narcissus Tulips	Crown Rot (Peltularia rolfsii) Black Rot (Sclerotinia sclerotiorum)	Use lower rates for light soils and higher rates for heavy soils.	
Dutch Iris	Bulb Rot Stem Rot (Sclerotium rolfsii)	3 1/4 to 4 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 2 - 3 inches.
Gladiolus	Neck or Bulb Dry Rot (Stromatinia or Sclerotinia gladioli)	3 3/4	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 4 - 6 inches. Bulbs may be planted immediately after incorporation.
Glendula Larkspur Snapdragon Sweet Pea	Stem Rot Root R. (Rhizoctonia spp.)	3 1/2	Apply specified rate in sufficient water to insure uniform ground coverage prior to planting and thoroughly incorporate to a depth of 4 inches.
Southern Pine Seedlings	Root/Stem Rot (Rhizoctonia spp.) Needle Blight (Dothistromia pini)	1 1/2	Apply specified rate in sufficient water to insure uniform ground coverage. An additional 1 1/2 inches of water per acre should be applied by irrigation prior to planting. Seed may be planted immediately after irrigation or within one week of this date.

Solution of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application of the fungicide.

TERRACLOR 75%WP as a Ground Spray on Established Woody Ornamentals

Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 150 sq. ft. area)	Application
Azaleas	Azalea petal blight (Ovulinia Azaleae)	1	Apply specified rate per 150 sq. ft. in sufficient water to assure uniform ground coverage. Apply to the ground beneath the bushes and immediate surrounding area. Begin application prior to opening of buds and repeat every 3 - 4 weeks during bloom season. Solution of TERRACLOR 75%WP and water should be continuously agitated to assure uniform application of fungicide.
Camelias	Camellia flower blight (Sclerotinia camelliae)	1	

TERRACLOR 75%WP as a Dip Application

Crop	Disease	Rate of TERRACLOR 75%WP (Lbs.)	Application
Iris Hyacinth Narcissus Tulips	Crown Rot (Pellicularia rolfsii) Black Rot (Sclerotinia sclerotiorum)	2	Dip bulbs for 5 minutes in 7.5% concentration which is obtained by adding 2 lbs. of TERRACLOR 75%WP to 3.2 gallons of water. Add 1% sticker to solution and maintain good agitation.
Easter Lilies	Root Rot (Rhizoctonia spp.)	4 - 6	Mix TERRACLOR 75%WP in 100 gallons of water and dip bulbs for 5 - 15 minutes. A sticking agent may be added. Maintain good agitation in dip tanks.
Roses	Storage Rot (Botrytis cinerea)	2	Mix TERRACLOR 75%WP in 100 gallons of water and dip dormant roses prior to storage. A spray application of the same solution may be substituted for dipping. Maintain good agitation.

TERRACLOR 75%WP as Band Application Prior to Planting Field Grown Crops

Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 linear ft. of row)	Application
Gladiolus	Neck or Bulb Dry Rot (Stomatium or Sclerotinia gladioli)	5 1/2	Apply as a 12 inch band centered on the row at planting time. Incorporate to depth of 4 inches. Bulb may be planted immediately after incorporation.
Dutch Iris	Bulb Rot Stem Rot (Sclerotium rolfsii)	1/2	Apply as a 12 inch band centered on the row at planting time. Follow with 3 additional applications at the same rate at 15 day intervals.

TERRACLOR 75%WP as In Furrow Application at Planting

Crop	Disease	Rate of TERRACLOR 75%WP (Lbs. per 1,000 linear ft. of row)	Application
Easter Lilies	Root Rot (Rhizotonia spp.)	4 1/2 to 6	Apply in 5 - 10 gallons of water. Direct spray into open furrow to thoroughly cover entire furrow and bulbs. Close furrow in normal manner. See notes below. Use lower rates for lighter soils and higher rates for heavier soils. For Black rot control, use high rates. Solution of TERRACLOR 75%WP and water should be agitated to assure uniform application of the fungicide.
Iris Hyacinth Narcissus Tulips	Crown Rot (Pellicularia rolfsii) Black Rot (Sclerotinia sclerotiorum)		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Paper Bags: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Cardboard Canisters: (Commercial) Completely empty canisters into application equipment. Then dispose of container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Household: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.