

WASG  
209477  
301  
18

JAN 18 1988

Mr. Raymond A. Cardona  
Uniroyal Chemical Company  
74 Amity Road  
Bethany, CT 06525

Dear Mr. Cardona:

Subject: Terraclor 2 lb Emulsifiable  
EPA Registration No. 400-400  
Your Letter Dated December 3, 1987

The proposed label, submitted October 28, 1987, is acceptable as long as it is understood that you must comply with the settlement agreement in paragraph II-1 on page 3. That is, the references to potatoes must be obscured on labels until such time as the use is reinstated.

Sincerely yours,

Lois A. Rossi  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (TS-767C)

**BEST AVAILABLE COPY**

10023-I-DeLuise-R-12-KENCO:1 12/31/87 12/31/87:CB:jh

CONSEQUENCES

SYMBOL	ORIGINATOR						
SURNAME							
DATE							

2064

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS  
CAUTION**

This material may cause skin irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Use with adequate ventilation. Do not contaminate feed or foodstuffs. The following protective clothing are required during mixing/loading operations: respirator, gloves, long-sleeve shirt, long pants, socks and shoes.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water or wetlands by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable. Do not use or store near heat sources or open flame. Do not use in the vicinity of pilot lights.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Protect from temperatures below 20° F.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

# TERRACLOR<sup>®</sup>

## 2 LB. EMULSIFIABLE SOIL FUNGICIDE

**UNIROYAL  
CHEMICAL**

**ACCEPTED  
COMMENTS**  
EPA Letter Dated:

JAN 19 1988

Under the Federal Insecticide, Fungicide, and Herbicide Act as amended, for products registered under FIFRA Act No. 400-400

COMPOSITION	
Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB) ...	23.8%
Inert ingredients: .....	76.2%
Total: .....	100.0%

Contains 2 lbs. of Pentachloronitrobenzene per U.S. Gallon at 68°F.

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush with plenty of water. Get medical attention.  
**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.  
**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

Uniroyal Chemical Company, Inc.  
Middlebury, CT 06749

8-30-87

EPA REG. NO. 400-400  
EPA EST. NO.

Terraclor is a Registered Trademark of Uniroyal, Inc.  
©Copyright 1987, Uniroyal, Inc.

Lot. No.

NET CONTENTS:

NOV 2 1987

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

Crops (diseases)	Rates of TERRACLOR 2 lb. Emulsifiable		Application	Remarks
	gals/acre	fl. oz/1000 Feet of Row		
DRY, SUCCULENT & SNAP BEANS (Root and Stem Rot - Rhizoctonia solani)	1/2 to 3/4 gallons Use in 8 to 10 gals water (36 inch row spacing - bush type beans; 62 inch row spacing - pole beans) In California and Arizona: 3 to 4 pts. in 8 to 10 gals water (spacings as above)	Bush: 4.4 to 6.6 Pole: 7.6 to 11.3  Bush: 2.2 to 4.4 Pole: 3.8 to 5.6	Spray planting furrow and covering soil at planting time	Avoid applying directly to seed; otherwise delayed emergence may occur. Use lower rate on lighter soils.
(White Mold - Sclerotinia sclerotiorum)	1 gallon Use in 8 to 10 gals water (36 inch row spacing - bush type beans; 62 inch row spacing - pole beans)	Bush: 8.8 Pole: 15	Spray an 8 inch wide band centered on the row immediately after or at time of seeding. If disease is severe, application may be repeated at 2 to 3 week intervals using 1 or 2 nozzles per row and directing spray at base of plants.	Remarks as above. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

Preharvest Intervals: Dry beans - Do not apply within 45 days of harvest; Succulent beans - Do not apply within 35 days of harvest; Snap beans - Do not apply within 14 days of harvest.

COTTON (Damping off - Rhizoctonia solani)	1/2 to 1 gallon Use in 15 gals water (40 inch row spacing)	4.8 to 9.7	Spray on seed and surrounding soil at planting time.	Do not allow the feeding or grazing of cotton foliage by livestock.
GARLIC (White Rot - Sclerotium cepivorum)	10 gallons Use in 100 gals water (24 inch row spacing)	58.7	Apply at planting Attach a suitable rig to the machine planter and spray 8-12 inch bands using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.	

Crops (diseases)	Rates of TERRACLOR 2 lb. Emulsifiable		Application	Remarks
	gals/acre	fl. oz/1000 Feet of Row		
POTATOES (Rhizoctonia solani)	9 to 12-1/2 gallons Use in sufficient water to total 25 gals final spray volume	—	Apply as a broad- cast to the soil surface prior to planting and thoroughly mix (disc and cross disc) to a depth of 4 inches. A drag harrow is usually attached behind the discs on the last discing.	It is very important to thoroughly mix the TERRACLOR into the tuber-forming zone of the soil. It is desirable to make treatment when the soil is slightly moist.
	5 gallons Use in sufficient water to total 20 gallons final spray volume (42 inch row spacing)	51.6	Direct the spray solution to several points in the row to insure optimum treat- ment of the stem, stolen and tuber- forming zones, using three nozzles per row. Place one fan nozzle ahead of the furrow opening discs and two cone spray nozzles above and slightly ahead of the closing discs. Select nozzles that will deliver approximately 1/3 of the recommended spray volume per acre through each nozzle at the desired planting speed and pressure.	

Rotation Restriction: Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.