


400-402

PM 21

10.7



# TERRACLOR<sup>®</sup>

10% GRANULAR  
SOIL FUNGICIDE

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

### CAUTION

Harmful if swallowed. Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. The following protective clothing are required during mixing/loading operations: gloves, a respirator, long-sleeved shirt, long pants, socks and shoes.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover, collect, or incorporate granules. Do not contaminate water by cleaning of equipment or disposal of wastes.

**UNIROYAL  
CHEMICAL**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB)	10%
Inert Ingredients	90%
Total	100%

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a dry location

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush eyes with plenty of water. Get medical attention.

**IF ON SKIN:** Wash with plenty of water. Get medical attention.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

UNIROYAL CHEMICAL COMPANY, INC.

Middlebury, CT 06749

EPA REG. NO. 400-402

EPA EST. NO.

10-11-88

Lot. No.

Terraclor is a Registered Trademark of Uniroyal Chemical Company, Inc.

©Copyright 1987, Uniroyal Chemical Company, Inc.

NET CONTENTS:

**RATE OF TERRACLOR 10% GRANULAR**

CROP	DISEASE	POUNDS/ACRE	POUNDS/1000 FEET OF ROW	RECOMMENDED APPLICATION
COTTON	Damping-off Seed and Seedling Rots <i>Rhizoctonia solani</i>	10-20 (40 inch row spacings)	$\frac{3}{4}$ to 1 $\frac{1}{2}$	In-furrow Application. Apply over seed and covering soil at planting time, using suitable application equipment. Do not allow feeding or grazing of cotton foliage by livestock.
BEANS	Root and Stem Rot - <i>Rhizoctonia solani</i>	10 to 15 (36 inch row spacing - bush type 62 inch row spacing - pole beans)	Bush $\frac{3}{4}$ to 1 Pole 1 $\frac{1}{2}$ to 1 $\frac{3}{4}$	In-furrow Application. Direct the material into the planting furrow and covering soil at planting time. NOTE: Do not feed treated vines to livestock.
CABBAGE, BRUSSELS SPROUTS, BROCCOLI	Clubroot - <i>Plasmodiophora brassicae</i>	200 (40 inch row spacing)	15	Row Application. Apply in a 12 to 15 inch band, by means of a suitable applicator, and rototill to a depth of 4 to 6 inches, immediately prior to planting.
		300	—	Broadcast Application. Apply to the soil surface prior to planting. Disc and cross disc to a depth of 4 to 6 inches.
For control of clubroot, it is very important to thoroughly mix TERRACLOR with the soil.				
CAULIFLOWER	Clubroot - <i>Plasmodiophora brassicae</i>	200 (48 inch row spacing)	18.3	Row Application. Apply in a 12 to 15 inch band, by means of a suitable applicator, and rototill to a depth of 4 to 6 inches, immediately prior to planting. It is very important to thoroughly mix TERRACLOR with the soil.
		300	—	Broadcast Application. Apply to the soil surface prior to planting. Disc and cross disc to a depth of 4 to 6 inches.
	Wire Stem or Black Root <i>Corticium solani</i>	110 to 150 (48 inch row spacing)	10 to 13.7	Row Application. Apply in an 8 inch band centered on the row, immediately prior to seeding. For smaller areas apply 12 lbs. per 1,000 sq. ft. Rototill to a depth of 2 inches.
PEANUTS	Southern blight, <i>Sclerotium rolfsii</i>	36 inch row spacing		
		100	6.9	Surface Band Treatment. Apply in an 8 to 12 inch band, centered on the row, at pegging time.
		100	6.9	Pre-plant Soil Mix. Apply in an 8 to 12 inch band, centered on the row, and mix to a depth of 2 to 4 inches. Application may be made simultaneously with planting, using suitable equipment.
		33 per application	2.3 pounds per application	Soil Mix Cultivation. Apply in split applications during three cultivations. Attach two delivery tubes per row ahead of inside cultivator sweeps (one on each side) and adjust so as to treat a total band of 12 inches wide, centered on the row. Do not pile untreated soil on the crown of the plant.
		50	3.4 pounds per application	Split Pegging Applications. Apply as a 12 to 18 inch band centered over the row, follow by shallow incorporation if possible. The initial application should be timed to coincide with early pegging. A second application at the same rate may be made if disease symptoms reappear. Depending on weather, this usually occurs 4 to 5 weeks after the first application.
NOTE: Do not exceed 10 lbs. actual PCNB per acre in any one season, regardless of the application methods used. Do not feed treated vines to livestock. Do not allow hogging down of peanuts.				
POTATOES	<i>Rhizoctonia solani</i>	180 to 250	—	Broadcast Application. Spread evenly over the soil surface prior to planting. Disc and cross disc to depth of 4 to 6 inches.
		100 (36 inch row spacing)	6.8	Row Application. Apply in-furrow and thoroughly work into top 4 to 6 inches of the soil. It is essential that the fungicide is worked into the tuber forming zone of the soil. It is desirable to make treatment when the soil is slightly moist.
Rotation Restriction: Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.				

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.