

TERRACLOR®

10% GRANULAR SOIL FUNGICIDE

ACCEPTED

FEB 13 1990

Use with Federal, State, and
 Regulatory Agency approval for
 pesticide registered under
 EPA Reg. No. 400-402

**UNIROYAL
CHEMICAL**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover, collect, or incorporate granules. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB)	10%
Inert Ingredients:	90%
Total:	100%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention.
IF ON SKIN: Wash with plenty of water. Get medical attention.
IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

UNIROYAL CHEMICAL COMPANY, INC.
 Middlebury, CT 06749
 EPA REG. NO. 400-402
 EPA EST. NO.

Terraclor is a Registered Trademark of Uniroyal Chemical Company, Inc.
 ©Copyright 1987, Uniroyal Chemical Company, Inc.

NET CONTENTS:

Lot. No.

DIRECTIONS FOR USE

When a rate range is shown the lower rate should be used on lighter soils or when disease pressure is expected to be low. The higher rates should be used on heavy soils, in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

RATE OF TERRACLOR 10% GRANULAR

CROP	DISEASE	RATE OF TERRACLOR 10% GRANULAR		RECOMMENDED APPLICATION
		POUNDS/ACRE	POUNDS/1000 FEET OF ROW	
COTTON	Damping-off, Seed and Seedling Rots Rhizoctonia solani	10-20 (40 inch row spacings)	¾ to 1½	In-furrow Application: Apply over seed and covering soil at planting time, using suitable application equipment. Do not allow feeding or grazing of cotton foliage by livestock.
BEANS	Root and Stem Rot - Rhizoctonia solani	10 to 15 (36 inch row spacing)	¾ to 1	In-furrow Application: Direct the material into the planting furrow and covering soil at planting time. NOTE: Do not feed treated vines to livestock.
CABBAGE, BRUSSELS SPROUTS, BROCCOLI, CAULIFLOWER, COLLARDS*, KALE* AND MUSTARD GREENS*	Clubroot - Plasmodiophora brassicae	200 (40 inch row spacing)	15	Row Application: Apply in a 12 to 15 inch band, by means of a suitable applicator, and rototill to a depth of 4 to 6 inches, immediately prior to planting. May be used on cole crops which are direct seeded.
		300	—	Broadcast Application: Apply to the soil surface prior to planting. Disc and cross disc to a depth of 4 to 6 inches. May be used on cole crops which are direct seeded.
	For control of clubroot, it is very important to thoroughly mix TERRACLOR with the soil.			
	Wire Stem or Black Root Corticium solani	110 to 150 (40 inch row spacing)	8.5 to 11.5	Row Application: Apply in an 8 inch band centered on the row, immediately prior to seeding. For smaller areas apply 12 lbs. per 1,000 sq. ft. Rototill to a depth of 2 inches.
*Based on the available residue data, the use of Terraclor on collards, kale and mustard greens is restricted to the state of Georgia.				
PEANUTS	Root and Stem Rot Rhizoctonia solani* Southern blight Sclerotium rolfsii	10 to 20 (36 inch row spacing)	0.7 to 1.4	In Furrow Application: Apply in furrow at planting using suitable equipment.
		100 (36 inch row spacing)	6.9	Surface Band Treatment: Apply in an 8 to 12 inch band, centered on the row, at pegging time.
		50 (36 inch row spacing)	3.4	Narrow Band Application: Apply in a narrow band so there is a 4-6 inch band of granules on the soil surface when the peanut canopy completely shades the soil (50-70 days). For best results, use the drop tube with no bander, and run it 1-2 inches into the peanut foliage; or run some type of drag over the top to sweep the foliage forward in front of the drop tube during application.
		33 per application	2.3 pounds per application	Soil Mix Cultivation: Apply in split applications during three cultivations. Attach two delivery tubes per row ahead of inside cultivator sweeps (one on each side) and adjust so as to treat a total band of 12 inches wide, centered on the row. Do not pile untreated soil on the crown of the plant.
		25-50 per application	1.7 to 3.4 pounds per application	Split Pegging Applications: Apply at 25 lbs. per acre as a 7 inch band or 50 lbs. per acre as a 12 to 18 inch band centered over the row follow by shallow incorporation if possible. The initial application should be timed to coincide with early pegging. A second application at the same rate may be made if disease symptoms reappear.
NOTE: Do not exceed 10 lbs. actual PCNB per acre in any one season, regardless of the application methods used. Do not feed treated vines to livestock. Do not allow hogging down of peanuts. *DO NOT USE IN CALIFORNIA				
POTATOES	Rhizoctonia solani	180 to 250	—	Broadcast Application: Spread evenly over the soil surface prior to planting. Disc and cross disc to depth of 4 to 6 inches.
		100 (36 inch row spacing)	6.8	Row Application: Apply in-furrow and thoroughly work into top 4 to 6 inches of the soil. It is essential that the fungicide is worked into the tuber forming zone of the soil. It is desirable to make treatment when the soil is slightly moist.
	Scab - Streptomyces scabies	Use Rates shown above	Use Rates shown above	Treat as above using broadcast or in-furrow application. For broadcast application material must be thoroughly incorporated to a depth of 8-10 inches by rototilling or cross disking for effective control. Under high moisture soil conditions, 80% or higher available moisture, use lower rate. Under dry conditions use higher rate. AVAILABLE SOIL MOISTURE IN THE HILL SHOULD BE MAINTAINED ABOVE 65% FOLLOWING TUBER INITIATION. DO NOT USE IN CALIFORNIA
Rotation Restriction: Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.				

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.