

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

400-405

124-

Willard F. Cummings
U.S. Registration Manager
Crop Protection Research & Development
Uniroyal Chemical Company, Inc.
74 Amity Road
Bethany, Connecticut 06525

MAY 21 1992

Subject: Terraclor Super 2E
EPA Reg. No. 400-405
RE: Label Amendment (Rate limitations)
Your Letter Dated February 22, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

- * Please delete the feeding restriction on cotton gin waste. Gin waste is not currently considered a significant feed item and since there are no established tolerances for any pesticide in or on gin waste, a tolerance is not required at this time. However, the Agency is revising its guidelines and tolerances/data for residues of PCNB in or on gin waste may be required at a future date. Please see attached review for additional comments.

Sincerely yours,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C						
SURNAME	M.../11						
DATE	5/20/92						

**EMULSIFIABLE
SOIL FUNGICIDE**
Controls Rhizoctonia, Pythium and Fusarium
Soil-Borne Seedling Diseases of Cotton

ACCEPTED
with COMMENTS
in EPA Letter Datab

BEST AVAILABLE COPY



Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
400-405

COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB)	23.2%
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole	5.8%
Inert Ingredients:	71.0%
TOTAL	100.0%

Contains 2 lbs. of Pentachloronitrobenzene and
0.5 lbs. of 5-Ethoxy-3-trichloromethyl-1,2,4-thiadiazole per U.S. gallon.

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, Connecticut 06749

EPA REG. NO. 400-405

EPA EST. NO.

TERRACLOR SUPER X® is a Registered Trademark of Uniroyal Chemical Company, Inc.

©Copyright 1991, Uniroyal Chemical Company Inc.

001

Net Contents:

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink an eight ounce glass of water containing 2 tablespoons of activated charcoal.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING

Harmful if swallowed. Do not inhale spray mist. Do not get on skin or in eyes. May cause skin irritation. Skin contact may be harmful. The following protective clothing are required during mixing/loading operations: respirator, gloves, long-sleeved shirt, long pants, socks and shoes. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Wash thoroughly all contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Do not use or store near heat sources or open flame. Do not use in the vicinity of pilot lights.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

COTTON

(Seedling Disease Complex - Rhizoctonia, Pythium and Fusarium)

SOIL IN-FURROW APPLICATION: At planting time, use 2 to 3 quarts per acre (per 13,000 linear feet based on 40 in. rows) mixed with 5 to 15 gallons of water. Use the higher treatment rate under conditions of expected heavy disease pressure. Spray all the soil that surrounds and covers the seed. Two nozzles work best; center the front nozzle over the

open furrow to spray soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow.

Do not allow the feeding or grazing of cotton foliage by livestock.

Do not feed gin waste to livestock.

Apply only at planting time.

Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.

ROTATION RESTRICTION: Do not plant root crops in PCNB treated fields within 12 months of application unless PCNB is registered for use on those crops.

PESTICIDE DISPOSAL - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.