TERRACLOR Super X°

GRANULAR SOIL FUNGICIDE FOF! COTTON



BEST AVAILABLE COPT

#### COMPOSITION

Active ingred:ant: (% by weight) Pentachloronitrobenzene (PCNB)	10.0%
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole	2.5%
Inert Incredients:	87.5%
TOTAL	100.0%
UNIROYAL CHEMICAL COMPANY, INC., Middlebury, Connecticut 06/49	
EPA REG. NO. 400-406	003

TERRACLO9 SUPER X® is a Registered Trademark of Uniroyal Chemical Company, Inc. Copyright 1991, Uniroyal Chemical Company Inc.

# KEEP OUT OF REACH OF CHILDREN WARNING

## STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Wash with plenty of water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink an eight conce glass of water containing 2 tablespoons of activated

### PRECAUTIONARY STATEMENTS HAZAROS TO HUMANS

Harmful if swallowed. Do not inhale dust. Do not get on skin or in eyes. May cause skin irritation. Contact may be harmful. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Wash thoroughly all contaminated clothing before reuse.

**ENVIRONMENTAL HAZAROS** 

This posticide is toxic to fish and aquatic organisms. Cover, collect or incorporate treated granule. Do not contaminate water when disposing of equipment washwaters.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its izbeling.

#### COTTON

(Seedling Disease Complex - Rhizoctonia, Pythium and Fusarium) **SOIL IN-FURROW APPLICATION:** At planting came, use 8 to 12 tbs. per acre (per 13,000 linear feet based on 40 in, rsws).

Apply the specified dosage to the soil around the seed and to the covering soil as it fills in the furrow. The soil around the seed and the covering soil should be thoroughly mixed with the product. Two nozzles work best; center the front nozzle over the open furrow to spray soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow. The higher rates should be used when weather conditions are expected to be untavorable for rapid germination, fields have a history of disease problems or minimum or low till programs are utilized. Apply only at planting time.

Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.

Do not allow the feeding or grazing of cotton foliage by livestock.

ROTATION RESTRICTION: Do not plant root crops in PCNE treated fields within 12 months of application unless PCNB is registered for use on those crops.

### STORAGE AND DISPOSAL

De not conteminate water, food or food by storage or disposal. **STORAGE** - Store in a dry location.

PESTICIOE DISPOSAL - Pesticide wastes are toxic, Improper disposal of excess publicide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Prejirodimental

Control Agency, or the Hazardous Wasti feptes intative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Completely, empty, bag info application equipment. Then dispose of empty bag in a fanitary landfill or by incineration, or if allowed by State and local authorities; by burning. If burned, stay out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes sucted on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warrant/finer any other warranty of merchantability or litheas for a perticular furpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably to reasonable to seller, and buyer assumes the rick of any such

# STORAGE AND DISPOSAL

De not contaminate water, feed or feed by storage or disposal.

STORAGE - Protect from temperatures below 29°F.
PESTICIDE DISPOSAL - Pasticide westes are toxic. Improper disposal of excess pesticide, spray mixture, or rineste is a violation of Federal Law, If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is researably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this variety der any either warranty of merchanishilly or fitness for a perfector purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably forecombile to soller, and buyer assumes the risks of any such use.

TERRACLOR SUPER XO is a Registered Trademork of Universal

Chemical Company, Inc.

**CCopyright 1991, Untrayel Chemical Company Inc.** 

