TERRACLOR Super X GRANULAR SOIL FUNGICIDE FOR COTTON



COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB)	
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole	
Inert Ingredients:	
TOTAL	
UNIROYAL CHEMICAL COMPANY, INC., Middlebury, Connectic	ut 06749
EPA REG. NO. 400-406	004
EPA EST, NO.	5-5-93
TERRACLOR SUPER X® is a Registered Trademark of Uniroyal Chemical (Company, Inc.
Copyright 1991, Uniroyal Chemical Company Inc.	

Net Contents:

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Wash with plenty of water. Get medical attention. IF SWALLOWED: Call a physician or Poison Control Center. Drink an eight ounce glass of water containing 2 tablespoons of activated charcoal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS WARNING

Harmful if swallowed. Do not inhale dust. Do not get on skin or in eyes. May cause skin irritation. Contact may be harmful. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes. Wash thoroughly all exposed skin. In case of eye contact, flush with plenty of water. Wash thoroughly all contaminated clothing before reuse.

ENVIRONMENTÁL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Cover, collect or incorporate treated granules. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

JUN 1 5 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act,

as amended, for the pesticide

400-406

registered under

EPA Reg. No.

Í

ſ

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

COTTON

(Seedling Disease Complex - Rhizoctonia, Pythium and Fusarium) SOIL IN-FURROW APPLICATION: At planting time, use 8 to 12 lbs. per acre (per 13,000 linear feet based on 40 in. rows).

Apply the specified dosage to the soil around the seed and to the covering soil as it fills in the furrow. The soil around the seed and the covering soil should be thoroughly mixed with the product. The higher rates should be used when weather conditions are expected to be unfavorable for rapid germination, fields have a history of disease problems or minimum or low till programs are utilized.

Apply only at planting time.

Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.

Do not allow the feeding or grazing of cotton foliage by livestock.

ROTATION RESTRICTION: Do not plant root crops in PCNB treated fields within 12 months of application unless PCNB is registered for use on those crops.

STORAGE AND DISPOSAL

1-1-1

Do not contaminate water, food or feed by storage or disposal. STORAGE - Store in a dry location.

PESTICIDE DISPOSAL - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.