

TURFCIDE®

10% GRANULAR SOIL FUNGICIDE

CONTROLS CERTAIN SOIL-BORNE
DISEASES OF LAWNS — TURF — ORNAMENTALS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. This material may cause skin irritation. Avoid contact with eyes, skin or clothing. Avoid contact with the skin by wearing the following protective clothing: long-sleeve shirt, long pants, socks and shoes. Wash hands thoroughly after using and prior to eating, drinking or smoking. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply product in areas which would permit runoff to contaminate water supplies. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL


Do not contaminate water, food or feed by storage or disposal.

STORAGE — Store in a dry location.

PESTICIDE DISPOSAL — Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL — Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

HOUSEHOLD: Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.



COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene	10.0%
Inert Ingredients:	90.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting.
IF ON SKIN: Remove contaminated clothing and wash skin with soap and warm water.
IF INHALED: Remove to fresh air.
IF IN EYES: Flush with water for 15 minutes.
For further treatment, get medical attention.
SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

CHEMICAL GROUP — Uniroyal, Inc.
Middlebury, CT 06749
EPA REG. NO. 400-407
EPA EST. NO.

Lot. No.

4-22-85

TURFCIDE is a Registered Trademark of Uniroyal, Inc.
©Copyright 1983, Uniroyal, Inc.
NET WEIGHT:

DIRECTIONS FOR USE

LAWNS AND TURF

Brown patch - Rhizoctonia solani: For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the fall or the spring at first indication of infection. Apply 7½ lbs. per 1,000 sq. ft. Following application, the treated areas should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later. For Dichondra: Treat as above, using 7½ lbs. per 1,000 sq. ft. For Northern or cool season lawns such as Rye Grass, Fescue, Bluegrass, Bent or similar grasses: Apply 2 to 2½ lbs. per 1,000 sq. ft. at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. Water treated area lightly to wash chemical down to soil. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.

Helminthosporium leaf spot: Apply at rate of 5 to 7½ lbs. per 1,000 sq. ft. Preventative applications should be made in spring or fall of each year. For control of existing infections, apply at the higher rate as soon as symptoms are first noted. Following applications, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3 to 4 weeks later.

Snow mold (Typhula incarnata, Fusarium nivale): Apply at rate of 5 to 7½ lbs. per 1,000 sq. ft. using a granular lawn spreader or comparable equipment. Application should be made prior to first snowfall.

Do not graze treated areas or feed clippings to livestock.

SPREADER SETTING NUMBER PER 1000 SQ. FT.

SPREADER MAKE	2 Lbs.	2½ Lbs.	5 Lbs.	7½ Lbs.
CYCLONE	3½	3¾	6	7½
SCOTT	5⅞	5¾	6½	7½
SEARS	6¼	6½	7½	9

These settings apply to 20/40 mesh granular carriers. The spreader settings listed above are approximate. Accurately weigh a few pounds into your spreader and apply to measured area. Then adjust spreader setting, if necessary, in order to assure correct rate of application

ORNAMENTALS

AFRICAN VIOLETS, BEGONIAS, CARNATIONS, CHRYSANTHEMUMS, POINSETTIAS, SNAPDRAGONS, and similar bedding plants (Stem Rot-Rhizoctonia): Apply 15 lbs. per 1,000 sq. ft. by spreading evenly on the soil surface and thoroughly mixing into the top 2 inches of soil. Apply to a well prepared seed bed.

BULBOUS IRIS, HYACINTH, NARCISSUS, AND TULIP (Crown Rot and Black Rot): Apply broadcast 1,000 to 2,000 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) and mix thoroughly in upper 6 to 7 inches of soil by double discing or rotovating. Note: Use higher rate for heavier soils and for Black Rot control.

CALENDULA, LARKSPUR, SNAPDRAGON, AND SWEET PEA (Stem Rot-Sclerotinia): Apply 1,000 lbs. per acre broadcast one week prior to planting. For smaller areas apply 2 lbs. per 100 sq. ft. Spread on the soil surface and uniformly mix in the soil to a depth of 4 inches. Terraclor was developed for this use in Charleston, South Carolina area.

DUTCH IRIS (Bulb and stem rot caused by Sclerotium rolfsii): Broadcast: Apply 1,000 to 1,500 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) prior to planting and mix with the top 2 or 3 inches of soil by discing or roto-tilling.

ROW: Apply 60 lbs. per 14,500 linear feet of row, in a 12 inch band, centered on the row, at planting time. Follow with three additional applications at this rate at 15 day intervals.

EASTER LILIES (Rhizoctonia root rot): Broadcast: Apply 1,000 to 2,000 lbs. per acre (2 to 4 lbs. per 100 sq. ft.) to soil surface prior to planting and mix thoroughly in upper 6 to 7 inches by double discing or rotovating. Note: Use the lower rate of application for light soils and higher rate of application for heavier soils.

GLADIOLUS (Neck or Bulb Dry Rot - Stromatinia or Sclerotinia gladioli): Broadcast: At planting time apply 1,200 lbs. per acre (1½ lbs. per 100 sq. ft.) to soil surface and mix thoroughly in upper 4 to 6 inches by double discing or rotovating. (Bulbs may be planted immediately.)

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

BEST AVAILABLE COPY