



# TERRACLOR<sup>®</sup>

## 30% GRANULAR SOIL FUNGICIDE

Controls Certain Soil-Borne Diseases of  
BEANS, CRUCIFERS, PEANUTS, POTATOES

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

#### CAUTION

Harmful if swallowed. Avoid inhaling dust. This material may cause skin irritation. Wash hands and face thoroughly after using. Wear gloves and a respirator when loading the spreader. The following protective clothing are required during mixing/loading operations: gloves, long-sleeved shirt, long pants, socks and shoes. Wash contaminated clothing before reuse.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a dry location  
**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACCEPTED  
MAY 30 1984  
UNIROYAL  
For the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the pesticide issued under Reg. No. 400-715

#### COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene	30%
Inert Ingredients	70%
Total	100.0

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering Syrup of Ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove victim to fresh air. Provide artificial respiration if needed. Get medical attention if needed.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

UNIROYAL CHEMICAL — Division of UNIROYAL, Inc.  
Naugatuck, Connecticut 06770  
EPA REG. NO. 400-415  
EPA EST. NO.

3 16-84

Lot. No.

TERRACLOR is a Registered Trademark of Uniroyal, Inc.  
Copyright 1983, Uniroyal, Inc.

NET CONTENTS:

#### DIRECTIONS FOR USE

**CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI (Clubroot - Plasmodiophora brassicae):** Row Application Apply 67 lbs per 13,100 feet of row for cabbage, broccoli and brussels sprouts or to 10,900 linear feet of row for cauliflower, in a 12 to 15 inch band, by means of a suitable applicator, and rototill to a depth of 4 to 6 inches, immediately prior to transplanting.

**BROADCAST.** Apply 100 lbs per acre to the soil surface prior to transplanting. Disc and cross disc to a depth of 4 to 6 inches.

**NOTE:** For clubroot of cabbage, cauliflower, broccoli and brussels sprouts, it is very important to thoroughly mix it with the soil.

**CAULIFLOWER (Wire Stem or Black root - Corticium solani):** Apply 37 to 50 lbs per 10,900 feet in an 8-inch band, centered on the row, immediately prior to seeding. For smaller areas apply 4 lbs per 1000 sq. ft. Rototill to a depth of 2 inches.

**BEANS (Root and Stem Rot - Rhizoctonia solani) FURROW TREATMENT:** Apply 4 to 5 lbs per 14,500 linear feet of row for bush beans, or per 8,430 linear feet of row for pole beans. Direct the material into the planting furrow and covering soil at planting time. Avoid getting the material concentrated around the seed otherwise delayed emergence may occur. **NOTE:** Do not feed treated vines to livestock.

**PEANUTS (Southern Stem and Root Rot - Sclerotium rolfsii): PRE- PLANT SOIL MIX:** Apply 34 lbs per 12,400 linear feet of row, in an 8 to 12 inch band, centered on the row, and mix to a depth of 2 to 4 inches. Application may be made simultaneously with planting, using suitable equipment.

**SURFACE BAND TREATMENT:** When pegging begins, apply 34 lbs per 12,400 linear feet of row, in an 8 to 12 inch band, directed at the plant crown and soil pegging zone.

**NOTE:** Do not use more than one application method during the same planting season. Do not feed treated vines to livestock.

**POTATOES (Rhizoctonia solani): BROADCAST TREATMENT:** Use 60 to 83 lbs per acre. Spread evenly over the soil surface prior to planting. Disc and cross disc to a depth of 4 to 6 inches.

**ROW (in-furrow) APPLICATION:** Apply 33 lbs per 12,400 linear feet of row and work thoroughly into the top 4 to 6 inches of the soil (it is essential that the fungicide is worked into the tuber forming zone of the soil). It is desirable to make treatment when the soil is slightly moist.

**NOTE:** Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

**IMPORTANT NOTICE —** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.