

ACCEPTED

AUG 4 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **400-417**

16F2
Pr. 21
400-417

TERRAZOLE®

25% EMULSIFIABLE

**For Commercial Use Only
Controls Certain Soil-Borne Diseases
of ORNAMENTALS**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
DANGER**

Corrosive, causes irreversible eye damage. Harmful if swallowed. Do not take internally. Do not inhale vapors or spray mist. Do not get in eyes, or on clothing. Wear goggles or face shield. May cause skin irritation. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.



COMPOSITION

Active Ingredient: (% by weight)	
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole*	25.0%
Inert Ingredients:**	75.0%
Total:	100.0%

*U.S. Patents 3,260,588, 3,260,725 and others.
**Contains Petroleum Distillates
(Contains 2 lbs. 5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole per U.S. Gallon at 68°F).

**KEEP OUT OF REACH OF CHILDREN
DANGER!**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Obtain prompt medical attention.

IF IN EYES: Flush with plenty of water. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove to fresh air.

NOTE TO ATTENDING PHYSICIAN: Probable mucosal damage and the presence of petroleum distillates in the formulation may contraindicate the use of gastric lavage.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

**Uniroyal Chemical Company, Inc.
Middlebury, CT 06749**

EPA REG. NO. 400-417
EPA EST. NO.

4-23-86

Terrazole® is a Registered Trademark of Uniroyal, Inc.
©Copyright 1983, Uniroyal, Inc.

NET CONTENTS:

GENERAL INFORMATION

Terrazole is useful as a soil fungicide for control of damping-off, root and stem diseases caused by Pythium and Phytophthora.

Use only at recommended rates as overdosage may be harmful to sensitive plants. Do not mix with other chemicals, other than Terraclor[®] unless prior use has proven compatibility.

RECOMMENDATIONS

In nurseries and greenhouses

For bedding and bench plantings use with water, as a soil drench at time of seeding and transplanting. The drench solution should be continually agitated to assure uniform application of the fungicidal material. To improve penetration of the fungicide, the drenching should be followed with an equal quantity of water applied either by sprinkling or by irrigation

<p>BEDDING PLANTS Alyssum, Calendula, Chrysanthemum, Celosia, Dahlia, Dianthus, Marigold, Pansies, Petunia, Salvia, Snapdragon, Vinca, Verbena, Zinnia.</p>	<p>BEDDING PLANTS: (Soil 2" to 3" Deep): - Add ¼ to 1 oz. to 25 gallons of water and apply to 100 sq. ft. of bed area. Repeat application may be made at one to two month intervals, if necessary.</p>
<p>FOLIAGE PLANTS Aglaonema, Anthurium, Chinese Evergreen, Dieffenbachia, Nephthytis, Peperomia, Pilea, Pothos, Philodendron Spp.</p> <p>HERBACEOUS AND WOODY: African Violet, Geranium, Rhododendron.</p>	<p>TROPICAL FOLIAGE, HERBACEOUS and WOODY PLANTS: (Peat and other high organic media, 6" deep) - Bed and Bench Plantings: Add 1 to 2 ozs. in 25 gallons of water and apply to 100 sq. ft. of bed or bench area. Repeat application at two to three month intervals to maintain disease control. Pot Plantings: Use ½ pint solution per 6" pot or in sufficient amount to saturate the soil.</p>

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.