

10F2  
PM 21  
400-419

**ACCEPTED**  
AUG 4 1986  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-419

# TERRAZOLE<sup>®</sup>

## 5% GRANULAR FUNGICIDE

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**  
**CAUTION**  
Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Do not handle or apply product with bare hands.  
**ENVIRONMENTAL HAZARDS**  
Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water.



**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store in a dry location.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.  
**HOUSEHOLD:** Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

**COMPOSITION**

Active Ingredient: (% by weight)	
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole*	5%
Inert Ingredients.	95%
Total:	100.0%

\*U.S. Patents 3,260,588 and 3,260,725: others pending.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**STATEMENT OF PRACTICAL TREATMENT**  
IF SWALLOWED: Induce vomiting.  
IF ON SKIN: Remove contaminated clothing and wash skin with soap and warm water.  
IF INHALED: Remove to fresh air.  
IF IN EYES: Flush with water for 15 minutes.  
For Further Treatment, Get Medical Treatment  
**SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT**

**Uniroyal Chemical Company, Inc.**  
Middlebury, CT 06749  
EPA REG. NO. 400-419  
EPA EST. NO.

4-24-86

TERRAZOLE is a Registered Trademark of Uniroyal, Inc.  
© Copyright 1983, Uniroyal, Inc.

Lot. No.

**NET CONTENTS:**

## DIRECTIONS FOR USE

**ORNAMENTALS** — For control of damping-off and root rot diseases caused by *Pythium* and *Phytophthora* at the time for seeding or transplanting in nurseries and greenhouses.

PLANT TYPE		RECOMMENDATIONS
<b>FOLIAGE PLANTS</b>		Dry Soil Mix:  To mix with potting, greenhouse or other transplanting soils, add Terrazole 5G at the rate of 10 ozs. per cubic yard of soil and mix thoroughly. Soil to be mixed should be free of large lumps and dry enough to prevent agglomeration during mixing. Where <i>Rhizoctonia</i> control is desired, add recommended quantities of Terraclor. Retreat as needed for continued control with soil drench applications of Terrazole 35% Wettable Powder or Terrazole 25% Emulsifiable.
Aglaonema Anthurium Brassaia Caladium Chinese Evergreen Dieffenbachia Dracaena	Echeveria Epipremnum Nepthytis Peperonia Pilea Philodendron Zygocactus	
<b>CONTAINER &amp; BEDGROWN</b>		NOTE: Overdosage may result in phytotoxicity. If plants are suspected of being sensitive, treat only a few plants and observe for 4 to 5 days prior to treating the entire planting.
<b>HERBACEOUS</b>	<b>WOODY</b>	
African Violet Begonia Calceolaria Chrysanthemum Easter Lily Geranium Gerbera Gloxinia Gypsophilia Orchids Sinningia	Arctostaphylos Azalea Camellia Ceanothus Ficus Fir Hemlock Ilex Juniper Leucothoe Magnolia Poinsettia Rhamnus Rhododendron Southern Pine Western Pine	

**AVOCADOS (California Only):** For control of Avocado root rot (*Phytophthora cinnamomi*). Apply Terrazole 5% Granular evenly to the soil surface surrounding the Avocado tree, in a circular area extending out to the drip line. The amount of material to be applied to each tree will be governed by the area to be covered, as per chart below:

Diameter of Circular Surface	Approximate Square Feet Covered	Amount of Terrazole 5% Granular per tree
20' diameter	314	86 ounces
15' diameter	176	48 ounces
10' diameter	79	22 ounces
5' diameter	20	6 ounces

Following application apply approx. one gallon of water per square foot of area treated. Repeat application once a month for 9 to 10 months during the growing season.

Do not treat within 30 days of harvest. During application, care should be exercised so that the treatment does not come in contact with the fruit.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.