

TERRAZOLE®

4 LB. EMULSIFIABLE ORNAMENTAL FUNGICIDE



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
WARNING**

May be fatal if absorbed through skin. Causes skin irritation. Harmful if swallowed or inhaled. Do not breathe vapors. Do not get on skin or on clothing. Wash contaminated clothing prior to reuse. When handling or using the concentrate, wear natural rubber gloves. Wash thoroughly with soap and water after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

COMPOSITION

Active Ingredient: (% by weight)	
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole	45.8%
Inert Ingredients:	54.2%
Total:	100.0%
Contains 4 lbs. of 5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole per U.S. Gallon at 68°F.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of water. Do not induce vomiting. Contains petroleum distillates. Call a physician.

IF ON SKIN: Remove contaminated clothing and wash skin with soap and warm water.

IF IN EYES: Flush with plenty of water for at least 15 minutes.

IF INHALED: Remove to fresh air.

If patient is unconscious, maintain breathing and heartbeat (CPR: cardiopulmonary resuscitation, if necessary). Contact your local Poison Control Center, hospital or physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

ACCEPTED

FEB 1 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-422

UNIROYAL CHEMICAL COMPANY, INC.
Middlebury, CT 06749
EPA REG. NO. 400-422
EPA EST. NO.

6-2-86

Lot. No.
TERRAZOLE is a Registered Trademark of UNIROYAL CHEMICAL COMPANY, INC.
© Copyright 1986, Uniroyal Chemical Company, Inc.

NET CONTENTS:

DIRECTIONS FOR USE

Terrazole is useful as a soil fungicide for control of damping-off, root and stem diseases caused by *Phythium* and *Phytophthora*. Use only at recommended rates as overdosage may be harmful to sensitive plants. Do not mix with chemicals other than Terraclor[®], unless prior use has proven compatibility.

ORNAMENTALS

For bedding and bench plantings use with water, as a soil drench at time of seeding and transplanting in nurseries and greenhouses. The drench solution should be continually agitated to assure uniform application of the fungicidal material. To improve penetration of the fungicide, the drenching should be followed with an equal quantity of water applied either by sprinkling or by irrigation.

<p>BEDDING PLANTS Alyssum Calendula Celosia Chrysanthemum Coleus Dahlia Dianthus Larkspur Marigold</p>	<p>Pansies Petunia Phlox Salvia Snapdragon Verbena Vinca Zinnia</p>	<p>BEDDING PLANTS: (Soil 2" to 3" deep) - Add 3 to 4 fl. oz. to 100 gallons of water and apply to 400 sq. ft. of bed area. Repeat application may be made at one to two month intervals, if necessary.</p>
<p>FOLIAGE PLANTS Aglaenema Anthurium Catus Caladium Calathea Chamadorea Chinese Evergreen Crassula Dieffenbachi Gynura HERBACEOUS AND WOODY African Violet Azalea Boxwood Crysanthemum Easter Lily Euonymous Ficus Geranium</p>	<p>Hedera Hoya Maranta Nephrolepis Nephtytis Peperomia Pilea Pothos Stromanthe Zygocatus</p> <p>Gerbera Juniper Kalanchoe Pieris Poinsettia Red Pine Rhododendron Sinningia</p>	<p>TROPICAL FOLIAGE HERBACEOUS AND WOODY PLANTS (Peat and other high organic media 6" deep).</p> <p>Bed and Bench Plantings: Add 2 to 4 fl. ozs. in 100 gallons of water and apply to 400 sq. ft. of bed or bench area. Repeat application at two to three month intervals to maintain disease control.</p> <p>Pot Plantings: Use 1/2 pint solution per 6" pot or in sufficient amount to saturate the soil.</p>

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

