

TERRAZOLE® 4EC

COMPOSITION

Active Ingredients: (% by weight)	
5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole*	44.3%
Inert Ingredients**	55.7%
Total:	100.0%

*Contains 4 lbs. of 5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole per U.S. Gallon at 20°C.

**Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Hold eyelids open and flush with water for 15 minutes. Call a physician if irritation persists.

IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Obtain prompt medical attention.

IF INHALED: Remove to victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage from the presence of petroleum distillates in the formulation may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if absorbed through skin. Causes substantial eye injury. Causes skin irritation. Harmful if swallowed or inhaled. Do not breathe vapors. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear: Coveralls over short-sleeved shirt and short pants; chemical-resistant gloves such as barrier laminate or viton; chemical-resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

ACCEPTED
COMMENTS
EPA Letter Dated
OCT 13 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 400-422



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Terrazole is useful as a soil fungicide for control of damping-off, root and stem diseases caused by Pythium and Phytophthora. Use only at recommended rates as overdosage may be harmful to sensitive plants. Do not mix with chemicals, other than Terraclor® unless prior use has proven compatibility.

ORNAMENTALS

For bedding and bench plantings use with water as a soil drench at time of seeding and transplanting in commercial nurseries and greenhouses. The drench solution should be continually agitated to assure uniform application of the fungicidal material. To improve penetration of the fungicide, the drenching should be followed with an equal quantity of water applied either by sprinkling or by irrigation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over short-sleeved shirt and short pants
- chemical-resistant gloves such as barrier laminate or viton
- chemical-resistant footwear plus socks
- protective eyewear
- chemical-resistant headgear for overhead exposure

<p style="text-align: center;">BEDDING PLANTS</p> <table border="0"> <tr> <td>Alyssum</td> <td>Dahlia</td> <td>Petunia</td> <td>Vinca</td> </tr> <tr> <td>Calendula</td> <td>Dianthus</td> <td>Phlox</td> <td>Zinnia</td> </tr> <tr> <td>Celosia</td> <td>Larkspur</td> <td>Salvia</td> <td></td> </tr> <tr> <td>Chrysanthemum</td> <td>Marigold</td> <td>Snapdragon</td> <td></td> </tr> <tr> <td>Coleus</td> <td>Pansies</td> <td>Verbena</td> <td></td> </tr> </table>	Alyssum	Dahlia	Petunia	Vinca	Calendula	Dianthus	Phlox	Zinnia	Celosia	Larkspur	Salvia		Chrysanthemum	Marigold	Snapdragon		Coleus	Pansies	Verbena		<p>BEDDING PLANTS: (Soil 2" to 3" Deep) -Add 3 to 4 fl. ozs. to 100 gallons of water and apply to 400 sq. ft. of bed area. Repeat application may be made at one to two month intervals, if necessary.</p>				
Alyssum	Dahlia	Petunia	Vinca																						
Calendula	Dianthus	Phlox	Zinnia																						
Celosia	Larkspur	Salvia																							
Chrysanthemum	Marigold	Snapdragon																							
Coleus	Pansies	Verbena																							
<p style="text-align: center;">FOLIAGE PLANTS</p> <table border="0"> <tr> <td>Aglaonema</td> <td>Chinese Evergreen</td> <td>Maranta</td> <td>Stromanthe</td> </tr> <tr> <td>Anthurium</td> <td>Crassula</td> <td>Nephrolepis</td> <td>Zygocactus</td> </tr> <tr> <td>Cactus</td> <td>Dieffenbachia</td> <td>Nephtytis</td> <td></td> </tr> <tr> <td>Caladium</td> <td>Gynura</td> <td>Peperomia</td> <td></td> </tr> <tr> <td>Calathea</td> <td>Hedera</td> <td>Pilea</td> <td></td> </tr> <tr> <td>Chamadorea</td> <td>Hoya</td> <td>Pothos</td> <td></td> </tr> </table>	Aglaonema	Chinese Evergreen	Maranta	Stromanthe	Anthurium	Crassula	Nephrolepis	Zygocactus	Cactus	Dieffenbachia	Nephtytis		Caladium	Gynura	Peperomia		Calathea	Hedera	Pilea		Chamadorea	Hoya	Pothos		<p>TROPICAL FOLIAGE, HERBACEOUS and WOODY PLANTS: (Peat and other high organic media 6" deep).</p> <p>Bed and Bench Plantings: Add 2 to 4 fl. ozs. in 100 gallons of water and apply to 400 sq. ft. of bed or bench area. Repeat application at two to three month intervals to maintain disease control.</p>
Aglaonema	Chinese Evergreen	Maranta	Stromanthe																						
Anthurium	Crassula	Nephrolepis	Zygocactus																						
Cactus	Dieffenbachia	Nephtytis																							
Caladium	Gynura	Peperomia																							
Calathea	Hedera	Pilea																							
Chamadorea	Hoya	Pothos																							
<p style="text-align: center;">HERBACEOUS AND WOODY PLANTS</p> <table border="0"> <tr> <td>African Violet</td> <td>Easter Lily</td> <td>Gerbera</td> <td>Poinsettia</td> </tr> <tr> <td>Azalea</td> <td>Euonymus</td> <td>Juniper</td> <td>Red Pine</td> </tr> <tr> <td>Boxwood</td> <td>Ficus</td> <td>Kalanchoe</td> <td>Rhododendron</td> </tr> <tr> <td>Chrysanthemum</td> <td>Geranium</td> <td>Pieris</td> <td>Sinningia</td> </tr> </table>	African Violet	Easter Lily	Gerbera	Poinsettia	Azalea	Euonymus	Juniper	Red Pine	Boxwood	Ficus	Kalanchoe	Rhododendron	Chrysanthemum	Geranium	Pieris	Sinningia	<p>Pot Plantings: Use ½ pint solution per 6" pot or in sufficient amount to saturate the soil.</p>								
African Violet	Easter Lily	Gerbera	Poinsettia																						
Azalea	Euonymus	Juniper	Red Pine																						
Boxwood	Ficus	Kalanchoe	Rhododendron																						
Chrysanthemum	Geranium	Pieris	Sinningia																						

BEST AVAILABLE COPY

COTTON

To control Water Mold seed decay (pre-emergence damping off) caused by *Pythium* spp.

Soil In-Furrow Application

At planting time use 6 to 12 oz. per acre (per 13,000 linear feet based on 40 inch rows) mixed with 5 to 15 gallons of water.

Apply the specified dosage to the soil around the seed and to the covering soil as it fills the furrow. The soil around the seed and the covering soil should be thoroughly mixed with the product. Two nozzles work best; center the front nozzle over the open furrow to spray the soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow. The higher rates should be used when weather conditions are expected to be unfavorable for rapid germination, fields have a history of disease problems or minimum or low till programs are utilized.

For control of *Pythium* and *Rhizoctonia*, apply 3 to 6 pints of Terraclor® 2E, 1.5 to 3.0 pints of Terraclor Flowable Fungicide or 1½ to 2½ lbs. Terraclor 75WP per 13,000 linear feet of row in a tank mixture with Terrazole 4EC.

NOTE:

When Terrazole 4EC is applied with other Terraclor products, observe all precautions and restrictions that appear on the Terraclor and Terrazole labels.

Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season.

Rotation Restriction: Do not plant root crops in PCNB treated fields within 12 months of application unless PCNB is registered for use on those crops.

Caution Statements

Do not allow the feeding or grazing of cotton foliage by livestock. Apply only at planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, **but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.**

®Terrazole and Terraclor are Registered Trademarks of UNIROYAL CHEMICAL COMPANY, INC.

©Copyright 1994, Uniroyal Chemical Company, Inc.

BEST AVAILABLE COPY