

OMITE®-30WS

THIS BAG CONTAINS 2 x 2½ LB. WATER SOLUBLE BAGS
SEE FOLDER FOR USE INSTRUCTIONS
ENCLOSED IN BOTTOM FOLD

AGRICULTURAL MITICIDE

Recommended for agricultural use only

Net contents:

COMPOSITION

Active Ingredients: (% by weight)

Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]	30.0%
Inert Ingredients	70.0%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER • PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Call a physician.

IF INHALED: Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Get medical attention. Do not induce vomiting. Do not give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; waterproof gloves; shoes plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing, or loading. Applicators, if applying more than 10 lbs. of OMITE-30W per acre in air-blast equipment must be in an enclosed cab.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

ACCEPTED

12 APR 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-427

**UNIROYAL
CHEMICAL**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) listed in chart below.

RESTRICTED ENTRY INTERVALS (REI)	
CROP	
Grapes (wine & raisin)	14 days
Grapes (table)	21 days
Citrus	21 days
All other labeled crops	7 days

Exception: After the first 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

US RESTRICTIONS

Do not plant leafy vegetables in rotation within 2 months after last application of propargite to cotton or corn.

Do not plant any other food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite.

Do not feed livestock on cover crops grown among the trees listed on this label.

GENERAL INSTRUCTIONS

OMITE-30WS is a dry wettable powder for preparing sprays to control mites.

OMITE-30WS is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. OMITE-30WS can be used in concentrate sprays.

OMITE-30WS performance is best when day temperatures average above 70°F.

MIXING INSTRUCTIONS

This package contains 2 x 2½ pound water soluble bags of OMITE-30WS. After opening package:

1. Fill spray tank with ½ the required amount of water and turn on agitation system.
2. Do not open or perforate the inner water soluble bags of OMITE-30WS. The water soluble bags and contents will dissolve in water.
3. Immediately put water soluble bag into spray tank and mix using vigorous agitation.
4. Fill tank with the remaining amount of required water.
5. Do not touch water soluble bags as handling may cause breakage.
6. Use only entire water soluble bag. Do not break bags or use partial bags.
7. Do not begin spraying until the bag is completely dissolved and the powder is thoroughly dispersed in the spray tank.
8. If only one water soluble bag is used, store the remaining bag in the original package in a dry location away from heat or cold.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Place water soluble bag directly into application equipment (see directions). Then dispose of empty outer bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FOR USE ON BEARING CROPS
DOSAGE INSTRUCTIONS (see application notes for specific directions)

CROP	MITES CONTROLLED	AMOUNT OMITE-30WS PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	EARLIEST HARVEST DAYS AFTER APPLICATION
			Ground	Aerial		
ALMONDS (CA & AZ)	clover European red two-spotted spider strawberry spider Pacific spider	5 - 10 lbs.	50	—	2	28
GRAPES (West of Rocky Mts.)	Pacific spider strawberry spider Willamette two-spotted spider	5 - 9 lbs.	40	—	2	21
HOPS ⁽¹⁾	two-spotted spider	5 lbs.	100	—	2	14
NECTARINES (West of Rocky Mts.)	Pacific spider brown almond	5 - 9 lbs.**	50	20	2	14
ORNAMENTALS ⁽³⁾ CARNATIONS CHRYSANTHEMUMS	two-spotted spider	1 - ½ lbs. per 100 gals. water	Spray to wet	—	—	—
CHRISTMAS TREES & CONIFERS ⁽⁴⁾ (for plantation & nursery use only)	Spider mites	4½ - 7½ lbs. (west)	100	10	3	—
		3 - 7½ lbs. (east)				
ROSES ⁽³⁾ (field grown)	two-spotted spider Pacific spider	2 - 5 lbs.	50	—	—	—
OTHER ORNAMENTALS ⁽³⁾ (field & nursery)	two-spotted spider Southern red	1 lbs. per 100 gallons of water	300 - 500	—	—	—
PEANUTS	two-spotted spider	3 - 5 lbs.	20	—	2	14
WALNUTS (CA only)	European red two-spotted spider	5 - 12½ lbs.	100	10	2	21

* Each water soluble bag contains 2½ lbs. of OMITE-30WS.

**Use a minimum of 7½ pounds of OMITE-30WS per acre when applied by air.

FOR USE AFTER HARVEST AND ON NON-BEARING TREES ONLY
 Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30WS PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR
CHERRIES (West of Rocky Mts.)	European red two-spotted spider Pacific spider McDaniel	6 lbs.	50	—
GRAPEFRUIT ⁽²⁾ (West of Rocky Mts.)	citrus red	7½ - 10½ lbs.	100	1
NAVEL ORANGES ⁽²⁾ (West of Rocky Mts.)	citrus red	7½ - 10½ lbs.	100	1

* Each water soluble bag contains 2½ lbs. of OMITE-30WS.