


PROCURE[®] 50W

AGRICULTURAL FUNGICIDE



COMPOSITION

Active Ingredients: (% by weight)	
1-[1-[[4-chloro-2-(trifluoromethyl) phenyl] imino]-2-propoxyethyl]-1H-imidazole	50.0%
Inert Ingredients	50.0%
Total:	100.0%

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
 EPA REG. NO. 400-431
 EPA EST. NO.

001WPS

NET WEIGHT

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for 15 minutes. Get medical attention.
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of Ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive, causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact with this product may cause allergic skin reactions in some individuals. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; waterproof gloves; shoes plus socks; protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables; use detergent and hot water. Keep and wash PPE separately from other laundry.
 When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label directions before using.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; waterproof gloves; shoes plus socks; and protective eyewear.

DIRECTIONS FOR USE

PROCURE 50W is a broad-spectrum foliar fungicide for control of certain diseases of grapes and pome fruits. PROCURE 50W is particularly effective against powdery mildew of grapes, apples and pears and scab of apples and pears. Against scab it is active as an eradicant by preventing disease symptoms after infection has occurred. It's anti-sporulant activity reduces scab spores after lesions become visible.

Read the entire label and directions before using PROCURE 50W Wettable Powder fungicide.

NOTE: Do not allow livestock to graze on treated orchard ground cover. Do not apply this product through any type of irrigation system.

RECOMMENDED APPLICATIONS PROCURE 50W

PROCURE 50W Wettable Powder can be applied by various ground spray equipment. Equipment should be frequently checked for calibration. Apply PROCURE 50W in sufficient water to ensure complete and thorough coverage of foliage, bloom and fruit.

Use higher recommended rates on more susceptible varieties and/or under higher disease pressure.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

GRAPES

2/3

PROCURE 50W should not be used as an eradicant for mildew-control. Use as a protectant according to the spray schedule below. Use PROCURE 50W at 4 - 8 oz./A. Select spray volume, nozzle adjustments, and ground speed to ensure complete coverage of fruit clusters and foliage. Tractor speed should not exceed 3 mph. Use a minimum of 50 gallons of water per acre for early season applications. Increase gallonage used as amount of foliage increases. Do not exceed 32 oz. PROCURE 50W/A/season. Do not apply within 7 days of harvest.

Recommended Times of Application and Rates of PROCURE 50W

Variety Susceptibility to Mildew	Type of Grape	Light to Moderate Disease Pressure	Heavy Disease Pressure
Light	Wine & Raisin	12" to 18" cane, bloom, 21 days post bloom. Use 4 - 6 oz./A. Do not exceed 21 days between applications.	12" to 18" cane, bloom, 14 to 21 days post bloom. 3 weeks after last application. Use 4 - 6 oz./A. Do not exceed 21 days between applications.
	Table	12" to 18" cane, bloom, 21 days post bloom. Use 4 - 6 oz./A. Do not exceed 21 days between applications.	12" to 18" cane, bloom, 14 to 21 days post bloom. 3 weeks after last application. Use 4 - 6 oz./A. Do not exceed 21 days between applications.
Additional applications may be needed for seasonal control.			
Moderate	Wine & Raisin	12" cane, bloom, 14 to 21 days post bloom and 3 weeks later. Use 4 - 6 oz./A. Do not exceed 14 days between applications at low rate of 4 oz./A.	12" cane, bloom, 14 to 21 days post bloom and 14 to 21 days later. Use 4 oz./A. in early sprays 6 oz./A in later.
	Table	12" cane, bloom, 14 to 21 days post bloom and 3 weeks later. Use 4 - 6 oz./A. Do not exceed 14 days between applications at low rate of 4 oz./A.	6" cane, 18" cane, bloom, 14 days post bloom, 14 days after last application. Use 4 oz./A. in early sprays, 6 oz./A. in later. Do not exceed 14 days between applications.
Additional applications may be needed for seasonal control.			
High	Wine & Raisin	12" cane, bloom, 14 days post bloom; additional application may be required 14 days after last application. Use 4 - 6 oz./A. Do not exceed 14 days between applications.	12" cane, bloom, 14 days post bloom, 14 days later. Do not exceed 14 days between applications; Use 6-8 oz./A.
	Table	6" cane, 18" cane, bloom, 14 days post bloom, 14 days after last application. Use 4 oz./A in early sprays, 6 oz./A in later. Do not exceed 14 days between applications.	6" cane and 18" cane at 4 oz./A. Bloom, 14 days post bloom, 14 days later, followed by an additional application. 14 days later at 6 oz./A.
Additional applications may be needed for seasonal control.			
Sulfur at recommended use rates can be applied before and/or between PROCURE 50W applications.			

APPLES & PEARS

On apples and pears, PROCURE 50W can be tank mixed with standard protectant fungicides during the season for maximum fruit disease control. Make applications at 7 to 10 day intervals through the first cover and at approximately 10 to 14 day intervals and thereafter as needed. PROCURE 50W may be applied up to 14 days prior to harvest.

CROP	DISEASE	RATE OF PROCURE 50W		COMMENTS
		oz/100 gal*	oz/acre	
Apples	Powdery Mildew	2 to 4	8 to 16	Initiate spraying at first sign of mildew, usually at the 1/2" green tip stage. Continue application at 7 to 14 day intervals as needed through the terminal growth period. PROCURE 50W may be applied up to 14 days prior to harvest.
	Scab and Cedar-Apple Rust	2 to 4	8 to 16	<p>Standard Spray Schedule Apply PROCURE 50W at the rate of 2 to 4 oz. per 100 gal. at 7 to 10 day intervals from 1/2" green tip stage through second cover or until sporulation is complete. Continue applications of PROCURE 50W in a standard 14 days cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the preliminary infection period, use 2 PROCURE applications, 7 days apart.</p> <p>Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 50W at the rate of 3 to 4 oz. per 100 gal. within 72 hours after the beginning of the scab infection period. Lower rate up to 3 oz. per 100 gal. will provide 48 "kick back" activity. (Use the higher rates under heavy disease pressure.</p> <p>After first cover, continue PROCURE 50W in a standard cover spray program at 2 to 3 oz. per 100 gal. in combination with protectant fungicides which control other diseases. Do not apply more than 64 oz. of PROCURE 50W per acre of apples per season.</p>
Pears	Powdery Mildew	2 to 4	8 to 16	Initiate spray at pink or white stage. Continue applications at 7 to 14 day intervals or as required. PROCURE 50W may be applied up to 14 days prior to harvest.
	Scab	2 to 4	8 to 16	<p>Standard Spray Schedule Apply PROCURE 50W at the rate of 2 to 4 oz. per 100 gal. at 7 to 10 day intervals from green tip stage through second cover. Continue applications of PROCURE 50W in a standard cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the preliminary infection period, use 2 PROCURE applications, 7 days apart.</p> <p>Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 50W at the rate of 3 to 4 oz. per 100 gal. within 72 hours after the beginning of the scab infection period. Lower rate up to 3 oz. per 100 gal. will provide 48 "kick back" activity. (Use the higher rates under heavy disease pressure.</p> <p>After first cover, continue PROCURE 50W in a standard cover spray program at 2 to 3 oz. per 100 gal. in combination with protectant fungicides which control other diseases. Do not apply more than 64 oz. of PROCURE 50W per acre of pears per season.</p>

*The rates of PROCURE 50W Wettable Powder are based on a standard of 400 gallons per acre of dilute spray. For less than standard size trees, apply the rate per 100 gallons until drip point. If using a concentrate spray, refer to the amount per acre to ensure equivalent amount of material is applied.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:


Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Paper: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

© PROCURE is a Registered Trademark of Uniroyal Chemical Company, Inc.



**UNIROYAL
CHEMICAL**

ACCEPTED
OCT 05 1994

COMPOSITION

Active Ingredients: (% by weight)	
1-[1-[[4-chloro-2-(trifluoromethyl) phenyl] imino]-2-propoxyethyl]-1H-imidazole	50.0%
Inert Ingredients	50.0%
Total:	100.0%

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
EPA REG. NO. 400-431
EPA EST. NO.

001WPS

NET WEIGHT

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive, causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact with this product may cause allergic skin reactions in some individuals. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; waterproof gloves; shoes plus socks; protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water.

Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Read entire label directions before using.
 Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; waterproof gloves; shoes plus socks; and protective eyewear

DIRECTIONS FOR USE

PROCURE 50W is a broad-spectrum foliar fungicide for control of certain diseases of grapes and pome fruits. PROCURE 50W is particularly effective against powdery mildew of grapes, apples and pears and scab of apples and pears. Against scab it is active as an eradicator by preventing disease symptoms after infection has occurred. Its anti-sporulant activity reduces scab spores after lesions become visible.

Read the entire label and directions before using PROCURE 50W Wettable Powder fungicide.

NOTE: Do not allow livestock to graze on treated orchard ground cover. Do not apply this product through any type of irrigation system.

RECOMMENDED APPLICATIONS PROCURE 50W

PROCURE 50W Wettable Powder can be applied by various ground spray equipment. Equipment should be frequently checked for calibration. Apply PROCURE 50W in sufficient water to ensure complete and thorough coverage of foliage, bloom and fruit.

Use higher recommended rates on more susceptible varieties and/or under higher disease pressure.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

GRAPES

PROCURE 50W should not be used as an eradicator for mildew-control. Use as a protectant according to the spray schedule below. Use PROCURE 50W at 4 - 8 oz./A. Select spray volume, nozzle adjustments, and ground speed to ensure complete coverage of fruit clusters and foliage. Tractor speed should not exceed 3 mph. Use a minimum of 50 gallons of water per acre for early season applications. Increase gallonage used as amount of foliage increases. Do not exceed 32 oz. PROCURE 50W/A/season. Do not apply within 7 days of harvest.

Recommended Times of Application and Rates of PROCURE 50W

Variety Susceptibility to Mildew	Type of Grape	Light to Moderate Disease Pressure	Heavy Disease Pressure
Light	Wine & Raisin	12" to 18" cane, bloom, 21 days post bloom. Use 4 - 6 oz./A. Do not exceed 21 days between applications.	12" to 18" cane, bloom, 14 to 21 days post bloom, 3 weeks after last application. Use 4 - 6 oz./A. Do not exceed 21 days between applications.
	Table	12" to 18" cane, bloom, 21 days post bloom. Use 4 - 6 oz./A. Do not exceed 21 days between applications.	12" to 18" cane, bloom, 14 to 21 days post bloom, 3 weeks after last application. Use 4 - 6 oz./A. Do not exceed 21 days between applications.
Additional applications may be needed for seasonal control.			
Moderate	Wine & Raisin	12" cane, bloom, 14 to 21 days post bloom and 3 weeks later. Use 4 - 6 oz./A. Do not exceed 14 days between applications at low rate of 4 oz./A.	12" cane, bloom, 14 to 21 days post bloom and 14 to 21 days later. Use 4 oz./A. in early sprays 6 oz./A. in later.
	Table	12" cane, bloom, 14 to 21 days post bloom and 3 weeks later. Use 4 - 6 oz./A. Do not exceed 14 days between applications at low rate of 4 oz./A.	6" cane, 18" cane, bloom, 14 days post bloom, 14 days after last application. Use 4 oz./A. in early sprays, 6 oz./A. in later. Do not exceed 14 days between applications.
Additional applications may be needed for seasonal control.			
High	Wine & Raisin	12" cane, bloom, 14 days post bloom; additional application may be required 14 days after last application. Use 4 - 6 oz./A. Do not exceed 14 days between applications.	12" cane, bloom, 14 days post bloom, 14 days later. Do not exceed 14 days between applications. Use 6-8 oz./A.
	Table	6" cane, 18" cane, bloom, 14 days post bloom, 14 days after last application. Use 4 oz./A in early sprays, 6 oz./A in later. Do not exceed 14 days between applications.	6" cane and 18" cane at 4 oz./A. Bloom, 14 days post bloom, 14 days later, followed by an additional application, 14 days later at 6 oz./A.
Additional applications may be needed for seasonal control.			
Sulfur at recommended use rates can be applied before and/or between PROCURE 50W applications.			

APPLES & PEARS

On apples and pears, PROCURE 50W can be tank mixed with standard protectant fungicides during the season for maximum fruit disease control. Make applications at 7 to 10 day intervals through the first cover and at approximately 10 to 14 day intervals and thereafter as needed. PROCURE 50W may be applied up to 14 days prior to harvest.

CROP	DISEASE	RATE OF PROCURE 50W		COMMENTS
		oz/100 gal*	oz/acre	
Apples	Powdery Mildew	2 to 4	8 to 16	Initiate spraying at first sign of mildew, usually at the 1/2" green tip stage. Continue application at 7 to 14 day intervals as needed through the terminal growth period. PROCURE 50W may be applied up to 14 days prior to harvest.
	Scab and Cedar-Apple Rust	2 to 4	8 to 16	<p>Standard Spray Schedule Apply PROCURE 50W at the rate of 2 to 4 oz. per 100 gal. at 7 to 10 day intervals from 1/2" green tip stage through second cover or until sporulation is complete. Continue applications of PROCURE 50W in a standard 14 days cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the preliminary infection period, use 2 PROCURE applications, 7 days apart.</p> <p>Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 50W at the rate of 3 to 4 oz. per 100 gal. within 72 hours after the beginning of the scab infection period. Lower rate up to 3 oz. per 100 gal. will provide 48 "kick back" activity. (Use the higher rates under heavy disease pressure.</p> <p>After first cover, continue PROCURE 50W in a standard cover spray program at 2 to 3 oz. per 100 gal. in combination with protectant fungicides which control other diseases. Do not apply more than 64 oz. of PROCURE 50W per acre of apples per season.</p>
Pears	Powdery Mildew	2 to 4	8 to 16	Initiate spray at pink or white stage. Continue applications at 7 to 14 day intervals or as required. PROCURE 50W may be applied up to 14 days prior to harvest.
	Scab	2 to 4	8 to 16	<p>Standard Spray Schedule Apply PROCURE 50W at the rate of 2 to 4 oz. per 100 gal. at 7 to 10 day intervals from green tip stage through second cover. Continue applications of PROCURE 50W in a standard cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the preliminary infection period, use 2 PROCURE applications, 7 days apart.</p> <p>Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 50W at the rate of 3 to 4 oz. per 100 gal. within 72 hours after the beginning of the scab infection period. Lower rate up to 3 oz. per 100 gal. will provide 48 "kick back" activity. (Use the higher rates under heavy disease pressure.</p> <p>After first cover, continue PROCURE 50W in a standard cover spray program at 2 to 3 oz. per 100 gal. in combination with protectant fungicides which control other diseases. Do not apply more than 64 oz. of PROCURE 50W per acre of pears per season.</p>

*The rates of PROCURE 50W Wettable Powder are based on a standard of 400 gallons per acre of dilute spray. For less than standard size trees, apply the rate per 100 gallons until drip point. If using a concentrate spray, refer to the amount per acre to ensure equivalent amount of material is applied.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Paper: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

® PROCURE is a Registered Trademark of Uniroyal Chemical Company, Inc.

