


PM 21 400-433

10/4

Terraguard 50W

ORNAMENTAL FUNGICIDE



ACCEPTED

MAY 01 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-433

50.0%
50.0%
100.0%

COMPOSITION

Active Ingredients: (% by weight)
 1-[1-[4-chloro-2-(trifluoromethyl) phenyl] propoxyethyl]-1H-imidazole 50.0%
 Inert Ingredients: 50.0%
 Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for 15 minutes.
IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of Ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

Net contents:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact with this product may cause allergic skin reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; waterproof gloves; shoes plus socks.
 Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.
 When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.



Uniroyal Chemical
 Company, Inc.
 Middlebury, CT 06749
 EPA REG. NO. 400-433
 EPA EST. NO.
 069WPS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Read entire label directions before using.
 Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; waterproof gloves; and shoes plus socks

GENERAL INFORMATION

TERRAGUARD 50W ornamental fungicide can be used in repeat applications as a cutting soak, soil drench, foliar spray or through chemigation for control of certain diseases on ornamental plants grown in enclosed commercial structures such as greenhouses, shadehouses, nurseries and interiorscapes.

When applied as a foliar spray, TERRAGUARD 50W has translaminal activity and is locally systemic within the plant. TERRAGUARD 50W provides excellent protectant activity and is most effective when applied prior to the onset of disease following recommended rates and application directions. However, when not used in preventive programs, TERRAGUARD 50W can also be applied after disease symptoms appear to provide good eradicant activity.

MIXING INSTRUCTIONS

Fill the mixing tank with half the required amount of water. Then add TERRAGUARD 50W with agitation running to insure proper mixing and suspension of product. Fill the tank with the remaining amount of required water.

The mixture of TERRAGUARD 50W and water should be continuously agitated to assure uniform application.

USE AND RATE RECOMMENDATIONS - ORNAMENTALS

FOLIAR DISEASES	OUNCES TERRAGUARD 50W PER 100 GALS. (TSP./GAL.)	APPLICATION NOTES
Aerial Blight (<i>Rhizoctonia solani</i>)	4 - 8 ($\frac{1}{2}$ - 1 $\frac{1}{2}$)	Apply sprays at weekly intervals as needed. Use high rates for initial applications under heavy disease pressure. Lower rates can be used for subsequent applications and preventive sprays.
Leaf Spot (<i>Alternaria</i> spp.) (<i>Exserohilum rostratum</i>) (<i>Helminthosporium rostratum</i>)		
Petiole Rot (<i>Myrothecium rotundum</i>)		
Powdery Mildew (<i>Erysiphe cichoracearum</i>) (<i>Erysiphe lagerstromiae</i>) (<i>Oidium</i> spp.) (<i>Podosphaera</i> spp.) (<i>Sphaerotheca pannosa</i>)	4 - 8 ($\frac{1}{2}$ - 1 $\frac{1}{2}$)	Apply sprays at weekly intervals as needed. Use higher rates under heavy disease pressure. Lower rates may be used for subsequent applications and preventive sprays.
	4 - 16 ($\frac{1}{2}$ - 2 $\frac{1}{2}$)	
		ROSES/POINSETTIAS ONLY: Apply sprays at weekly intervals as needed. Use the 16 oz./100 gal. rate for initial applications to existing infections. Lower rates of 4 - 8 oz./100 gals. should be used for subsequent applications and preventive sprays.

APPLICATION NOTES

FOLIAR SPRAYS: The recommended rate of application is 1 gallon of spray mixture to cover 200 sq. ft. of area or 200 gallons per acre. This will provide foliage coverage just to the point of drip. Do not handle treated plants until sprays have dried.

SOIL DRENCHES: Apply the recommended rate of TERRAGUARD 50W according to the guidelines below. The application should be made to plants which have been well watered one day prior to application. For best results, irrigation with additional water should not be made until 24 hours after application.

FOR CONTAINER GROWN PLANTS

Pot Diameter (inches)	Minimum Drench Volume (fl. oz./pot)
4	1.5
5	3
6	4
8	7
10	13
12	18

FOR BED AND BENCH GROWN PLANTS

Soil Depth (inches) to be Drenched	Coverage for TERRAGUARD 50W Drench Mix	
	Fl. oz./Sq. ft.	Sq. ft./100 gals.
2 or less	16	800
3	24	530
4 or more	32	400

DISEASES CONTROLLED

Common Name	Causal Agent
Aerial Blight/Root Rot	<i>Rhizoctonia solani</i>
Black Root Rot	<i>Thielaviopsis</i> spp.
Leaf Spot	<i>Alternaria</i> spp.
Leaf Spot	<i>Exserohilum rostratum</i>
Petiole Rot	<i>Myrothecium rotundum</i>
Powdery Mildew	<i>Erysiphe</i> spp.
Powdery Mildew	<i>Oidium</i> spp.
Powdery Mildew	<i>Podosphaera</i> spp.
Powdery Mildew	<i>Sphaerotheca pannosa</i>
Root Rot/Crown Canker	<i>Cylindrocladium scoparium</i>
Root Rot/Damping Off	<i>Cylindrocladium theae</i>
Root Rot/Petiole Rot	<i>Cylindrocladium spathiphylli</i>
Scab	<i>Venturia inaequalis</i>

3 of 4

USE AND RATE RECOMMENDATIONS - OF

TERRAGUARD 50W

SOIL BORNE DISEASES	OUNCES TERRAGUARD 50W PER 100 GALS. (TSP./GAL.)	APPLICATION NOTES
Black Root Rot (Thielaviopsis spp.)	1 - 4 (1/4 - 1/2)	Apply soil drenches at 3 to 4 week intervals as needed. Use higher rate under heavy disease pressure.
Cylindrocladium Root Rot and Petiole Rot (Cylindrocladium spathiphylli)	4 - 8 (1/2 - 1)	Apply soil drenches at 3 to 4 week intervals as needed. Use higher rate under disease pressure which can occur under warmer conditions. NOTE: Applications may be made as a heavy spray over the foliage however, the volume applied should be the same as that required for the soil drench to insure adequate soil penetration.
Cylindrocladium Root Rot, Wilt and Crown Canker (Cylindrocladium scoparium) (Cylindrocladium theae)	12 - 16 (2 - 2 1/2)	FOR PLANTS BEING PROPAGATED: The cutting soak plus soil drench application may be necessary to provide control, therefore treatment should consist of both methods of application. Pre-stick cutting soak Soak cuttings for 10 minutes. Slight agitation of the mixture is required to maintain proper suspension.
	5 - 8 (2 - 1 1/2)	Soil Drench: The higher rate should be used under heavier disease pressure. Apply additional drenches at 3 to 4 week intervals as needed. Normally 2 to 4 applications are sufficient. NOTE: Do not exceed 8 ozs./600 sq. ft. of bed or bench area as injury and/or inhibition of rooting may occur.
	6 - 12 (1 - 2)	FOR PLANTS WITH ESTABLISHED ROOT SYSTEMS: Soil Drench: Use higher rate under heavier disease pressure. Lower rates may be used in subsequent application. Apply additional drenches at 3 to 4 week intervals as needed. Normally 2 to 4 applications are sufficient. If reinfection occurs at a later date, retreatment may be necessary.
Root Rot (Rhizoctonia solani)	4 - 8 (1/2 - 1)	Apply soil drenches at 3 to 4 intervals as needed. Use higher rate under heavy disease pressure.

PLANT TOLERANCE:

When used on bedding plant plugs, do not exceed 2 ozs./100 gals. water. Some cultivars of impatiens have shown a sensitivity to applications of TERRAGUARD 50W.

DO NOT USE ON IMPATIENS PLUGS.

ON IMPATIENS TRANSPLANTS, DO NOT EXCEED 2 OZS./100 GALS.

Neither the manufacturer nor the seller has determined whether or not TERRAGUARD 50W can be used safely on all ornamental plants. Prior to any large scale application, the user should determine the safety of TERRAGUARD 50W by testing a small number of the type of plants to be treated at the recommended rates. Observe the treated plants for symptoms of phytotoxicity which may occur as stunting, foliage burn or, for plants being propagated, as an inhibition of rooting. This may take up to two months for species that do not root readily. The user assumes all risks arising out of application to unlabeled plants.

USE DIRECTIONS FOR CHEMIGATION:

In addition to the above use rates and recommendations, the following precautions must be observed when using this product in any type of irrigation system:

Apply this product only through the following systems:

- 1) Overhead sprinklers such as impact or micro-sprinklers,
- 2) Micro-irrigation such as spaghetti-tube or individual tube irrigation,
- 3) Mist-type irrigation such as fog systems,
- 4) Hand-held calibrated irrigation equipment such as the hand-held wand with injector.

Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system, (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection

pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically

484

shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

IMPORTANT NOTICE

The directions and recommendations on this label are derived from research to ensure correct product usage. Insufficient control may result from extremes in weather conditions, or lack of following label instruction. The use of this product is beyond the control of Uniroyal and seller. Buyer is responsible for proper use as per directions and acceptance of product performance under extraordinary or unusual weather conditions.

Uniroyal is not responsible for losses or damage resulting from using TERRAGUARD in any manner not specifically recommended.

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paper: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

© TERRAGUARD is a Registered Trademark of Uniroyal Chemical Company, Inc.

© Copyright 1996, Uniroyal Chemical Company, Inc.