

400-453

02-20-2004

19

EPA	United States	<input type="checkbox"/> Registration	OPP Identifier Number
	Environmental Protection Agency	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

## Application for Pesticide - Section I

1. Company / Product Name 400-453	2. EPA Product Manager Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company / Product (Name) Terraclor® Flowable Fungicide	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Crompton Corporation 74 Amity Road Bethany, Connecticut 06524-3402 <input type="checkbox"/> check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to: EPA Reg No. _____ Product Name _____	

## Section II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application. <b>NOTIFICATION</b>
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below. <b>FEB 20 2004</b>

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

- This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

- modification of potato footnote for California

## Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt	No. Per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling Accompanying product
6. Manner in Which Label is Applied to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled Other _____			

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Betsy K. Katzman	Title Registration Specialist	Telephone No. (Include Area Code) (203) 393-2163
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Betsy K. Katzman</i>	3. Title Manager U.S. Registration	
4. Typed Name Willard F. Cummings	5. Date February 4, 2004	

# Certification with Respect to Label Integrity

version 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
400-453	02/04/04	000400-00453.20040204015.potato.pdf


I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

*Willard F. Cummings*  
 Signature


02/04/2004  
 Date

Willard F. Cummings  
 Name (typed)

U.S. Registration Manager  
 Title



# Terraclor<sup>®</sup> Flowable Fungicide



**Active Ingredient:** (% by weight)  
 Pentachloronitrobenzene (PCNB) ..... 40%  
 Inert Ingredients: ..... 60%  
 TOTAL ..... 100%

Contains 4 lbs. of Pentachloronitrobenzene per U.S. gallon.

**Net Contents:**

## KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR ON CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor immediately for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>CROMPTON EMERGENCY PHONE</b>	<b>800-292-5898</b>
<b>SAFETY DATA AND INFORMATION</b>	<b>203-573-3303</b>
<b>TRANSPORTATION EMERGENCY (CHEMTREC)</b>	<b>800-424-9300</b>

Precautionary Statements and Directions for Use are in the attached booklet. If booklet is missing, contact Crompton or an authorized dealer.

EPA REG. NO. 400-453  
EPA EST. NO.  
015CM/020404

Crompton Manufacturing Company, Inc.  
Middlebury, CT 06749

Crompton<sup>®</sup>, and Uniroyal Chemical<sup>®</sup>  
are Registered Trademarks of a  
subsidiary of Crompton Corporation



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

**Applicators and Other Handlers (other than mixers and loaders) Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks.

**Mixers and Loaders Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks. For exposures in enclosed areas: A respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter. For exposures outdoors: Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified by WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

DO NOT PLANT ROOT CROPS IN PCNB TREATED FIELDS WITHIN 12 MONTHS OF BROADCAST AND BANDING APPLICATIONS UNLESS PCNB IS REGISTERED FOR USE ON THOSE CROPS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical-resistant gloves such as barrier laminate or nitrile rubber, or neoprene rubber, or viton
- shoes and socks.

**GENERAL INFORMATION**

TERRACLOR FLOWABLE is a protective fungicide recommended for control of certain soil borne diseases of vegetable and field crops. Best results are achieved when used in a preventative program following recommended rates and application directions. Carefully read, understand and follow label directions.

**TANK MIXTURE COMPATIBILITY**

Combinations of TERRACLOR FLOWABLE with fungicides, insecticides, plant growth regulators or fertilizers should be checked for compatibility before tank mixing. Use a jar test to mix the recommended proportions of each product in a volume of water which is proportionate to the expected field use rate. The mix is considered compatible if it remains stable or can be easily reconstituted by shaking after standing for 5 minutes.

**VEGETABLES**

TERRACLOR FLOWABLE should be applied as a preventative fungicide to vegetable crops. It is important when using TERRACLOR FLOWABLE as a protective fungicide that material be placed into the soil. This can be accomplished as a soil drench, mechanical incorporation or as an in-furrow application. See use and rate recommendations for specific crop use recommendations and diseases controlled by TERRACLOR FLOWABLE.

When rate range is shown the lower rate should be used on lighter soils or disease pressure is expected to be low. The higher rates should be used on heavy soils, in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

**CROP**

**DISEASES CONTROLLED**

BEANS (snap & dry)	Root & Stem Rot ( <i>Rhizoctonia solani</i> )
PEPPERS, TOMATOES	Southern Blight ( <i>Sclerotium rolfsii</i> )
GARLIC	White Rot ( <i>Sclerotium cepivorum</i> )
BROCCOLI	Wire Stem or Black Root ( <i>Rhizoctonia solani</i> )
BRUSSELS SPROUTS	Club Root ( <i>Plasmodiophora brassicae</i> )
CABBAGE	
CHINESE BROCCOLI	
CHINESE CABBAGE	
CAULIFLOWER	
COLLARDS	
KALE	
MUSTARD GREENS	
COTTON	Damping Off ( <i>Rhizoctonia solani</i> )
HOT PEPPERS (direct seeded)	Root & Stem Rot ( <i>Rhizoctonia solani</i> )
PEANUTS	Root & Stem Rot ( <i>Rhizoctonia solani</i> ) Southern Stem Rot/White Mold ( <i>Sclerotium rolfsii</i> )
POTATOES	Stem Canker/Black Scurf ( <i>Rhizoctonia solani</i> )
VEGETABLE & BEDDING PLANTS	Root & Stem Rot, Damping Off ( <i>Rhizoctonia solani</i> )

CROP	DISEASE	RATE OF TERRACLOR FLOWABLE FUNGICIDE		RECOMMENDED APPLICATION
		AMOUNT/ACRE	FL. OUNCE/1000 FEET OF ROW	
Peanuts	Root and Stem Rot ( <i>Rhizoctonia solani</i> )	2 to 4 pints Apply in 8 to 10 gallons of water. (36 inch row spacing)	2.2 to 4.4	<b>At Planting:</b> Apply specified dosage as a directed in-furrow spray to the seed and surrounding soil at planting time. The soil around the seed and the covering soil should be thoroughly mixed with the product. The higher rate should be used when weather conditions are expected to be unfavorable for rapid germination, fields have a history of disease problems or minimum or low till programs are utilized. Do not use in California
	Southern Stem or White Mold ( <i>Sclerotium rolfsii</i> )	1.5 to 3 pints Apply in 8 to 10 gallons of water. (36 inch row spacing)	1.8 to 3.7	
<b>Do not exceed 2 lbs. PCNB active ingredient per acre in any one season, regardless of the method of application or formulation used.</b> <b>Apply only at planting time.</b> <b>Do not feed treated vines to livestock.</b> <b>Do not allow hogging down of peanuts.</b>				
Potatoes*	Stem Canker/ Black Scurf ( <i>Rhizoctonia solani</i> )	5 to 10 pints per acre. Apply in 10 to 20 gallons of water. (34 inch row spacing)	5.2 to 10.4	<b>Band Application:</b> Spray an 8½ inch band in seed furrow at time of planting. Direct spray into furrow over the seed and cover as a part of the tilling operation during planting.
<b>Do not exceed 20 lbs. of actual PCNB per acre in any one season regardless of method of application or formulation used.</b> <b>*Use not registered by the California Department of Pesticide Regulation.</b>				
Broccoli, Chinese broccoli (gai lan, white flowering broccoli), Brussels sprouts, Cabbage, Chinese cabbage (tight-heading varieties only), Cauliflower, Collards*, Kale*, Mustard greens*	Clubroot ( <i>Plasmodiophora brassicae</i> )	Mix 3 pints in 100 gallons of water. Use ½ pint of this solution per plant.		<b>Transplant Solution:</b> The applied dosage of TERRACLOR should be based on the severity of the disease infection. Maintain agitation to hold in suspension.
		- OR -		
		5.6 gal. Apply in 25 gallons of water. (40 inch row spacings)	55.0	<b>Band Application:</b> Spray as a 12 to 15 inch band centered on the row and incorporate to a depth of 4 to 6 inches immediately prior to planting. May be used on cole crops which are direct seeded.
	- OR -			
	7.5 gal. Apply in 30 gallons of water.	-	<b>Broadcast Application:</b> Apply as a preplant broadcast prior to planting. Thoroughly incorporate to a depth of 4 to 6 inches using a disc or other suitable equipment. May be used on cole crops which are direct seeded.	
	<b>Note: for effective control of clubroot, it is extremely important to thoroughly mix TERRACLOR with the soil.</b>			
- OR -				
Wire Stem or Black Root ( <i>Rhizoctonia solani</i> )	2.8 to 3.8 gal. Apply in 50 gallons of water.	-	<b>Broadcast Drench Application:</b> Apply as a soil drench at the time of or immediately after seeding. For smaller areas use 1 level tablespoon per gallon of water as a soil drench per 50 sq. ft. of seed bed. Apply with water can or similar equipment.	
	- OR -			
	1.9 to 2.8 gal. Apply in 35 gallons of water. (40 inch row spacings)	18.6 to 27.4	<b>Row Drench Treatment:</b> Spray as an 8 inch band centered on the row at the time of or immediately after seeding.	
<b>* Based on the available residue data, the use of TERRACLOR on collards, kale and mustard greens is restricted to the state of Georgia. Apply only at planting time. Do not exceed 30 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.</b>				
Beans dry, succulent & snap (excluding cowpeas)	Root and Stem Rot ( <i>Rhizoctonia solani</i> )	2 to 3 pints Use in 8 to 10 gallons of water. (36 inch row spacing)	2.2 to 3.3	<b>At Planting:</b> Spray planting furrow and covering soil at planting time. Avoid spraying directly to seed; otherwise delayed emergence may occur. Use lower rate on lighter soils.
		In California & Arizona: 1.5 to 2.0 pints in 8 to 10 gals. (spacings as above)	1.7 to 2.2	
<b>Apply only at planting time.</b>				
Hot Peppers (direct seeded)	Root and Stem Rot ( <i>Rhizoctonia solani</i> )	1.5 to 3.0 pints Use in 8 to 10 gallons of water. (40 inch row spacing)	2.2 to 3.3	<b>In-Furrow Application:</b> Apply only at planting time. Apply to the soil around the seed and to the covering soil as it fills the furrow.

CROP	DISEASE	RATE OF TERRACLOR FLOWABLE		RECOMMENDED APPLICATION
		AMOUNT/ACRE	FL. OUNCE/1000 FEET OF ROW	

<b>Cotton</b>	Damping-off or Soreshin ( <i>Rhizoctonia solani</i> )	1.5 to 3.0 pints Use in 3 to 10 gallons of water. (40 inch row spacings)	1.85 to 3.70	<b>In-Furrow Application:</b> Apply to soil around the seed and to the covering soil as it fills the furrow using suitable spray equipment. Two (2) nozzles work best, center the front nozzle to spray the open furrow to cover the soil around the seed and the rear nozzle to spray the covering soil as it fills the furrow.
---------------	---	--	--------------	--

**NOTE:** Use maximum rate of TERRACLOR FLOWABLE for no-till cotton or when disease pressure is expected to be high.

For broader spectrum disease control (*Rhizoctonia* and *Pythium* spp.), TERRACLOR FLOWABLE can be mixed with pythium control products such as Terrazole® 4EC or Ridomil Gold®. Because local conditions can influence the compatibility of TERRACLOR FLOWABLE with these products, a jar test should be done before tank mixing larger quantities. See TANK MIXTURE COMPATIBILITY section of this label for guidance. TERRACLOR FLOWABLE is not compatible with Di-Syston® 8. When TERRACLOR FLOWABLE is applied with either Terrazole 4EC or Ridomil Gold®, all precautions and restrictions on those labels must be observed.

Do not exceed 2 lbs. of PCNB active ingredient per acre in any one season applied as spray or granular formulations.

Apply only at planting time.

Do not allow the feeding or grazing of cotton foliage by livestock.

Ridomil Gold® is a registered trademark of Syngenta

Di-Syston® is a registered trademark of Bayer AG, Germany

<b>Garlic</b>	White Rot ( <i>Sclerotium cepivorum</i> )	5 gal. Apply in 100 gallons of water. (24 inch row spacings)	29.4	<b>At Planting:</b> Apply at planting. Attach a suitable rig to the planter and spray 8 to 12 inch bands using 2 nozzles per planting furrow. Direct front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.
---------------	---	--	------	--

— OR —

2.5 gals. per 1,000 lbs. of "seed" cloves.

**Clove Treatment:** Mix in sufficient water to apply dosage as a uniform mist-spray for complete coverage of treated cloves. For optimum results add a commercial sticker to the spray solution.

**Apply only at planting time. Do not exceed 20 lbs. of PCNB active ingredient per acre in any one season regardless of the method of application or formulation used.**

<b>Peppers Tomatoes</b>	Southern Blight ( <i>Sclerotium rolfisii</i> )	Mix 4.5 pints in 100 gallons of water. Use ½ pint of this solution per plant.		<b>Transplant Solution:</b> Apply at the time of transplanting. The solution should be agitated often to maintain a uniform suspension and assure proper dosage. Apply only at planting time. TOMATOES - Do not exceed 7.0 lbs. PCNB active ingredient per acre in any one season (based on 5000 plants/acre) or 16.5 lbs. PCNB active ingredient per acre in any one season (based on 11,733 plants/acre). PEPPERS - Do not exceed 20 lbs. PCNB active ingredient per acre in any one season.
-----------------------------	--	--	--	---

— OR —

1.2 to 1.8 gal.  
Apply in 100 gallons of water. (36 inch row spacings)

10.6 to 15.8

**In-Furrow:** Apply as an in-furrow spray to the open "V" trench, just prior to planting. When cultivating, set plows as flat as possible to avoid getting untreated soil against stems of plants. Apply only at planting time. Do not exceed 7.2 lbs. of PCNB active ingredient per acre in any one season.

**ROTATION RESTRICTION:** Do not plant root crops in PCNB treated fields within 12 months of broadcast and banding applications unless PCNB is registered for use on those crops.

CROP	DISEASE	RATE OF TERRACLOR FLOWABLE FUNGICIDE	APPLICATION NOTES															
<b>Vegetable Bedding Plants</b> Beans (excluding cowpeas) Broccoli Brussel sprouts Cabbage Cauliflower Peppers Tomatoes	Root & Stem Rot, Damping-Off ( <i>Rhizoctonia solani</i> )	¾ - 1 pints 6 - 12 fl. oz.	<b>Soil drench applications to seedlings grown in containers or beds prior to transplanting into the field:</b> Mix the specified amount in 100 gallons of water and apply according to the following guidelines. One repeat application can be made 4 to 6 weeks after the initial application.  <table border="1"> <thead> <tr> <th colspan="3">Coverage for TERRACLOR Flowable Fungicide Drench Mix</th> </tr> <tr> <th>Soil Depth Inches</th> <th>Fl. oz./Sq. ft.</th> <th>Sq. ft./100 gals.</th> </tr> </thead> <tbody> <tr> <td>2 or less</td> <td>16</td> <td>800</td> </tr> <tr> <td>3</td> <td>24</td> <td>530</td> </tr> <tr> <td>4 or more</td> <td>32</td> <td>400</td> </tr> </tbody> </table>	Coverage for TERRACLOR Flowable Fungicide Drench Mix			Soil Depth Inches	Fl. oz./Sq. ft.	Sq. ft./100 gals.	2 or less	16	800	3	24	530	4 or more	32	400
Coverage for TERRACLOR Flowable Fungicide Drench Mix																		
Soil Depth Inches	Fl. oz./Sq. ft.	Sq. ft./100 gals.																
2 or less	16	800																
3	24	530																
4 or more	32	400																

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not reuse empty containers. Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Pressure rinse is preferred. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State authorities.

**IMPORTANT NOTICE** – Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

®TERRACLOR is a Registered Trademark of Crompton Corporation

© Copyright 2004, Crompton Corporation

**Crompton**  
Uniroyal Chemical

9/19  
EPA  
February 4, 2004

Document Processing Desk (NOTIF)  
Office of Pesticide Programs  
U.S. Environmental Protection Agency  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, Virginia 22202-4501

Attention: Ms. Mary Waller, PM 21  
Registration Division (7505C)

Subject: **Terraclor® Flowable Fungicide**  
**EPA Reg No. 400-453**  
**Label Notification - modification of potato footnote for California**

Dear Ms. Waller:

Crompton Corp. is submitting revised labels for the registration of Terraclor Flowable Fungicide (EPA Reg. No. 400-453). The footnote associated with the potato use directions has been modified. No other change has been done to this label.

This submission includes the following:

- 1) Application for Pesticide Label Amendment (form 8570-1).
- 2) Five copies of the revised Terraclor FF (400-453) label.
- 3) One copy of the revised Terraclor FF label with changes highlighted.
- 4) One copy of the current Terraclor FF label.
- 5) Certification with Respect to Label Integrity form.
- 6) PDF file of proposed product label on CD-ROM named:  
000400-00453.20040204015.potato.pdf

Should you have any questions regarding this submission, please contact Betsy Katzman at (203) 393-2163.

Sincerely,

CROMPTON MANUFACTURING COMPANY, INC.

*Betsy K. Katz* for  
Willard F. Cummings  
U.S. Registration Manager  
Crop Protection  
Research & Development