

PM 21 400-454 08/31/99

# Turficide® 4F

FLOWABLE  
TURF & ORNAMENTAL SOIL FUNGICIDE

ACCEPTED

AUG 31 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-454

### COMPOSITION

Active Ingredient: (% by weight)	
Pentachloronitrobenzene (PCNB)	40%
Inert Ingredients:	60%
TOTAL	100%

Contains 4 lbs. of Pentachloronitrobenzene per U.S. gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.  
**IF ON SKIN:** Wash with plenty of soap and water.  
**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing vapor or spray mist.

#### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

**Applicators and Other Handlers (other than mixers and loaders) Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks.

**Mixers and Loaders Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks. For exposures in enclosed areas: A respirator with an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter. For exposure outdoors: Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Net contents:

**UNIROYAL  
CHEMICAL**

Uniroyal Chemical  
Company, Inc.  
Middlebury, CT 06749  
EPA REG. NO. 400-454  
EPA EST. NO.  
017WPS

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product to turf through any type of irrigation system.

Do not plant root crops in PCNB treated fields within 12 months of broadcast or banding applications unless PCNB is registered for use on those crops.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves such as barrier laminate or nitrile rubber, or neoprene rubber, or viton
- shoes plus socks

**GENERAL INFORMATION**

TURFCIDE 4F is a preventative fungicide recommended for control of certain soil borne diseases of turf and ornamental crops.

Best results are achieved when used in a preventative program following recommended rates and application directions.

Carefully read, understand and follow label directions.

**COMPATIBILITY**

For broad spectrum disease control, this product can be tank mixed with TERRAZOLE® 35WP\* (EPA Reg. No. 400-416) for use on turf and ornamentals in accordance with the more restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not tank mix with other chemicals unless prior use has proven compatibility.

**LAWNS AND TURF**

TURFCIDE 4F is a fungicide for the prevention and control of certain soil borne diseases on golf courses, sod farms, home lawns and institutional areas where turf is grown.

Best results are obtained when TURFCIDE 4F is used as a preventative application before diseases appear, however, if you do not apply as a preventative treatment, TURFCIDE 4F should be applied at first sign of disease.

Areas treated with TURFCIDE 4F should be watered following application to move material down to soil level. If treated area is subjected to flooding or unusually heavy rainfall, or if disease is severe or reappears, the area should be treated again.

See use and rate recommendations for retreatment intervals. Do not graze treated areas or feed clippings to livestock.

Under certain growing conditions, a temporary discoloration of the grass may occur. This causes no harm and will disappear in a short time.

**USE AND RATE RECOMMENDATIONS FOR TURF**

DISEASE	TURF TYPE	FL. OZS. PER 1000 SQ. FT.	APPLICATION NOTES
Brown Patch ( <i>Rhizoctonia solani</i> )	Warm climate grasses such as: St. Augustine Bermuda	16-24	Mix specified amount in 1-5 gals. of water and apply a single application in the spring or fall or at first indication of disease. If disease is severe or reappears, area should be retreated 3 - 4 weeks later.
	Dichondra	16-24	Mix specified amount in 5-10 gals. of water and treat as above.
	Cool climate grasses such as: Bluegrass Ryegrass Fescue Bent	4½ to 6	Mix specified amount in 1-5 gals. of water and apply at first sign of disease. Repeat every 7 to 10 days during warm, damp weather. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85° or higher.
Melting Out/ Leaf spot* ( <i>Drechslera spp.</i> , <i>Bipolaris spp.</i> , <i>Exserchilum spp.</i> )  Dollar spot* ( <i>Sclerotinia homocarpa</i> )	All	10½ to 15	Mix specified amount in 1-5 gals. of water and apply as a preventative application in the spring or fall of each year. For control of existing infections, apply at higher rates as soon as symptoms appear. Do not apply at daytime temperatures of 85°F or higher.
		3	For maximum spring disease protection of Melting Out/Leaf Spot on cool season grasses, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications at two week intervals. Do not make more than a total of 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.
Gray Snow Mold ( <i>Typhula spp.</i> )  Pink Snow Mold ( <i>Microdochium nivale</i> )	Cool climate grasses	12-16	Mix specified amount in 1-5 gals. of water and apply one application prior to first snowfall or when temperatures remain below 60°F and extended wet conditions are expected. In the absence of snowcover, supplemental applications of 3-6 fl. ozs. can be applied at 4 to 6 week intervals to maintain control when disease pressure is severe. *Note: TURFCIDE must be moved into the soil thatch layer to be effective. Applications made in less than 5 gals. of water must receive 0.25" of irrigation or rainfall prior to snow cover.
		3	For maximum spring disease protection of pink snow mold, apply 3 fl. ozs. in 1-5 gals. of water per 1000 sq. ft. at the time of first mowing and repeat applications at two week intervals. Do not make more than a total 5 applications. Tank mix applications with benzimidazole or sterol inhibitor fungicides late in the program may enhance activity against Dollar Spot and Brown Patch disease.

\*Use not registered by the California Department of Pesticide Regulation

**ORNAMENTALS**

**DISEASES CONTROLLED**

TURFCIDE 4F can be used as a soil treatment for protection against a variety of soil borne diseases of ornamentals grown in greenhouses, shadehouses, nurseries, landscape plantings and interiorscapes.  
 The mixture of TURFCIDE 4F and water should be continuously agitated to assure uniform application.  
 It is important that the mixture of TURFCIDE 4F and water be drenched or incorporated thoroughly into the soil for best results.  
 See use and rate recommendations for specific plants and diseases.  
 When a rate range is shown, the lower rate should be used on lighter or coarse textured soils or when disease pressure is expected to be low.  
 The higher rates should be used on heavy or fine textured soils in fields with a history of disease problems, or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

**Causal Agent**

- Botrytis cinerea*
- Dothistroma pini*
- Ovulinia azaleae*
- Pellicularia filamentosa*
- Pellicularia rolfsii*
- Phyllostica cookeri*
- Rhizoctonia solani*
- Rhizoctonia spp.*
- Sclerotinia camelliae*
- Sclerotinia gladioli*
- Sclerotinia sclerotiorum*
- Sclerotium rolfsii*
- Stromatinia gladioli*

**Common Name**

- Storage Rot
- Needle Blight
- Azalea Petal Blight
- Root Rot / Stem Rot
- Bulb Rot / Crown Rot
- Magnolia Leaf Spot
- Root Rot / Stem Rot
- Root Rot / Stem Rot
- Camellia Flower Blight
- Dry Rot
- Stem Rot / Black Rot
- Bulb Rot / Crown Rot
- Dry Rot

**USE AND RATE RECOMMENDATIONS FOR ORNAMENTALS**

CROP	DISEASE	PINTS OF TURFCIDE 4F	APPLICATION NOTES																										
Bedding Plants Flowering Plants Foliage Plants Woody Plants such as but not limited to: Anthurium Azalea Chrysanthemum Dianthus Dieffenbachia Dracaena Ficus Gardenia Impatiens Lilies Marigold Pansy Petunia Poinsettia Pothos Rhododendron Schefflera Spathiphyllum	Root / Stem Rot, Damping Off <i>(Rhizoctonia solani)</i> <i>(Pellicularia filamentosa)</i> <i>(Sclerotinia sclerotiorum)</i>	$\frac{3}{8}$ - $\frac{3}{4}$ (6 - 12 fl. oz.)	<b>SOIL DRENCH APPLICATIONS: Mix specified amount in 100 gals. of water and apply according to the following guidelines:</b>  <b>For Bed, Bench, Container and Field Grown Plants:</b> <table border="1" style="width: 100%;"> <thead> <tr> <th>Soil Depth (inches)</th> <th>Coverage for TURFCIDE 4F Fl. oz./Sq. ft.</th> <th>Sq. ft./100 gals.</th> </tr> </thead> <tbody> <tr> <td>2 or less</td> <td>16</td> <td>800</td> </tr> <tr> <td>3</td> <td>24</td> <td>530</td> </tr> <tr> <td>4 or more</td> <td>32</td> <td>400</td> </tr> </tbody> </table> <b>For Potted Plants:</b> <table border="1" style="width: 100%;"> <thead> <tr> <th>Pot Diameter (inches)</th> <th>Minimum Drench Volume (fl. oz./pot)</th> </tr> </thead> <tbody> <tr> <td>4</td> <td>1.5</td> </tr> <tr> <td>5</td> <td>3</td> </tr> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>8</td> <td>7</td> </tr> <tr> <td>10</td> <td>13</td> </tr> <tr> <td>12</td> <td>18</td> </tr> </tbody> </table> One repeat application may be made 4 to 6 weeks later, if necessary.	Soil Depth (inches)	Coverage for TURFCIDE 4F Fl. oz./Sq. ft.	Sq. ft./100 gals.	2 or less	16	800	3	24	530	4 or more	32	400	Pot Diameter (inches)	Minimum Drench Volume (fl. oz./pot)	4	1.5	5	3	6	4	8	7	10	13	12	18
		Soil Depth (inches)	Coverage for TURFCIDE 4F Fl. oz./Sq. ft.	Sq. ft./100 gals.																									
2 or less	16	800																											
3	24	530																											
4 or more	32	400																											
Pot Diameter (inches)	Minimum Drench Volume (fl. oz./pot)																												
4	1.5																												
5	3																												
6	4																												
8	7																												
10	13																												
12	18																												
Azaleas Camellias	Azalea Petal Blight <i>(Ovulinia azaleae)</i>  Camellia Flower Blight <i>(Sclerotinia camelliae)</i>	$1\frac{1}{2}$	<b>BROADCAST INCORPORATED PRIOR TO PLANTING IN THE FIELD:</b> Apply specified amount to 1000 sq. ft. in sufficient water to assure uniform ground coverage. Thoroughly incorporate to a depth of 4 inches.  <b>GROUND SPRAY ON ESTABLISHED PLANTS:</b> Apply specified amount to 150 sq. ft. in sufficient water to assure uniform ground coverage. Apply to ground beneath the bushes and immediate surrounding area. Begin application prior to opening of buds and repeat every 3-4 weeks through bloom period.																										
Hyacinth Iris (bulbous) Narcissus Tulip	Root Rot / Stem Rot / Bulb Rot <i>(Rhizoctonia solani)</i> <i>(Pellicularia solani)</i> Bulb Rot / Crown Rot <i>(Sclerotium rolfsii)</i> <i>(Pellicularia rolfsii)</i> Black Rot <i>(Sclerotinia sclerotiorum)</i>	5 - $9\frac{1}{4}$	<b>BROADCAST INCORPORATED PRIOR TO PLANTING:</b> Apply specified amount to 1000 sq. ft. in sufficient water to insure uniform ground coverage. Thoroughly incorporate to a depth of 6 to 7 inches. <b>NOTE:</b> For bulbous iris, use 5 - 7 pints per 1000 sq.ft. and incorporate to a depth of 2 - 3 inches. Use lower rate for coarse textured (light) soils and higher rate for fine textured (heavy) soils.																										
		$6\frac{3}{4}$ - 9	<b>IN-FURROW APPLICATION AT PLANTING:</b> Apply specified amount to 1000 linear ft. of row in 5 to 10 gals. of water. Direct spray into open furrow to thoroughly cover entire furrow and bulbs. Close furrow in normal manner. Use lower rate for coarse textured (light) soils and higher rate for fine textured (heavy) soils. For Black Rot control, use higher rate.																										
		3	<b>BULB SOAK:</b> Mix specified amount in 3.2 gals. of water (7.5% concentration). Add 1% sticker to mixture and maintain good agitation. Soak bulbs for 5 minutes.																										
		$\frac{3}{4}$	<b>BAND APPLICATION PRIOR TO PLANTING:</b> Apply specified amount to 1000 linear ft. as a 12 inch band spray centered on the row at planting time. Follow with 3 additional applications at 15 day intervals.																										

**USE AND RATE RECOMMENDATIONS FOR ORNAMENTALS**

CROP	DISEASE	PINTS OF TURFCIDE 4F	APPLICATION NOTES											
Gladiolus	Dry Rot ( <i>Stromatinia gladioli</i> ) ( <i>Sclerotinia gladioli</i> )	5½	<b>BROADCAST APPLICATION PRIOR TO PLANTING:</b> Apply specified amount to 1000 sq. ft. in sufficient water to insure uniform ground coverage. Thoroughly incorporate to a depth of 4 to 6 inches. Bulbs may be planted immediately after incorporation.											
		8¼	<b>BAND APPLICATION PRIOR TO PLANTING:</b> Apply specified amount to 1000 linear ft. as a 12 inch band centered on the row. Thoroughly incorporate to a depth of 4 inches. Bulbs may be planted immediately after incorporation.											
Lilies (Easter) (Asiatic) (Oriental)	Root Rot / Stem Rot ( <i>Rhizoctonia solani</i> ) ( <i>Pellicularia filamentosa</i> ) Bulb Rot / Crown Rot ( <i>Sclerotium rolfsii</i> ) ( <i>Pellicularia rolfsii</i> ) Stem Rot / Black Rot ( <i>Sclerotinia sclerotiorum</i> )	5 - 9¼	<b>BROADCAST INCORPORATION PRIOR TO PLANTING:</b> Apply specified amount to 1000 sq. ft. in sufficient water to assure uniform ground coverage. Thoroughly incorporate to a depth of 6 to 7 inches. Use lower rate for coarse textured (light) soils and higher rate for fine textured (heavy) soils.											
		6¾ - 9	<b>IN-FURROW APPLICATION AT PLANTING:</b> Apply specified amount to 1000 linear ft. of row in 5 to 10 gals. of water. Direct spray into open furrow to thoroughly cover entire furrow and bulbs. Close furrow in normal manner. Use lower rate for coarse textured (light) soils and higher rate for fine textured (heavy) soils.											
		6 - 9	<b>BULB SOAK:</b> Mix specified amount in 100 gals. of water. A sticking agent may be added. Maintain good agitation in dip tanks. Soak bulbs in mixture for 5 - 15 minutes.											
Magnolia grandiflora	Magnolia Leaf Spot ( <i>Phyllostica cookeri</i> )	3	<b>FOLIAR SPRAY APPLICATION:</b> Apply specified amount in 100 gallons of water. Add a spreader-sticker at the rate of 1 pint per 100 gallons. Begin applications approximately 1 week before disease normally appears. Apply at least 4 sprays at 2 week intervals. <b>NOTE:</b> Do not use on Magnolia fuscata as injury may result.											
Cut Flowers such as but not limited to: Carnation Chrysanthemum Roses	Storage Rot ( <i>Botrytis cinerea</i> )	1 - 3	<b>DIP APPLICATION:</b> Mix specified amount in 100 gallons of water. Maintain good agitation in dip tanks. Dip prior to storage.											
		3	<b>SPRAY APPLICATION:</b> Mix specified amount in 100 gallons of water. Thoroughly spray prior to storage.											
Southern Pine Seedlings	Root / Stem Rot, Damping Off ( <i>Rhizoctonia spp.</i> ) ( <i>Pellicularia filamentosa</i> )  Needle Blight ( <i>Dothistroma pini</i> )	1¾	<b>BROADCAST APPLICATION PRIOR TO PLANTING:</b> Apply specified amount to 1000 sq. ft. in sufficient water to insure uniform ground coverage. An additional 1/2 inch of water should be applied by irrigation following treatment. Seed may be planted immediately following irrigation or within one week of this date.											
Vegetable Bedding Plants Limited to only container grown Broccoli Brussel Sprouts Cabbage Cauliflower Peppers Tomatoes	Root/Stem Rot, Damping Off ( <i>Rhizoctonia solani</i> ) ( <i>Pellicularia filamentosa</i> )	¾ - ¾ 6 - 12 fl. oz.	<b>SOIL DRENCH APPLICATIONS IN THE NURSERY OR GREENHOUSE:</b> Mix specified amount in 100 gals. of water and apply according to the following guidelines: For Container Grown Plants:											
			<table border="1"> <thead> <tr> <th rowspan="2">Soil Depth inches</th> <th colspan="2">Coverage for TURFCIDE 4F Drench Mix</th> </tr> <tr> <th>Fl. oz./Sq. ft.</th> <th>Sq. ft./100 gals.</th> </tr> </thead> <tbody> <tr> <td>2 or less</td> <td>16</td> <td>800</td> </tr> <tr> <td>3</td> <td>24</td> <td>530</td> </tr> <tr> <td>4 or more</td> <td>32</td> <td>400</td> </tr> </tbody> </table>	Soil Depth inches	Coverage for TURFCIDE 4F Drench Mix		Fl. oz./Sq. ft.	Sq. ft./100 gals.	2 or less	16	800	3	24	530
Soil Depth inches	Coverage for TURFCIDE 4F Drench Mix													
	Fl. oz./Sq. ft.	Sq. ft./100 gals.												
2 or less	16	800												
3	24	530												
4 or more	32	400												

**PLANT TOLERANCE:**

Neither the manufacturer nor the seller has determined whether or not TURFCIDE 4F can be used safely on ornamental plants not specified on this label. Prior to any large scale application on such plants, the user should determine the safety of TURFCIDE 4F fungicide by testing a small number of the type of plants to be treated at the recommended rates for that particular group, i.e., bedding plants, foliage, etc. and observing the treated plants over a one to two week period for symptoms of phytotoxicity, which may occur as foliage burn or stunted growth. The user assumes all risks arising out of application to unlabeled plants.

**USE DIRECTIONS FOR CHEMIGATION OF ORNAMENTALS**

In addition to the above use rates and recommendations, the following precautions must be observed when using this product in any type of irrigation system:  
Apply this product only through the following irrigation systems:  
1) Pressurized drench (flood) or drip (trickle), 2) Micro-irrigation such as spaghetti-tube or individual tube irrigation, 3) Hand-held calibrated irrigation equipment such as the hand-held wand with injector. 4) calibrated overhead watering booms, 5) ebb and flow or bench flooding subirrigation systems.  
Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system, (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

**SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not reuse empty containers. Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep container closed when not in use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:**

**Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**EMERGENCY ASSISTANCE:**

**UNIROYAL CHEMICAL EMERGENCY PHONE 203-723-3670**

**SAFETY DATA AND INFORMATION 203-573-3303**

**TRANSPORTATION EMERGENCY (CHEMTREC) 800-424-9300**

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

®TURFCIDE is a Registered Trademark of Uniroyal Chemical Company, Inc.

©Copyright 1999, Uniroyal Chemical Company, Inc.