

400-461

12/14/2010

10 of 13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Judith Ball
Chemtura USA Corporation
199 Benson Road
Middlebury, Connecticut 06749

Dear Ms. Ball:

DEC 14 2010

Subject: Labeling Amendment; Rate Correction
Dimilin 2L
EPA Registration No. 400-461
Date Submitted: December 13, 2010

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. The following comment applies.

1. Should you wish to retain a reference to the company's website on your label, please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this label, please contact Kable Bo Davis at (703) 306-0415 or davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "John Hebert".

John Hebert
Product Manager (07)
Insecticide-Rodenticide Branch
Registration Division (7505P)


Enclosure- Stamped Label

2013

GROUP 15 INSECTICIDE

Restricted Use Pesticide. Due to toxicity to aquatic invertebrate animals. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Dimilin® 2L



Net Contents:

Insect Growth Regulator Aqueous Flowable

For use on **barley, oats, triticale, wheat, cotton, grassland and non-crop areas, leafy brassica and turnip greens, livestock/poultry premises, oranges, grapefruit, tangerine, peanuts, pears, peppers, rice, soybeans, stonefruit (excluding cherries), tree nuts, and turfgrass**


COMPOSITION

Active Ingredient: (% by weight)	
diflubenzuron	
N-[[[4-Chlorophenyl]amino]carbonyl]-2,6-difluorobenzamide*	22%
Inert Ingredients:	78%
TOTAL	100%

*Contains 2 lbs. diflubenzuron per gallon.
*U.S. Patent Number: 6,057,370; and 6,376,430B1 and other patents pending.

EMERGENCY ASSISTANCE:	
EMERGENCY PHONE	800-292-5898
SAFETY DATA AND INFORMATION	866-430-2775
TRANSPORTATION EMERGENCY (CHEMTREC)	800-424-9300
Have the product container or label with you when calling a doctor or going for treatment.	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**



EPA REG. NO. 400-461
EPA EST. NO.
052A/042910

Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749



www.chemtura.com

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistant selection chart.

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber, natural rubber, polyethylene, PVC, or viton, when mixing and loading and also when using hand-held equipment; shoes plus socks.

Mixers and Loaders Using Fixed-Wing Aircraft Must Wear: A long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, PVC or viton; shoes plus socks; dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C or a NIOSH approved respirator with any R, P or HE filter).

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems (including water soluble bags), enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to terrestrial juvenile insects and aquatic invertebrates/mollusks/insects. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

**DIRECTIONS FOR USE
Restricted Use Pesticide**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.
- shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE - Store in original container only.

PESTICIDE DISPOSAL - Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic containers: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse (or equivalent) promptly after emptying.

Triple rinse as follows: For containers small enough to shake: Empty the remaining contents into a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and then recap. Shake for 10 seconds. Pour rinsate into a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. For containers too large to shake: Empty remaining contents into a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into a mix tank or store for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into a mix tank and continue to drain for 10 seconds after the flow continues to drip. Hold container upside down over mix tank to collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

INSTRUCTIONS AND INFORMATION

Restriction: Do not apply this product through any type of irrigation system.

SPRAY DRIFT LABELING

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination. Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to ULV applications on grassland and non-crop areas, for the control of grasshoppers and Mormon crickets.

The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher-rated flows produce larger droplets.

- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for the displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speed of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evapora-

tion is most severe when conditions are hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

INFORMATION

DIMILIN 2L is an insect growth regulator which is effective on a wide variety of insect pests, predominately from the families Lepidoptera and Diptera. Because of its mode of action, which results in a disruption of the normal molting process of the insect larvae, the action of DIMILIN is slow and several days may elapse before the full effect is seen. Because of its specificity, DIMILIN is an excellent product for use in IPM programs.

RESISTANCE MANAGEMENT: When used as directed DIMILIN 2L provides control of a number of important insect pests as well as providing a margin of safety to beneficial insects and pollinators. DIMILIN 2L should be part of an IPM program that follows good management practices that include:

- Scouting regularly and use DIMILIN 2L against early immature stages for best results
- Always follow the label rate and timing directions
- Use chemical alternatives such as oil and preserve beneficial arthropods as part of an IPM program
- Maintain good coverage of all leaf surfaces with adequate water volume
- Alternate treatments to classes of insecticides with different modes of action

RESTRICTIONS

Do not apply this product to bodies of water where swimming is likely to occur.

For Field Crops, Row Crops, Orchard Uses, Grassland and Non-Crop Areas: Do not apply within 25 feet by ground or 150 feet by air of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries. All applications must include a 25 foot vegetative buffer strip within the buffer zone to decrease runoff.

RESTRICTIONS ON ROTATIONAL CROPS: Do not plant food or feed crops in DIMILIN treated soils within 1 month following last application, unless DIMILIN is authorized for use on these crops.

APPLICATION INSTRUCTIONS

USE AND MIXING DIRECTIONS IF USED WITH WATER:

1. Fill tank with half of the required amount of water.
2. Begin agitation and add required amount of DIMILIN 2L.
3. Continue agitation while adding remainder of water.
4. If permitted for the use site, add proper quantity of oil slowly. To avoid formation of an invert emulsion, use at least 2 parts of water for each part of oil.

USE AND MIXING DIRECTIONS IF USED WITHOUT WATER:

Always evaluate any potential mixture for compatibility and sprayability. To ensure thorough mixing of DIMILIN 2L with insecticides or other carriers, it is recommended that ingredients be premixed in a nurse tank prior to being transferred to aerial or ground ULV application equipment. If nurse tank is not available, or unable to simultaneously mix:

1. Fill tank with the required amount of oil and/or oil based insecticide.
2. Begin agitation and add required amount of DIMILIN 2L.
3. After the contents of the tank have been thoroughly agitated, a volume of carrier sufficient to fill the booms and piping system should be drained and then added back to the tank.

Aerial or ground application: Apply spray with aerial or ground equipment designed or modified to insure full uniform coverage of the entire plant. Adjust equipment to provide droplets with a diameter of 150 to 220 microns. Provide agitation prior to, during, and after blending and while applying.

5013

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Barley Oats Triticale Wheat	BARLEY, OATS, TRITICALE & WHEAT RESTRICTIONS: Do not make more than 1 application per season. Do not exceed 4 fl oz per acre. Do not apply after boot stage of growth. For Use in The Following States Only: Alaska, Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming, Western North & South Dakota and Western Nebraska (West of Route 281 in ND, SD & NE) Pre-harvest Interval: Do not harvest grain and straw within 50 days of application. Do not harvest forage within three days of application. Do not harvest hay within 15 days of application.		
	Grasshopper	1 - 2	For best results, apply when the majority of infesting grasshoppers have reached the 2nd to 3rd nymphal stage of development. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding; a tank mix with a knockdown insecticide is recommended under these conditions.
	Cereal leaf beetle	4	For best results, apply at first sign of egg laying. Do not apply if infestation has advanced into later instar larvae.
	Aerial Application: Apply in 2 to 5 gallons total volume per acre. Ground Application: Apply in 5 to 15 gallons of total volume per acre. Use sufficient application volume to assure adequate coverage. Because of the unique mode of action of DIMILIN 2L, its visible effects on larvae and nymphs may not be seen until 5 to 7 days following application.		
Cotton	COTTON RESTRICTIONS: Do not exceed 6 applications per season. Do not exceed 24 fl oz per acre. Do not exceed 3 applications and 12 fl oz post boll opening. Pre-harvest Interval: Do not harvest within 14 days of application.		
	Beet armyworm -early season before first bloom	2 - 4	For early infestations on young cotton, apply DIMILIN 2L at the first sign of beet armyworm activity (2 egg masses or hatch outs/100 feet of row) in multiple applications, either as directed or broadcast spray. Use on a 5 to 7 day interval until 8 fl oz per acre have been applied. Multiple applications of DIMILIN 2L will provide acceptable beet armyworm control and because it has little activity on beneficial insects (parasites and predators) and has good persistence, will help prevent populations of beet armyworm from building up later in the growing season. Use of DIMILIN 2L in this way allows for more complete coverage of new foliage during the period of rapid vegetative growth.
	Beet armyworm - mid season	4 - 8	Apply starting around first bloom and through mid-bloom. Repeat application until up to 8 fl oz per acre have been applied, using a 5 to 7 day interval between applications. Use higher application rate on larger cotton and/or under conditions of greater larval pressure. Apply first application should coincide with peak beet armyworm moth catches in pheromone traps, indicating another generation of larvae is imminent. DIMILIN 2L is more effective on early stages of larval development, therefore treat cotton leaves should be treated before populations become established.
	Beet armyworm - late season	6 - 8	Apply after mid-bloom and prior to 14 days before harvest. Use higher application rate on larger cotton and/or under conditions of greater larval pressure. Coincide application with peak beet armyworm moth catches in pheromone traps. Additional applications may be needed if larval pressure continues.
	Fall armyworm Yellowstriped armyworm Southern armyworm Soybean looper * Cabbage looper * Saltmarsh caterpillar * *suppression	4-8	Apply during early stages of larval development. Repeat application until at least 8 fl oz per acre have been applied using a 5 to 7 day interval.
	Boll weevil - early season (before first bloom)	4 - 8	DIMILIN 2L will control boll weevil by suppressing reproduction. Apply with 2 to 4 qt of emulsified cottonseed oil, vegetable oil, or paraffinic crop oil. For ULV application, use 4 fl oz in a minimum of 8 fl oz of emulsified cottonseed oil, oil based insecticide, or vegetable or petroleum based oil carrier. A compatibility agent may be needed if a non-emulsified cotton-seed oil is used. Consult your supplier or Chemtura representative for oil specifications. For best suppression of boll weevil reproduction, make first application at pinhead square stage of cotton growth when overwintering boll weevils are entering the fields. Repeat applications must allow a minimum of 7 days between applications. DIMILIN 2L does not kill the adult boll weevil, however, eggs deposited by affected female weevils will not hatch, thus limiting reproduction. The control of egg hatch and larval development within the square prevents its shedding and will then allow normal boll development. After the initial treatment of the female weevil, 7 to 10 days are required before non-hatching eggs are laid; however, once affected, non-hatching eggs will be laid for approximately 10 days, and longer if the female encounters more DIMILIN 2L. Thus treat early and use multiple applications.
	Boll weevil	2 - 4	DIMILIN 2L will reduce the number of weevils that emerge in the following spring if applications are made when adult weevils are going into diapause to overwinter. Apply when cotton plant has reached full vegetative growth or when it begins blooming out the top. For LV application spray in combination with 2 to 4 qt of an emulsifiable vegetable or paraffinic oil per acre. For ULV application combine in a minimum of 8 oz of emulsified cottonseed oil, oil based insecticide, or vegetable or petroleum based oil carrier. A compatibility agent may be needed if a non-emulsified cottonseed oil is used. Apply at least 2, but not more than 3, applications at 7 to 14 day intervals should be made.

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Cotton (cont.)	: Emulsifiable concentrate insecticide formulations used in tank mixes, in the presence of oil, may result in phytotoxicity. Care must be taken where such mixtures are used.		
	Grasshopper	2	Apply when the majority of infesting grasshoppers have reached the 2nd to 3rd nymphal stage of development. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding; a tank mix with a knockdown insecticide is recommended under these conditions.
	<p>Aerial application: Apply in 3 to 5 gallons total volume per acre. For ULV application, use a total volume of 20 to 48 oz per acre.</p> <p>Ground application: Apply in 10 to 20 gallons of total volume per acre. For ULV application, use a total of 20 to 64 oz per acre.</p> <p>Adjuvant usage: Always use oil (1 to 2 qt) with DIMILIN 2L for larval/nymphal control if conditions are favorable for water evaporation (e.g. high air temperature and/or low humidity). For ground or aerial LV application, 1 pt to 2 qt of emulsified vegetable or paraffinic crop oil is recommended to enhance canopy penetration and to reduce spray droplet evaporation and subsequent drift. For ULV application, use DIMILIN 2L in a minimum of 20 oz of emulsified cottonseed, vegetable or petroleum based oil carrier. A compatibility agent may be needed if non-emulsified cottonseed oil is used.</p> <p>Consult your supplier or Chemtura representative for oil specifications.</p> <p>Use sufficient application volume to assure adequate coverage. DIMILIN 2L may be mixed with other insecticides being applied for other cotton insects. When emulsifiable concentrate insecticide formulations are used with oil and DIMILIN 2L in tank mixes, they may result in phytotoxicity. Care must be taken where such mixture is used. Because of the unique mode of action of DIMILIN 2L, its visible effects on larvae/nymphs may not be seen for 5 to 7 days following application.</p>		
Grassland (includes rangeland, pastures, improved pastures and similar areas used for production of native, domesticated forage grasses for harvest for live stock primarily for grazing or mechanical harvest)	GRASSLAND RESTRICTIONS: Do not exceed a total of 2 fl oz per acre per cutting. Do not exceed a total of 6 fl. oz. per acre per year. Allow at least 1 day after treatment before cutting grass. Apply only when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).		
	Grasshopper	1 - 2	Use 1 application on early instar (majority in the 2nd through 4th instar nymphal stages); use high rate for pastureland.
	Mormon cricket	0.75 - 1	Use on rangeland only, in a RAATs (Reduced Area and Agent Treatment) application on early instars. A RAATs application is an IPM strategy that takes advantage of grasshopper movement and conservation biological control to allow DIMILIN 2L to be applied on rangeland on a reduced treated area and at reduced rates, while sustaining acceptable control. RAATs may provide ranchers with an economic means to reduce competition by these insects on their rangeland, depending on insect age and plant canopy. Using this program DIMILIN 2L may be applied on as little as 50% of the infested acreage (e.g. skipping a 100 ft swath for every 100 ft treated), up to 100% infested acreage. The rate range to use per acre and amount of area treated will depend on grasshopper/Mormon cricket age, plant canopy and topography. Skip up to 50% of the infested area and use the lower rate under uniform topography with early instar ages and sparse vegetation. If the majority of the population is late instars, vegetation is dense, terrain is considered rough, and conditions are hot during treatment, increase the coverage and rate of DIMILIN 2L up to a blanket (100%) coverage with 1 fl oz per acre.
		0.5 - 1	If a second application is made, typically apply 2 to 3 weeks after the first application.
	Lepidopteran foliage feeding caterpillars such as: Fall armyworm Striped grass looper	2	For maximum control use DIMILIN 2L at first sign of hatch outs and prior to larvae reaching fourth instars (<1/2 inch). DIMILIN 2L must be ingested and larvae must molt before populations are reduced.
	Horn fly Face fly	2	Apply Dimilin 2L for the control of Horn fly and face fly emergence from cattle manure patties for two weeks or longer.
<p>Aerial application: For low/high volume application, apply in 2 to 10 gallons of water per acre. For rangeland ULV application, apply in a minimum of 12 fl. ozs. total volume per acre.</p> <p>Ground application: For low/high volume application, apply in 2 to 30 gallons of water per acre. For rangeland ULV application, apply in a minimum of 12 fl. ozs. total volume per acre.</p> <p>Regardless of application type, total spray volume used must ensure thorough coverage of the target crop. For aerial and ULV spray mixtures include an evaporation/drift retardant product at use rates prescribed on the specific product label, particularly when conditions are favorable for water evaporation (e.g., high air temperature and/or low humidity). When using oil type evaporation/drift retardant products, be sure to maintain a ratio of at least 2 parts water to 1 part oil. For low volume and ULV applications, make sure that the spray mixture in the boom contains the correct concentration of Dimilin 2L before application begins, and be sure that good agitation is maintained throughout mixing and application.</p> <p>Higher rates and gallonages are suggested for areas with dense vegetation, when nymphs are beyond the 3rd instar stage, and when climatic conditions are favorable for grasshopper/Mormon cricket survival and increase.</p> <p>Apply anytime after eggs begin to hatch through early instars. DIMILIN 2L remains active on the foliage and will continue to control larvae and grasshoppers/Mormon crickets that hatch later in the season. DIMILIN 2L is not effective in controlling larvae and grasshoppers/Mormon crickets once they have reached the adult stage. Since it is an insect growth regulator, effects may not be seen until these insects have molted at least once. If adult grasshoppers/Mormon crickets from early hatching and/or overwintering species are present, tank-mix DIMILIN 2L with a registered adulticide to control later hatching species.</p> <p>Check mixing compatibility and sprayability prior to transferring to the main spray tank.</p> <p>Besides a fatal incomplete molting, adult grasshoppers/Mormon crickets may exhibit missing posterior legs, hernias, abdominal segments malformed, twisted antennae, hemolymph exudation, and wrinkled wings. Additionally, they may move slower, have limited jumps and unsteady landings, show a reduction in feeding, have atrophy of posterior legs or be unable to fly. Any nymph/adult possessing these symptoms is likely more susceptible to predatory insects, birds and mammals.</p>			

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Leafy brassica group includes: Broccoli raab Cabbage Chinese (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens	LEAFY BRASSICA RESTRICTIONS: Do not make more than 4 applications per season. Do not exceed 16 fl oz per acre. Do not use on turnip cultivars or varieties which produce a harvestable root. Pre-harvest Interval: Do not harvest within 7 days of application.		
	Grasshopper	2 - 4	Apply to grasshoppers in the 2nd to 3rd nymphal stage of development. Reapply in 7 day intervals if nymphal hatchout/crop reinfestation continues. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. Use the higher rate in the range if the area has a history of heavy infestations, dense foliage is present, or greater residual control is desired. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding; a tank mix with a knockdown insecticide is recommended under these conditions. Ground Application: Use a minimum of 30 gallons of water per acre to give uniform coverage. Additional applications allow for more complete coverage of newly expanding foliage. Since DIMILIN 2L is an insect growth regulator, larvae and nymphs must ingest treated plant material and then molt before populations are reduced. Thus initial signs of control may not be seen until 5 to 7 days after treatment.
Livestock / Poultry Premises includes- - Litter - Stale / waste feed - Manure - Manure / straw mixtures - Feed muck / spoilage - Spoiled organic refuse - Bedding material - Floors - Walls / wall footings - Posts - Cage frames - Ceilings	RESTRICTIONS: Do not apply directly to livestock or poultry. Do not contaminate feed or water through application- cover or remove exposed feed and water from the area to be treated.		
	Carrion Beetle Darkling Beetle Hide Beetle	12 fl.ozs./ 1000 ft. ² in 2 - 20 gals. water per 1000 ft. ²	Broadcast Application: Apply as a whole house broadcast spray to the litter following de-caking, as well as to floors, walls, posts, cage frames, and cracks and crevices around insulation. When treating the litter, pay particular attention to areas under feed and water lines. Apply in sufficient volume to uniformly and thoroughly wet the litter and other surfaces- spray volume will vary depending on the depth of litter being treated. Band Application: When the whole house is not being treated, application can be made to areas where pests are concentrated, such as under feed and water lines, as well as along perimeter walls and side / end walks. Apply in sufficient volume to thoroughly wet litter following de-caking in a 2-4 foot wide band under and next to these areas- spray volume will vary depending on depth of litter. Lower sections of walls, posts and cage frames should also be treated at least 1 foot up from the floor.
	House fly Stable fly Face Fly Horn Fly	12 fl.ozs./ 1000 ft. ² in 2 - 20 gals. water per 1000 ft. ²	Broadcast Application: Apply as a whole house broadcast spray or spot treatment to the litter following de-caking, as well as to floors, walls, posts, cage frames and ceilings. When treating the litter, pay particular attention to moist areas under feed and water lines. Apply in sufficient volume to uniformly and thoroughly wet the litter and other surfaces- spray volume will vary depending on the depth of litter being treated.
		5 fl.ozs. in 10 gals. water	Spot Treatments: Apply as a directed spray at a volume of 1 quart of spray solution to 10 sq.ft. of surface area. 100 gallons of spray solution will treat 4000 sq.ft. Begin applications when flies first appear. Reapply as needed when adult fly numbers begin to increase, typically at 2- 3 week intervals.
Livestock / poultry operations includes farms, farm buildings, barns, feedlots, dairies, equine facilities, poultry houses, and other production facilities. Application sites within these operations also include fence lines of holding pens, feed troughs, feed bunks, hay bale feeders, water troughs; and marginal areas of waste retention ponds.			
For insect control around hay feeding sites, treat the entire area where manure and waste hay are mixed at the soil surface by livestock activity.			
DIMILIN 2L will not control adult or pupal stages, but does provide extended control of eggs and developing larvae. Exposure to adults, however, through contact or ingestion, does impact their reproductive potential, resulting in reduced numbers and viability of eggs. If a large adult population already exists at the time treatment is to be made, application with a knockdown insecticide either alone or in a tank mix with DIMILIN 2L may be desirable to achieve rapid reduction of that population.			
Non-crop areas [includes field border, fence rows, roadsides, farmsteads, ditchbanks, wasteland, Conservation Reserve Program CRP Land]	NON-CROP AREA RESTRICTIONS: See Grassland section for restrictions.		
	Grasshopper Mormon cricket	2	Apply DIMILIN 2L to manage these insects in their breeding areas before they move into cropland. See Grassland section for timing of application.
	Lepidopteran Foliage feeding caterpillars such as: Fall Armyworms Striped Grass Looper,	2	For maximum control use DIMILIN 2L at first sign of hatch outs and prior to larvae reaching fourth instars (<1/2 inch). DIMILIN 2L must be ingested and larvae must molt before populations are reduced.
See Aerial Application section of Grassland.			
Ground application: Apply in 5 to 30 gallons of water per acre. Include 1 pt to 2 qt of emulsified vegetable or paraffinic crop oil if conditions are favorable for water evaporation.			

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Turfgrass (For use in sod farms only)	Turfgrass Restrictions: Do not exceed a total of 4 applications per year.		
	Lepidopteran Foliage feeding caterpillars such as: Sod webworm Armyworms including Fall, True, Southern, Beet, Yellow-striped, Striped Grass Looper, Granulate Cutworm	2	Apply Dimilin 2L at first sign of hatchouts and prior to larvae reaching 4 th instars (>1/2 inch). Apply in 20 to 50 gallons of water per acre depending on density of turf and caterpillar pressure. Dimilin 2L must be ingested and larvae must molt before populations are reduced. Repeat applications at 14 day intervals or as needed to protect new foliage growth.
Peanuts	PEANUT RESTRICTIONS: Do not make more than 3 applications per season. Do not exceed 24 fl oz per acre. Pre-harvest Interval: Do not harvest within 28 days of application.		
	Velvet bean caterpillar Mexican bean beetle Green cloverworm	2 - 4	Make applications when larvae are small (< 0.5 inches) to give greater control and minimum insect damage to leaves. Repeat application if damaging numbers reappear. The minimum reapplication interval is 14 days. Use the higher rate in the range if the crop has a history of heavy infestations, dense foliage is present, or greater residual control is desired.
	Armyworms, such as: Beet armyworm Fall armyworm Southern armyworm Yellow-striped armyworm Lesser cornstalk borer Soybean looper (suppression)	4 - 8	
	Grasshopper	2	For best results, apply when the majority of infesting grasshoppers have reached the 2nd to 3rd nymphal stage of development. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding. A tank mix with a knockdown insecticide is recommended under these conditions.
	Aerial Application: Apply in sufficient water (3 to 5 gallons per acre) to achieve uniform coverage of foliage. Ground Application: Apply in 9 to 35 gallons of water per acre to give uniform coverage. Adjuvant Usage: See Cotton section. Since Dimilin 2L is an insect growth regulator, larvae/nymphs must ingest treated plant material and then molt before populations are reduced. Thus initial signs of control may not be seen until 5 to 7 days after treatment.		
Oranges Grapefruit Tangerine Pummelo and their hybrids	CITRUS RESTRICTIONS: Do not apply more than 60 fl. oz. of Dimilin 2L per acre per year. Do not apply within 21 days of harvest. Do not harvest cover crops for animal feed or graze livestock in treated groves. In the State of Florida, do not apply within 100 feet of estuarine/marine bodies of water by ground application or 1000 feet by air application. Spray last three rows windward of surface water using nozzles on one side only, with spray directed away from surface water. Prevent spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove when spraying the outside row. Shut off nozzles when turning at ends of rows and passing tree gaps in rows.		
	Citrus Leafminer (<i>phyllocnistis citrella</i>)	20	Apply when oviposition begins on new growth flush. Use sufficient spray volume for thorough coverage of leaf surfaces (ground = 50 to 1,000 gallons per acre; aerial = 5 to 20 gallons per acre). Repeat application no closer than 90 days apart for subsequent leaf flushes. The addition of a spray oil, such as FC435-66, enhances coverage and may enhance control of citrus leafminers. Dimilin 2L will not kill adult stages of leafminers. Dimilin 2L has activity on eggs, larval and pupal stages.
	Citrus Peelminer (<i>marmara spp.</i>)	20	Apply when oviposition begins on peel surface. Use sufficient spray volume for thorough coverage of leaf surfaces (ground = 50 to 1,000 gallons per acre; aerial = 5 to 20 gallons per acre). The addition of a spray oil, such as FC435-66, enhances coverage and may enhance control of peelminers. Dimilin 2L prevents development of peelminer eggs laid on protected fruit tissues. Protection may last only a few weeks when new tissue is exposed on rapidly expanding fruit.
	Citrus Root Weevil Complex West Indian Sugarcane rootstock borer weevil (<i>Diaprepes abbreviatus</i>) Southern bluegreen citrus root weevil (<i>Pachnaeus litus</i>) Bluegreen citrus weevil (<i>Pachnaeus opalus</i>) Fuller rose beetle (<i>Asynonychus godmani</i>) Little leaf notcher (<i>Artipus fl oridanus</i>)	20	Apply Dimilin 2L to newly expanded flush on citrus and/or when adult weevils are present. Use sufficient spray volume for thorough coverage of leaf surfaces (ground - 50 to 1,000 gallons per acre; aerial = 5 to 20 gallons per acre). Repeat application no closer than 90 days apart for subsequent leaf flushes and/or when adult weevils are present. The addition of a spray oil, such as FC435-66, enhances coverage and penetration of Dimilin 2L into the adult weevils and eggs. Also oil will deter attachment of weevil egg masses to leaf surfaces. Dimilin 2L will not kill adult weevils. The activity of Dimilin 2L is through ingestion or contact and will result in reduction of the reproductive potential of weevils, it prevents eggs from hatching, thus preventing larvae from entering soil and feeding on citrus tree roots. Also, the grubs from eggs laid on treated leaves are reduced in number.

9013

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Orange (cont.)	Citrus rust mite	20	Use in sufficient water to ensure thorough coverage (50 to 1,000 gallons per acre by ground application; aerial=5 to 20 gallons per acre). Repeat application no closer than 90 days apart to maintain full season rust mite control. Dimilin 2L has activity on eggs and nymphal stages of citrus rust mites. Adults that have passed all molting stages are not susceptible to Dimilin 2L. Due to the unique mode of action of Dimilin 2L the full effect of the treatment may not be apparent until 3 to 10 days after application.
Pear	PEAR RESTRICTIONS: Do not apply more than 4 applications per year. Do not apply more than 64 fl oz per acre per year. Pre-harvest Interval: Do not harvest within 14 days of application. Do not use oil in tank mix in late season treatments (3rd and 4th applications).		
	Pear psylla (pre-bloom)	40 - 48	Apply in 80 to 400 gallons of water per acre during the delayed dormant to the popcorn stage period. Complete uniform coverage of the tree is essential to achieve insect control. A horticultural mineral oil should be used at a rate of 4 to 6 gallons per acre during the delayed dormant period. After this period and through the popcorn stage, apply oil at a concentration of 0.25%, but use no more than 1 gallon per acre. A surfactant may be used to improve coverage. Follow manufacturer's label specifications. DIMILIN 2L should be applied during egg deposition so that it will come in contact with pear psylla eggs and/or 1st and 2nd instar nymphs.
	Pear psylla (post-bloom)	12 - 16	Applications at normal codling moth rates and timings will provide suppression of pear psylla.
	Pear rust mite (pre-bloom)	40 - 48	Apply in 80 to 400 gallons of water per acre from delayed dormant to the popcorn stage. See 'Pear psylla (pre-bloom)' for the use of oil.
	Codling moth	12 - 16	Apply in a minimum of 80 gallons of water per acre. Use the lower rate where there is light codling moth pressure and/or on small trees. Complete coverage of the fruit and foliage in all areas of the trees is essential for insect control. Timing of application is extremely important because DIMILIN 2L controls codling moth by prohibiting the hatching of eggs. It must be applied prior to egg laying so that eggs are laid on treated plant parts. Apply first application as soon as possible after first moths are caught (biofix) or observed, or about 50-75 degree-days after biofix. This timing can be determined by your local pest control consultant and/or fruit specialist with the aid of pheromone traps. Normally this timing occurs at late petal fall or about 10-14 days earlier than the timing used for organophosphate insecticides. Apply second application about 14-18 days after the first. If necessary, apply third and fourth application, timed prior to egg laying of the 2nd generation by using the same method as for the 1st generation. If traps are not used, make the 3rd application 21-30 days after the second, followed by the 4th application 21-30 days later. If a degree-day model is used the 3rd spray should be timed at 1000 degree-days after biofix. Combination with organophosphates for codling moth control: DIMILIN 2L can be used in combination with an organophosphate insecticide, to save a trip through the orchard and to make timing of the DIMILIN 2L sprays easier. The combination is more effective than DIMILIN 2L alone when controlling moderate to heavy codling moth infestations and/or treating large trees. The combination will provide residual control of eggs laid after application. Apply DIMILIN 2L and the organophosphates at their labeled rates. Apply at the beginning of egg hatch of 1st generation codling moth. This is the normal timing for the first organophosphate cover spray (250 degree-days following biofix for 1st generation and 1250 degree days for the 2nd generation). This program can be repeated for the 2nd or 3rd generation of codling moth or use DIMILIN 2L alone prior to egg laying. Do not use oil in tank mix with DIMILIN 2L in late season treatments. With light codling moth populations, as indicated by monitoring, this combination may offer control of an entire generation with 1 application. When populations are heavy, this combination will improve control, but it may not control an entire generation with one spray. A second spray of DIMILIN 2L alone or in combination may be applied 14-18 days later.
Leafminer	8 - 16	Apply in a minimum of 80 gallons of water just prior or during egg laying to control eggs and larvae. Timing for control of the 1st or 2nd generation can be determined by your local pest control consultant or fruit specialist. Should later generations of leafminers occur, apply DIMILIN 2L in the same manner. It is desirable to have DIMILIN 2L in place at the time of egg laying. It will continue to give control through the early sap feeding stage. Complete coverage of the foliage is essential to achieve control of the larvae through the early sap feeding stage.	
Oil may cause injury to certain pear varieties. Check compatibility of oil mixtures with your local tree fruit specialist.			

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Pepper Bell and Non Bell	PEPPER RESTRICTIONS: Up to five applications per growing season may be made as long as 24 fl oz. per acre, per season are not exceeded. Allow a minimum of seven days between any two applications. Do not apply within seven days of harvest. Do not apply more than 24 fl oz. per acre per season.		
	Beet armyworm Fall armyworm Southern armyworm and other foliage feeding Lepidopteran insects	4 - 8	Make initial application of 4 to 8 fl oz. DIMILIN 2L per acre when larvae are small to give greater control and minimum damage to leaves and/or to fruit. Use a higher rate if being applied alone and/or infestation is considered heavy. A knockdown tank-mix partner should be used if late instar larvae are present. Use a minimum of 30 gallons of water per acre to give uniform coverage. Additional applications allow for more complete coverage of new foliage and expanding fruit.
	Pepper weevil	4 - 8	Apply DIMILIN 2L at 4 to 8 fl oz. per acre starting at initial flowering. Use at the higher rate if adult infestation is considered moderate to heavy. Apply additional applications at 7 day intervals up to 7 days before harvest. Additional applications allow for more complete coverage of new foliage and expanding fruit. Note that DIMILIN 2L will not control adults; however eggs laid by adults will exhibit reduced hatching in fruits once adults have consumed or contacted residues of DIMILIN 2L on pepper tissue.
	Aerial application: Apply in sufficient water (3 to 10 gallons per acre) to achieve uniform coverage of foliage. Ground application: Use a minimum of 30 gallons of water per acre to give uniform coverage. Adjuvant Usage: See Cotton Section. Since DIMILIN 2L is an insect growth regulator, larvae and nymphs must ingest treated plant material and then molt before populations are reduced. Thus initial signs of control may not be seen until 5 to 7 days after treatment.		
Rice	RICE RESTRICTIONS: Pre-harvest Interval: Do not harvest within 80 days of application. Do not use on rice fields in which crayfish (crawfish) farming is included in the cultural practice. Do not drain treated water into fields where crayfish farming is intended Do not apply to rice immediately adjacent to sites of crayfish aquaculture. Do not use treated rice flood waters for irrigated crops except for uses currently established for Dimilin 2L. Do not impregnate on granular materials. Do not use on wild rice (<i>Zizania spp.</i>).		
	Rice water weevil (Southern U.S. Rice Belt) -for drill seeded; dry seeded; or water seeded, delayed flood rice	12 - 16	Make a single application of DIMILIN 2L per acre per year to control larvae when adult infestations reach economic threshold and/or at initial oviposition, usually within a time frame of 2-5 days after permanent flood establishment. If adult weevil infestations are historically high and/or migration into the field is prolonged, use the higher application rate.
	Rice water weevil (Southern U.S. Rice Belt) water seeded, pinpoint flood, or continuous flood rice	8 + 8	To control larvae, apply split applications. Apply 8 fl oz per acre after the permanent flood when adult infestations reach economic threshold and/or at initial oviposition, usually when rice leaves are exposed above the water surface. The 2nd 8 fl oz treatment must be made 5-7 days after the 1st application. Failure to make the second application within the above timeframe could result in inadequate control of rice water weevil larvae, especially if adult infestations are high and/or migration into the field is prolonged.
	Rice water weevil (California)	8 - 16	To control larvae apply DIMILIN 2L once per year at initiation of oviposition by adults. During a typical year this coincides with 2 to 8 days after rice emergence above the water. Target the application for 2 to 5 days after rice emergence above the water (2 to 4 leaf stage). Use 12 to 16 fl oz DIMILIN 2L if infestations have been historically high.
Consult your local extension service for determination of economic threshold and/or determination of oviposition. Dimilin 2L does not appear to control adult weevils. It controls rice water weevil by preventing larval emergence from the egg. Eggs laid under the surface of treated water are controlled. Additionally, adults feeding on treated plant surfaces do not lay viable eggs. Apply DIMILIN 2L by air using at least 5 gallons total volume per acre. Do not apply DIMILIN 2L if flooding is in progress. Activity will be reduced. Since DIMILIN 2L is water active, the entire field must be treated. For maximum activity of DIMILIN 2L do not disturb flood after a single application for at least 7 days. With split applications in water seeded, pinpoint or continuous flood rice, flood must not be disturbed for a minimum of 4 days following the 1st treatment and 7 days following the 2nd application. Hold treated water at least 14 days to allow for dissipation of DIMILIN 2L. DIMILIN 2L is not phytotoxic to rice. DIMILIN 2L can be safely applied in combination with post permanent flood herbicides such as FACET®, GRANDSTAND® and LONDAX®. However, before using a tank-mix combination, read each product label carefully and follow Precautionary Statements on each label. ®Facet is a registered trademark of BASF AG; ®Grandstand is a registered trademark of Dow AgroSciences; ®Londax is a registered trademark of E.I. DuPont de Nemours and Company.			

110A13

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Soybean (Except California)	SOYBEAN RESTRICTIONS: Use on soybeans not registered by the California Department of Pesticide Regulation. Do not make more than 2 applications per season. Pre-Harvest Interval: Do not harvest within 21 days of application.		
	Velvet bean caterpillar Mexican bean beetle Green cloverworm	2-4	Make applications when larvae are small (< 0.5 inches) to give greater control and minimum insect damage to leaves. Repeat application if damaging numbers reappear. The minimum reapplication interval is 30 days. DIMILIN 2L may be applied at the lower rate (2 fl oz) to prevent velvetbean caterpillar build-up when the vegetative growth of soybeans is completed and as pod formation begins. Consult local Extension Service regarding infestation levels requiring treatment.
	Beet armyworm Soybean looper (suppression)	4	Application must be made when worms are small before populations build.
	Grasshopper	2	Apply when the majority of infesting grasshoppers have reached the 2nd to 3rd nymphal stage of development. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding; a tank mix with a knockdown insecticide is recommended under these conditions.
Aerial application: apply in sufficient water (3 to 5 gallons per acre) to achieve uniform coverage of foliage. Ground application: apply in 9 to 35 gallons of water per acre to give uniform coverage. Adjuvant usage: See Cotton Section. Since DIMILIN 2L is an insect growth regulator, larvae/nymphs must feed on it and then molt before populations are reduced. Thus initial signs of control may not be seen until several days after treatment. Soybean yield enhancement: In the absence of significant insect pressure and under certain growing conditions, an increase in soybean seed yield has been demonstrated with DIMILIN 2L under field conditions on both determinate and indeterminate cultivars. Application of 2 to 4 fl oz per acre to high yield potential soybean plants at the R3 to R3.5 growth stage period has been more consistent in increasing yields than applications at other reproductive stages of the soybean plant. This reproductive period represents beginning pod growth (pod 3/16 inch long at one of the uppermost nodes on the main stem with a fully developed leaf) to just prior to full pod elongation (pod 3/4 inch long at one of the 4 uppermost nodes on the main stem with a fully developed leaf).			
Stonefruit (excluding cherries) includes: apricot nectarine peach plum prune	STONEFRUIT RESTRICTIONS: Do not apply after petal fall. Do not exceed 2 applications in any given season. Do not exceed 0.50 lb ai (32 fl oz) per acre in any given season. Allow at least 21 days between applications.		
	Peach twig borer	12 - 16	Dormant/delayed dormant: Apply DIMILIN 2L with 4 to 6 gallons per acre (1.5 to 2.0 gallons per 100 gallons in a dilute spray) narrow range oil. Always use the higher rate of DIMILIN 2L if the crop has a history of heavy infestations. Bloom: Apply starting at early bloom. Vegetable oil may be used at the rate of 1 qt per acre. Always use the higher rate in the range if the crop has a history of heavy infestations.
Fall webworm Filbert leafroller Oblique banded leafroller Omnivorous leafroller Omnivorous leaf-tier Oriental fruit moth Redhumped caterpillar Variegated leafroller Walnut caterpillar Winter moth		8 - 16	Apply Dimilin 2L at first sign of larval infestation. Use the higher rate for longer residual control, higher pest infestations, low crop load, larger trees or heavy, dense foliage.
Ground applications must be made in sufficient water for thorough coverage, using at least 50 gallons per acre for small trees (10 feet tall) and at least 100 gallons per acre for larger trees. Using insufficient water for thorough coverage and/or using an uneven spray pattern across the canopy will likely result in less than desired efficacy.			

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
Tree nuts group includes: almond beech nut Brazil nut butternut chestnut chinquapin filbert (hazelnut) hickory nut macadamia nut (bush nut) pecan pistachio walnut (black & English)	TREE NUT RESTRICTIONS: Pre-harvest Interval: Do not harvest within 28 days of application. Do not exceed 4 (3 for walnuts) applications. Do not exceed 64 fl. oz. per acre per growing season.		
	Codling moth	16	DIMILIN 2L is most effective when applied prior to egg laying. DIMILIN 2L must be present on the surface upon which eggs are laid; therefore, full coverage spray is necessary. Apply first application when moth flights begin or when moths are found in pheromone traps. Apply the 2nd application approximately 21 days after the 1st application. For control of the 2nd brood, application should be timed prior to egg laying, similar to 1st brood. Because of fluctuations in temperature, the emergence and moth flights of the over-wintering population may be extended over a long period of time. Under such circumstances, DIMILIN 2L should be tank mixed with an organophosphate insecticide at its lowest label rate. This tank mix should be applied at normal 1st organophosphate timing. Later in the season, if egg laying has already occurred before application of DIMILIN 2L, it is recommended that DIMILIN 2L be tank mixed with an organophosphate as previously described.
	Filbert worm	12 - 16	The lower rate may be used where filbert worm pressure is low and/or the trees are small. The higher rate is necessary when worm pressure is moderate to high and/or the trees are large. Apply DIMILIN 2L 2 to 3 days after the 1st moth is caught in pheromone detection traps. Mating takes place within several days of emergence and egg laying begins the next day. DIMILIN 2L must be applied prior to egg deposition on the treated foliage. Good uniform coverage of the tree is essential to achieve optimum control of filbert worm with DIMILIN 2L. Normally DIMILIN 2L will give season long control. If moth pressure remains high, additional applications should be made.
	Hickory shuckworm	8 - 16	Apply DIMILIN 2L starting at half-shell hardening. Make subsequent applications at 21-day intervals to shuck split, or while nuts are susceptible to hickory shuckworm under heavy infestations. Use the higher rate under higher pest infestations, low crop load, larger trees or heavy, dense foliage.
	Peach twig borer	12 - 16	Dormant/delayed dormant: Apply DIMILIN 2L at the rate of 12 to 16 fl oz per acre with 4 to 8 gallons per acre (1.5 to 2.0 gallons per 100 gallons in a dilute spray) narrow range oil. Always use the higher rate of DIMILIN 2L in the rate range if the crop has a history of heavy infestations. Bloom: Apply DIMILIN 2L at the rate of 12 to 16 fl oz per acre starting at early bloom. Always use the higher rate of DIMILIN 2L in the rate range if the crop has a history of heavy infestations. Spring flight ("May Spray"): Using pheromone traps to determine flight activity, apply DIMILIN 2L at the rate of 16 fl oz per acre at initial flight activity. Summer flight: Using pheromone traps to determine flight activity, apply DIMILIN 2L at the rate of 16 fl oz per acre at initial flight activity.
	Pecan nut case-bearer	8 - 16	Apply DIMILIN 2L at the initiation of each adult generation to target egg hatch. Note for the 1st generation this is approximately 8 to 15 days following the first prolonged moth catch (biofix which is defined as the date on which the total of 5 moths are captured in 3 pheromone traps within a 7-day period). States may have a different recommendation for initiation of spraying; please consult authorities such as county and university extension specialists on current recommendations. Use the higher rate for longer residual control, higher pest infestations, low crop load, larger trees or heavy, dense foliage.
	Pecan weevil (suppression)	8 - 16	Use the higher rate if weevils are attacking fruit and for higher infestations.
	Others, including: Fall webworm Filbert leafroller Oblique banded leafroller Omniverous leafroller Omniverous leaf-tier Oriental fruit moth Redhumped caterpillar Variegated leafroller Walnut caterpillar Winter moth	8 - 16	Apply DIMILIN 2L at the first sign of larval infestations. Use the higher rate for longer residual control, higher pest infestations, low crop load, larger trees or heavy, dense foliage.
Apply ground applications in sufficient water for thorough coverage, using at least 50 gallons per acre for small trees (10 feet tall) and at least 100 to 300 gallons per acre for larger trees. Using insufficient water for thorough coverage and/or using an uneven spray pattern across the canopy will likely result in less than desired efficacy. If 4 applications are used, application timing should correspond to dormant to pre-bud swell, at bloom to petal fall, at flowers/leaves/immature nut fruit formation and at hull split.			

13 of 13

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and to the extent consistent with applicable law, the buyer assumes the risk of any such use.

®DIMILIN is a Registered Trademark of Chemtura Corporation
©Copyright 2010, Chemtura Corporation