

400-503

3-25-2008

✓



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

R. Tim Weiland, Ph.D.  
Chemtura Corporation  
199 Benson Road, 2-5  
Middlebury, CT 06749  
USA

March 25, 2008

Dear Dr. Weiland,

Subject: Label Amendments: Adding caneberry subgroup 13A, wild raspberry and several tropical fruits.  
Acramite-50 WS  
EPA Registration No. 400-503  
Submission Date: Nov. 7, 2006


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

- 1- As a condition of registration, Chemtura Corporation is required to submit crop field trial data for the representative commodities of edible-podded legume vegetables, subgroup 6a, (9 trials) and succulent-shelled pea and bean, subgroup 6b, (12 trials). The field trials should be conducted according to the proposed use directions of this petition (not at an exaggerated rate) and residue decline data should be generated. These field trial data must be submitted to the Agency by March 25, 2010.
- 2- Add the statement "Wear protective eyewear" after the statement "Causes moderate eye irritation" under the "Precautionary Statements" section.
- 3- Add "Protective eyewear" to the Personal Protective Equipment (PPE) section. The new sentence should read: "Applicators and Other Handlers Must Wear: Protective eyewear, long-sleeved shirt & long pants; shoes plus socks. Follow manufacturer's....."
- 4- Change the subtitle "General information" on page 2 of the label to "Use information".
- 5- Change the subtitle "Use Rates and Recommendations" on page 2 of the label to "Use Rates and Directions".
- 6- Replace the word "should" with "must" under the subtitle "Use Rates and Directions" on page 2 of the label. The new sentence should read: "To provide maximum residue control, application must be made....."

2/7

A stamped copy of the label is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this letter, please contact Amer Al-Mudallal at (703) 605-0566.

Sincerely yours,


A handwritten signature in black ink, appearing to read "John Hebert". The signature is fluid and cursive, with a large initial "J" and a long horizontal stroke at the end.

John Hebert  
Product Manager (07)  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

3/7

GROUP 25 ACARICIDE

# Acramite®-50WS



**Miticide in Water Soluble Bags**  
For agricultural use only

Net Contents:

Active Ingredient: (% by weight)

bifenazate: hydrazine carboxylic acid, 2-(4-methoxy-[1,1'-biphenyl]-3-yl) 1-methylethyl ester .....	50%
Inert Ingredients .....	50%
Total .....	100%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID	
<b>IF IN EYES:</b> <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lens, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call the following telephone numbers for emergency medical treatment.	
<b>EMERGENCY PHONE</b>	<b>800-292-5898</b>
<b>SAFETY DATA AND INFORMATION</b>	<b>866-430-2775</b>
<b>TRANSPORTATION EMERGENCY (CHEMTREC)</b>	<b>800-424-9300</b>

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**MAR 25 2008**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, you are prohibited from the sale, distribution, or use of any pesticide which is found to be violative of FIFRA Act.

400-503

EPA REG. NO. 400-503  
EPA EST. NO. 33967-NJ-1  
015A

Chemtura Corporation  
199 Benson Road  
Middlebury, CT 06749



www.chemtura.com

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum or using tobacco.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

**Applicators and Other Handlers Must Wear:** Long-sleeved shirt & long pants; shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds, estuarine/marine invertebrates and fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). **The REI and exceptions are listed under the USE RATES AND RECOMMENDATIONS associated with each crop.** Notify workers of the exception (including when entry is permitted for each of the tasks named in the exception).

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; shoes plus socks; chemical-resistant gloves made of any waterproof material.

**GENERAL INFORMATION**

ACRAMITE®-50WS is a wettable powder in water soluble bags. ACRAMITE-50WS is a selective miticide for the control of a variety of mite pests on the crops listed on this label. When used as directed and applied to the foliage, it provides quick knockdown through contact activity, and long residual control. Due to its unique chemistry, mode of action and selective nature, ACRAMITE-50WS is relatively inactive against beneficial/predaceous mites and insects and therefore is compatible with IPM and resistance management programs.

**ACRAMITE-50WS is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.**

**MIXING INSTRUCTIONS**

Always reseal the outer bag in a manner that protects the remaining packets from moisture. Fill spray tank with 1/2 the desired amount of water. Then add the required number of water soluble bags of ACRAMITE-50WS with agitation running to fully disperse the product. Then fill the tank with the remaining amount of required water. When tank mixing, thoroughly mix the water soluble bags before adding other products in the following order: other water soluble bags, wettable powders, dry flowables, liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to disperse fully before adding the next product. Do not add products that release free chlorine or contains boron with water soluble bag formulations. Boron and free chlorine will inhibit the solubility of the water soluble bag material causing it to precipitate and form insoluble residue inside the spray tank.

Like many pesticides, ACRAMITE-50WS stability can be impacted by high pH and high temperature. For optimum performance, maintain spray mixtures containing ACRAMITE-50WS within a range of pH 5.5 to 6.5.

**Restrictions:** Do not remove the water soluble bags from the container except for immediate use. Use the entire contents of one water soluble bag; do not break open to use partial contents of a bag. Do not sell individual water soluble packets. Do not handle the inner bag with wet hands or wet gloves. Do not allow pouches to become wet prior to adding to the spray tank. Tank mixtures are permitted only in those states where the tank mix partner is registered. When tank mixing, follow the label directions for most restrictive of label precautions and limitations.

**Compatibility:** To obtain broad spectrum insect control, ACRAMITE-50WS can be tank-mixed with other insecticide products. However, due to variations in water quality, e.g., hardness and pH, it is required that users conduct small scale trials under local conditions to ensure compatibility prior to any large scale use.

**USE RATES AND RECOMMENDATIONS**

To provide maximum residual control, application should be made as soon as mites appear. Use the low rate where mite infestations are light. The higher rate may be required for heavy infestations or for extended residual control.

When used as directed, ACRAMITE-50WS is effective for the control of a variety of mites species, especially spider mites, red mites and grass mites. **NOTE:** It is not effective against rust mites, broad mites and flat mites. ACRAMITE-50WS is primarily active on the motile stage of mites, but also has ovicidal activity against spider mites (*Tetranychus* species).

**Restrictions:**

- Do not plant another crop within 30 days after last ACRAMITE application due to chances of bifentazate residues showing up in rotational crops.
- ACRAMITE-50WS should not be mixed with oil when applying to Golden Delicious apples.

**MITES CONTROLLED**

Banks grass  
Brown almond  
Clover  
European red (use maximum rate)

McDaniel  
Pacific spider  
Pecan leaf scorch  
Strawberry spider

Two-spotted spider  
Willamette

**BEARING CROPS  
DOSAGE INSTRUCTIONS**

<b>CROP*</b>	<b>AMOUNT ACRAMITE-50WS PER ACRE<sup>(A)</sup></b>	<b>MINIMUM GALLONS PER ACRE GROUND APPLICATION</b>	<b>TOTAL NUMBER OF SPRAYS PER SEASON</b>	<b>HARVEST DAYS AFTER APPLICATION (PHI DAYS)</b>	<b>RESTRICTED ENTRY INTERVAL (REI)</b>
CANEBERRY SUBGROUP 13A, <sup>(B)</sup> WILD RASPBERRY	0.75 - 1.0 lbs	50	1	1	12 hours
CUCURBIT VEGETABLES (See footnote C for crops in addition to those listed below) CUCUMBERS EDIBLE GOURDS <sup>(D)</sup> MUSKMELON <sup>(E)</sup> PUMPKIN SQUASH (summer <sup>(F)</sup> & winter) <sup>(G)</sup> WATERMELON	0.75 - 1.0 lbs	50	1	3	12 hours
FRUITING VEGETABLES (See footnote H for crops in addition to those listed below) EGGPLANT PEPPERS (all varieties of <i>capsicum</i> spp.) <sup>(H)</sup> TOMATOES (field grown)	0.75 - 1.0 lbs	50	1	3	12 hours
GRAPES <sup>(I)</sup>	0.75 - 1.0 lbs	50	1	14	12 hours except 5 days for cane turning, tying and girdling of table grapes.
HOPS	0.75 - 1.5 lbs	50	1	14	12 hours
MINT	0.75 - 1.5 lbs	50	1	7	12 hours
OKRA	0.75 - 1.0 lbs	50	1	3	12 hours
POME FRUITS: (See footnote K for crops in addition to those listed below) APPLES CRABAPPLES PEARS QUINCE	0.75 - 1.0 lbs	50	1	7	12 hours
STRAWBERRIES <sup>(J)</sup>	0.75 - 1.0 lbs	100	2	1	12 hours
STONE FRUIT: (See footnote M for crops in addition to those listed below) APRICOTS CHERRIES (Sweet & Tart) NECTARINES PEACHES PLUMS/PRUNES	0.75 - 1.0 lbs	50	1	3	12 hours
TREE NUTS (See footnote N for crops in addition to those listed below) ALMOND	0.75 - 1.0 lbs	50	1	7	12 hours
FILBERT (Hazelnut) PISTACHIOS WALNUT (Black & English)	0.75 - 1.0 lbs	50	1	14	12 hours
TROPICAL FRUIT: PAPAYA, STAR APPLE, BLACK SAPOTE, MANGO, SAPODILLA, CANISTEL, MAMEY SAPOTE, LYCHEE, LONGAN, SPANISH LIME, RAMBUTAN, PULASAN, GUAVA, FEIJOA, JABOTICABA, WAX JAMBU, STARFRUIT, PASSIONFRUIT, ACEROLA	0.75 - 1.0 lbs	50	1	1	12 hours

\*Use the higher rate under heavy mite pressure.

- (A) Each water soluble bag contains 0.5 lbs. of ACRAMITE-50WS
- (B) CANEBERRY: Blackberry; loganberry; red and black raspberry; cultivars and/or hybrids of these.
- (C) CUCURBIT VEGETABLES: Other crops which may be treated with ACRAMITE-50WS with the same use instructions are: Chayote (fruit) (*Sechium edule*), Chinese waxgourd (Chinese preserving melon) (*Benincasa hispida*), Citrus melon (*Citrullus lanatus var. citroides*), Gherkin (*Cucumis anguria*), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber).
- (D) EDIBLE GOURD (*Lagenaria* spp.) (includes *hyotan* and *cucuzza*). Also (*Luffa acutangula* and *L. cylindrical*) which includes hechima and Chinese okra.
- (E) MUSKMELON (hybrids and/or cultivars of *Cucumis melo*) includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon.
- (F) SQUASH, SUMMER (*Cucurbita pepo var. melopepo*) includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini.
- (G) SQUASH, WINTER (*Cucurbita maxima: C. moschata*) includes butternut squash, calabaza, hubbard squash; (*C. mixta; C. pepo*) includes acorn squash, spaghetti squash.
- (H) FRUITING VEGETABLES: Other crops which may be treated with ACRAMITE-50WS with the same instructions are: groundcherry (*Physalis* spp.), pepino (*Solarium muricatum*), tomatillo (*Physalis ixocarpa*)
- (I) PEPPER (*Capacium* spp.) includes bell peppers, chili peppers, cooking peppers, pimento, and sweet peppers.
- (J) GRAPES — Exception: In addition to the early entry exceptions allowed by WPS, you may enter or allow workers to enter treated areas to perform all tasks other than tying, turning and girdling after 12 hours following applications, as long as at least long pants, long-sleeved shirts, shoes and socks are worn. Notify workers of the exception (including when entry is permitted for each of the tasks named in the exception).
- (K) POME FRUIT: Other crops which may be treated with ACRAMITE-50WS with the same instructions are: Loquat (*Eriobotrya japonica*), Mayhaw (*Crataegus aestivalis, C. opaca, and C. rufula*), Pear, oriental (*Pyrus pyrifolia*)
- (L) STRAWBERRIES: Minimum period between applications is 21 days.
- (M) STONE FRUIT: Other crops which may be treated with ACRAMITE-50WS with the same instructions are: Plum (*Prunus domestica, Prunus* spp.), Chickasaw Plum (*Prunus angustifolia*), Damson Plum (*Prunus domestica* spp.) (*Instititia*), Japanese Plum (*Prunus salicina*) and Plumcot (*Prunus armeniaca X P. domestica*).
- (N) TREE NUTS: Other crops which may be treated with ACRAMITE-50WS with the same use instructions (14 day PHI) are: Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Hickory nut, Macadamia nut (bush nut).

**NON-BEARING CROPS  
DOSAGE INSTRUCTION**

ACRAMITE-50WS use on non-bearing crops, i.e. those crops which will not bear fruit within one year of application, **includes all the crops listed in the bearing crops dosage instructions plus crops listed below.** These include fruit trees and berries in both commercial plantings and nurseries, including transplant uses but excluding residential areas. **The listed non-bearing crops all have a 12 hour restricted entry interval (REI).** Application is by ground equipment only. Additional non-bearing crops that may be so treated are:

CROP	MITES CONTROLLED	AMOUNT ACRAMITE-50WS PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR
Berries (boysenberries, raspberries, etc.) Citrus (grapefruit, lemons, limes, oranges, tangerines, etc.) Currants Dates, Figs Persimmons	European red Two-spotted spider Pacific spider McDaniel Strawberry spider Citrus red Southern red	0.75 to 1.0 lbs.	50	1

7/7

### STRATEGIES FOR RESISTANCE PREVENTION

When used as directed, ACRAMITE-50WS combines high activity on mites with safety to beneficial/predaceous mites and insects. In addition, the unique chemistry of ACRAMITE-50WS provides a means of controlling mites which have developed resistance to commonly used products. These properties can result in fewer miticide/insecticide applications as well as general reduction in the problems caused by resistance.

ACRAMITE-50WS has demonstrated no cross resistance with other commercial miticides. ACRAMITE-50WS contains an active ingredient with a unique mode of action and is classified as a Group 25 acaricide. ACRAMITE-50WS is suitable to be used as a rotational partner with other miticides.

The following mite control strategies are recommended:

- Incorporate IPM techniques into your insect control program.
- Ensure thorough spray coverage to all foliage.
- Scout regularly and apply ACRAMITE-50WS as soon as infestations are observed. Do not wait until large populations have established.
- Always apply ACRAMITE-50WS at the recommended rates and according to label recommendations.
- Unless labeled otherwise, use only one application of ACRAMITE-50WS per year, and rotate to a product with a different mode of action grouping.
- Because of its selectivity, ACRAMITE-50WS can be used in conjunction with most biological control organisms available for mite control. ACRAMITE-50WS, when used as directed, does not adversely affect populations of beneficial/predaceous mites and insects including:

Common Name	Scientific Name
Predatory mite	<i>Amblyseius fallacis</i>
Predatory mite	<i>Phytoseiulus persimilis</i>
Western predatory mite	<i>Typhlodromus occidentalis</i>
Predatory mite	<i>Typhlodromus pyri</i>
Predatory mite	<i>Zetzellia mali</i>
Sevenspotted lady beetle	<i>Coccinella septempunctata</i>
Spider mite destroyer	<i>Stethorus punctum</i>
Common lacewing	<i>Chrysopa carnea</i>
Insidious flower bug	<i>Orius insidiosus</i>
Sixspotted thrips	<i>Scolothrips sexmaculatus</i>
Western flower thrips	<i>Frankliniella occidentalis</i>

The use of these organisms in conjunction with ACRAMITE-50WS is encouraged as a means of reducing the number of chemical applications.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a dry location.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Place water soluble bag directly into application equipment (see directions). Then dispose of empty outer bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT NOTICE**—Sellers warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and to the extent consistent with applicable law, the buyer assumes the risk of any such use.

®ACRAMITE is a Registered Trademark of Chemtura Corporation  
© Copyright 2008, Chemtura Corporation