

400-514

05-26-2010

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MAY 26 2010

R. Tim Weiland, Ph.D.
Chemtura Corporation
199 Benson Rd.
Middlebury, CT 06749

Subject: Notification of Change to Packaging Size Supporting the Addition of 1 Gallon
Container per PR Notice 98-10
Acramite®-4SC
EPA Registration No. 400-514

Dear Dr. Weiland:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide
Registration Notice (PRN) 98-10 dated April 30, 2010 for the product Acramite®-4SC.

The Registration Division (RD) has conducted a review of this request for its
applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN
98-10. The Confidential Statement of Formula (CSF) submitted with the application has been
stamped "Notification" and will be placed in our records.

If you have any questions, please contact me directly at 703-305-5967 or via e-mail at
gaines.jennifer@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Jennifer Gaines". The signature is written in a cursive style with a large initial "J".

Jennifer Gaines
Wildlife Biologist
Insecticide-Rodenticide Branch
Registration Division (7505P)
Office of Pesticide Programs



Chemtura Corporation
 199 Benson Road
 Middlebury, CT 06749
 203-573-2000 tel
 203-573-2958 fax
 www.chemtura.com

3
9

April 30, 2010

Document Processing Desk (NOTIF)
 Office of Pesticide Programs (7505P)
 U.S. Environmental Protection Agency
 2777 S. Crystal Drive
 Room S-4900, One Potomac Yard 4th Floor
 Arlington, Virginia 22202-4501

Attention: Mr. Kable Davis, PM 7
 Insecticide-Rodenticide Branch (7505P)

Subject: **AMENDED LABEL via Notification - ACRAMITE[®]-4SC (EPA Reg. No. 400-514); Storage Disposal section altered to support additional package container sizes.**

Dear Mr. Davis,

Chemtura Corporation is submitting a notification label for Acramite-4SC (EPA Reg. No. 400-514). Container Disposal verbiage headers have been amended to support existing and requested container sizes captured in the attached EPA forms (8570-1). This request is in concurrence with guidance sent to Chemtura by EPA's Jennifer Gaines in an April 29, 2010 email.

Should you have any questions regarding this submission, please contact R. Tim Weiland at 203-573-2027 or email: tim.weiland@chemtura.com.

Sincerely,

Chemtura Corporation

R. Tim Weiland / for

Michael R. Dupré
 Product Registration Manager, North America

Attachments:


1. EPA Form 8570-1 for ACRAMITE-4SC (EPA Reg. No. 400-514) for 1 gallon container
2. EPA Form 8570-1 for ACRAMITE-4SC) for 15 & 30 gallon containers
3. One (1) copy, ACRAMITE-4SC current label, version 024/012610
4. Two (2) copies, ACRAMITE-4SC pending label, version 026
5. One (1) copy, ACRAMITE-4SC pending label with changes identified by shading, ver. 026

MAY 26 2010

GROUP **un** ACARICIDE

216

Acramite®-4SC



Agricultural Miticide For agricultural use only

Net
Contents:

Active Ingredient: (% by weight)

| | |
|---|--------|
| bifenazate: hydrazine carboxylic acid, 2-(4-methoxy-[1,1'-biphenyl]-3-yl) 1-methylethyl ester | 43.2% |
| Other Ingredients:..... | 56.8% |
| Total..... | 100.0% |

Contains 4 lbs. active ingredient bifenazate per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

- IF SWALLOWED:**
- Call poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.

EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

| | |
|--|---------------------|
| EMERGENCY PHONE | 800-292-5898 |
| SAFETY DATA AND INFORMATION | 866-430-2775 |
| TRANSPORTATION EMERGENCY (CHEMTREC) | 800-424-9300 |

EPA REG. NO. 400-514
EPA EST. NO.
026

Chemtura Corporation
199 Benson Road
Middlebury CT 06749



www.chemtura.com

512

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and Other Handlers Must Wear: Long-sleeved shirt & long pants; shoes plus socks. When not using a closed system, wear apron and chemical-resistant gloves made of any waterproof material for mixing/loading activities. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, estuarine/marine invertebrates, and fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 12 hours.** [Exceptions are listed under the Use Instructions for each crop.]

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- shoes plus socks
- chemical-resistant gloves made of any waterproof material

USE INFORMATION

ACRAMITE®-4SC is a suspension concentrate (flowable). It is a selective miticide for the control of a variety of mite pests on the crops listed on this label. When used as directed and applied to the foliage, it provides quick knockdown through contact activity, and long residual control. Due to its carbazate chemistry, mode of action and selective nature, ACRAMITE-4SC is relatively inactive against beneficial/predaceous mites and insects and therefore is compatible with IPM and resistance management programs.

ACRAMITE-4SC is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces is necessary for effective control.

ALWAYS SHAKE OR STIR THIS PRODUCT WELL BEFORE USE

MIXING INSTRUCTIONS

Fill spray tank with 1/2 the desired amount of water. Then add the required amount of ACRAMITE-4SC with agitation running to fully disperse the product. Then fill the tank with the remaining amount of required water.

Like many pesticides, ACRAMITE-4SC stability can be impacted by high pH and high temperature. For optimum performance, maintain spray mixtures containing ACRAMITE-4SC within a range of pH 5.5 to 6.5.

Restrictions: Tank mixtures are permitted only in those states where the tank mix portion is registered. When tank mixing, follow the label directions for most restrictive of label precautions and limitations.

Compatibility: To obtain broad spectrum insect control, ACRAMITE-4SC can be tank-mixed with other insecticide products. However, due to variations in water quality, e.g., hardness and pH, it is required that users conduct small scale trials under local conditions to ensure compatibility prior to any large scale use.

USE RATES AND DIRECTIONS

Refer to **USE INSTRUCTIONS** table for application rates, application number, and PHI for labeled crops.

For ground application, also refer to **USE INSTRUCTIONS** table for minimum gallons of spray solution per acre using equipment such as, but not limited to, compressed air, hydraulic ground boom or air-blast sprayers.

For aerial application, also refer to **USE INSTRUCTIONS** table for minimum gallons of spray solution per acre (or the minimum permitted by your state, but not less than shown) using either a fixed-wing aircraft or helicopter. Always use a spray volume adequate to assure complete coverage of the crop canopy.

For chemigation application, refer to **CHEMIGATION USE PRECAUTIONS FOR MINT AND POTATOES** section; **USE INSTRUCTIONS** table exhibits application rate range. Only one application may be made per year. Sprinkler systems must be operated at 80 to 100% during treatment application to apply the minimum amount of water possible.

To provide maximum residual control, make application as soon as mites appear. Use the lower rate where mite infestations are light. The higher rate may be required for heavy infestations or for extended residual control.

When used as directed, ACRAMITE-4SC is effective for the control of a variety of mites species, especially spider mites, red mites and grass mites. NOTE: It is not effective against rust mites, broad mites and flat mites. ACRAMITE-4SC is primarily active on the motile stage of mites, but also has ovidical activity against spider mites (*Tetranychus* species).

Restrictions:

- Rotational Crops - This product has a plantback restriction of 30 days. Do not plant another crop within 30 days after last ACRAMITE application due to chances of bifentazate residues showing up in rotational crops.
- Do not exceed the maximum amount of bifentazate allowed per crop per season, regardless of the bifentazate-containing product(s) used.
- Do not tank mix oil with ACRAMITE-4SC when applying to Golden Delicious apples.

MITES CONTROLLED

Banks grass
Brown almond
Clover

European red (use maximum rate)
McDaniel
Pacific spider

Pecan leaf scorch
Strawberry spider
Southern red mite

Spruce spider mite
Two-spotted spider
Willamette

**BEARING CROPS
USE INSTRUCTIONS**

| CROP | AMOUNT ACRAMITE-45C PER ACRE (A) FL. OZ. | MINIMUM GALLONS PER ACRE | | CHEMIGATION ACRE INCHES OF WATER | TOTAL NUMBER OF SPRAYS PER SEASON | MINIMUM DAYS BETWEEN APPLICATIONS | HARVEST DAYS AFTER APPLICATION (PHI DAYS) |
|---|---|--------------------------------|-------------------|--|--|--|--|
| | | GROUND | AIR | | | | |
| CANEBERRY SUBGROUP 13-07A, (B) WILD RASPBERRY | 12 - 16 | 50 | — | — | 2 | 30 (N) | 1 |
| CHRISTMAS TREES/CONIFER PLAN- TATIONS AND NURSERIES | 12 - 16 | 100 | 10 (C) | — | 1 | — | — |
| COTTON | 16 - 24 | 20 | 5 (C) | — | 1 | — | 60 |
| CUCURBIT VEGETABLES (See footnote D for crops in addition to those listed below) CUCUMBERS; EDIBLE GOURDS (E); MUSKMELON (F); PUMPKIN; SQUASH (summer (G) & winter (H)); WATERMELON | 12 - 16 | 50 | 10 (C) | — | 1 | — | 3 |
| DRIED SHELLED BEAN (EXCEPT SOYBEAN) SUBGROUP (I) | 16 - 24 | 20 | 10 ^(D) | — | 2 | 14 (N) | 7 |
| FRUITING VEGETABLES (See footnote J for crops in addition to those listed below) EGGPLANT PEPPERS (all varieties of <i>Capsicum</i> spp.) (K) TOMATOES (field grown) | 12 - 16 | 50 | 10 (C) | — | 1 | — | 3 |
| GRAPES (L) | 12 - 16 | 50 | — | — | 1 | — | 14 |
| HOPS | 12 - 24 | 50 | — | — | 1 | — | 14 |
| LEGUME VEGETABLES SUBGROUP 6A (succulent); BEANS SUBGROUP 6B; SUCCULENT SHELLED SOYBEAN (M) | 16 - 24 | 20 | 10 (C) | — | 2 | 14 (N) | 3 |
| MINT | 12 - 24 | 50 | 10 (C) | 0.1 - 0.2 (O) | 1 | — | 7 |
| OKRA | 12 - 16 | 50 | — | — | 1 | — | 3 |
| POME FRUITS: (See footnote P for crops in addition to those listed below) APPLES; CRABAPPLES; PEARS; QUINCE | 12 - 16 | 50 | — | — | 1 | — | 7 |
| POTATOES (Q) TUBEROUS AND CORM VEGETABLES | 16 - 24 | 20 | 10 (C) | 0.1 - 0.2 (O) | 2 | 14 (N) | 14 |
| STRAWBERRIES | 12 - 16 | 100 | — | — | 2 | 21 (N) | 1 |
| STONE FRUIT: (See footnote R for crops in addition to those listed below) APRICOTS; CHERRIES (Sweet & Tart); NECTARINES; PEACHES; PLUMS/ PRUNES | 12 - 16 | 50 | — | — | 1 | — | 3 |
| SUCCULENT PEA (S) PEA, SUCCULENT SHELLED; PEA, EDD- IBLE PODDED; PEA, PIGEON | 16 - 24 | 20 | 10 (C) | — | 2 | 14 (N) | 3 |
| TREE NUTS: (See footnote T for crops in addition to those listed below) ALMOND | 12 - 24 | 50 | — | — | 1 | — | 7 |
| FILBERT (Hazelnut), PECANS, PISTA- CHIOS, WALNUTS (Black & English) | 12 - 24 | 50 | — | — | 1 | — | 14 |
| TROPICAL FRUIT: PAPAYA; STAR APPLE; BLACK SAPOTE, MANGO; SAPODILLA; CANISTEL; MAMEY SAPOTE; LYCHEE; LONGAN; SPANISH LIME; RAMBUTAN; PULASAN; GUAVA; FEIJOA; JABOTICABA; WAX JAMBU; STARFRUIT (CARAMBOLA); PASSIONFRUIT; ACEROLA | 12 - 16 | 50 | — | — | 2 | 21 (N) | 1 |

- (A) Use the higher rate under heavy mite pressure.
- (B) CANEBERRY: Blackberry; loganberry; red and black raspberry; cultivars and/or hybrids of these.
- (C) Minimum gallonage per acre or the minimum permitted by your state, but not less than shown.
- (D) CUCURBIT VEGETABLES: Other crops which may be treated with ACRAMITE-4SC with the same use instructions are: Chayote (fruit) (*Sechium edule*), Chinese waxgourd (Chinese preserving melon) (*Benincasa hispida*), Citrus melon (*Citrullus lanatus var. citroides*), Gherkin (*Cucumis anguria*), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber).
- (E) EDIBLE GOURD (*Lagenaria* spp.) (includes *hyotan* and *cucuzza*). Also (*Luffa acutangula* and *L. cylindrical*) which includes hechima and Chinese okra.
- (F) MUSKMELON: (hybrids and/or cultivars of *Cucumis melo*) includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon.
- (G) SQUASH, SUMMER: (*Cucurbita pepo var. melopepo*) includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini.
- (H) SQUASH, WINTER: (*Cucurbita maxima: C. moschata*) includes butternut squash, calabaza, hubbard squash; (*C. mixta; C. pepo*) includes acorn squash, spaghetti squash.
- (I) Cultivars of *Lupinus* spp. include grain lupin, sweet lupin, white lupin, and white sweet lupin; *Phaseolus* spp. include field bean, kidney bean, lima beans (dry), navy bean, and pinto bean; tepary bean; *Vigna* spp. include adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, and urd bean; broad bean (dry); chickpea; guar; lablab bean; and lentil.
- (J) FRUITING VEGETABLES: Other crops which may be treated with ACRAMITE-4SC with the same instructions are: groundcherry (*Physalis* spp.), pepino (*Solarium muricatum*), tomatillo (*Physalis ixocarpa*)
- (K) PEPPER (*Capacium* spp.) includes bell peppers, chili peppers, cooking peppers, pimento, and sweet peppers.
- (L) GRAPES: The REI is 5 days for cane turning, tying and girdling of table grapes. Exception: In addition to the early entry exceptions allowed by WPS, you may enter or allow workers to enter treated areas to perform all tasks other than tying, turning and girdling after 12 hours following applications, as long as at least long pants, long-sleeved shirts, shoes and socks are worn. Notify workers of the exception (including when entry is permitted for each of the tasks named in the exception).
- (M) SUCCULENTS: Bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broadbean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; soybean (immature seed); sword bean
- (N) A miticide with a different mode of action should be used between any 2 applications of ACRAMITE.
- (O) Refer to USE RATES AND DIRECTIONS and CHEMIGATION USE PRECAUTIONS sections.
- (P) POME FRUIT: Other crops which may be treated with ACRAMITE-4SC with the same instructions are: Loquat (*Eriobotrya japonica*), Mayhaw (*Crataegus aestivalis, C. opaca, and C. rufula*), Pear, oriental (*Pyrus pyrifolia*)
- (Q) POTATO: Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; tanier; turmeric; yam bean; yam, true
- (R) STONE FRUIT: Other crops which may be treated with ACRAMITE-4SC with the same instructions are: Plum (*Prunus domestica, Prunus* spp.), Chickasaw Plum (*Prunus angustifolia*), Damson Plum (*Prunus domestica* spp. *Instititia*), Japanese Plum (*Prunus salicina*) and Plumcot (*Prunus armeniaca X P. domestica*).
- (S) SUCCULENT PEA: Sugar pea, Chinese pea, Pois mange tout, Snap pea, Ming pea, Podded pea, Snow pea, China pea, Chicharo, Shi Hia wandou, Saya-endo, Sugar snap pea, Congo pea, No-eye pea, Red gram, Arher, Grandul, Gandules, Dhal, Toor, Gunds pea, Porto Rico pea, Urher Gandul, Guandu, Pois-d'angole, Gungo pea
- (T) TREE NUTS: Other crops which may be treated with ACRAMITE-4SC with the same use instructions (14 day PHI) are: Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Hickory nut, Macadamia nut (bush nut).

**NON-BEARING CROPS
USE INSTRUCTION**

ACRAMITE-4SC use on non-bearing crops, i.e. those crops which will not bear fruit within one year of application, **includes all the crops listed in the bearing crops use instructions plus crops listed below.** These include fruit trees and berries in both commercial plantings and nurseries, including transplant uses but excluding residential areas. **The listed non-bearing crops all have a 12 hour restricted entry interval (REI).** Application is by ground equipment only. Additional non-bearing crops that may be so treated are:

| CROP | MITES CONTROLLED | AMOUNT ACRAMITE-4SC PER ACRE | MINIMUM GALLONS SPRAY SOLUTION PER ACRE | TOTAL NUMBER OF SPRAYS PER YEAR |
|---|---|------------------------------|---|---------------------------------|
| Berries (blueberry, highbush and lowbush; currant; elderberry; gooseberry; huckleberry) Citrus (grapefruit, lemons, limes, oranges, tangerines, etc.) Dates, Figs Persimmons | European red Two-spotted spider Pacific spider McDaniel Strawberry spider Citrus red Southern red | 12 - 16 fl.oz | 50 | 1 |

CHEMIGATION USE PRECAUTIONS FOR MINT AND POTATOES

- A. Apply this product only through sprinkler systems, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make the necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Constant agitation must be maintained in the chemical supply tank during the entire period of miticide application.
- N. Inject the product with a positive replacement pump into the main line ahead of a right angle turn, to insure adequate mixing.
- O. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.
- P. Do not apply when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.
- Q. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute mixture per hour. Pesticide should be applied continuously for the duration of the water addition.
- R. Where sprinkler irrigation patterns do not overlap sufficiently unacceptable mite control may result. Where sprinkler distribution patterns overlap excessively crop injury may result.
- S. Check with state lead agencies for state specific chemigation requirements.

STRATEGIES FOR RESISTANCE PREVENTION

When used as directed, ACRAMITE-4SC combines high activity on mites with safety to beneficial/predaceous mites and insects. In addition, the carbazate chemistry of ACRAMITE-4SC provides a means of controlling mites which have developed resistance to commonly used products. These properties can result in fewer miticide/insecticide applications as well as general reduction in the problems caused by resistance.

ACRAMITE-4SC has demonstrated no cross resistance with other commercial miticides. ACRAMITE-4SC contains an active ingredient classified as a Group un acaricide. ACRAMITE-4SC is suitable to be used as a rotational partner with other miticides.

Follow the mite control strategies below:

- Incorporate IPM techniques into your insect control program.
- Ensure thorough spray coverage to all foliage.
- Scout regularly and apply ACRAMITE-4SC as soon as infestations are observed. Do not wait until large populations have established.
- Always apply ACRAMITE-4SC at the required rates and according to label information.
- Unless labeled otherwise, use only one application of ACRAMITE-4SC per year, and rotate to a product with a different mode of action grouping.
- Because of its selectivity, ACRAMITE-4SC can be used in conjunction with most biological control organisms available for mite control. ACRAMITE-4SC, when used as directed, does not adversely affect populations of beneficial/predaceous mites and insects including:

| Common Name | Scientific Name |
|---------------------------|-----------------------------------|
| Predatory mite | <i>Amblyseius fallacis</i> |
| Predatory mite | <i>Phytoseiulus persimilis</i> |
| Western predatory mite | <i>Typhlodromus occidentalis</i> |
| Predatory mite | <i>Typhlodromus pyri</i> |
| Predatory mite | <i>Zetzellia mali</i> |
| Seven-spotted lady beetle | <i>Coccinella septempunctata</i> |
| Spider mite destroyer | <i>Stethorus punctum</i> |
| Common lacewing | <i>Chrysopa carnea</i> |
| Insidious flower bug | <i>Orius insidiosus</i> |
| Six-spotted thrip | <i>Scolothrips sexmaculatus</i> |
| Western flower thrip | <i>Frankliniella occidentalis</i> |

The use of these organisms in conjunction with ACRAMITE-4SC is encouraged as a means of reducing the number of chemical applications.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable containers equal to or less than 5 gallons.

Do not reuse or refill this container. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If pressure rinsing, empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration. Do not burn, unless allowed by state and local ordinances.

Nonrefillable containers greater than 5 gallons. Do not reuse or refill this container. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration. Do not burn, unless allowed by state and local ordinances.

IMPORTANT NOTICE—Sellers warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and to the extent consistent with applicable law, the buyer assumes the risk of any such use.

®ACRAMITE is a Registered Trademark of Chemtura Corporation
© Copyright 2010, Chemtura Corporation

2/8