

400-518

09/26/2006

Jacket  
1/7



**UNITED STATES ENVIRONMENTAL PROTECTION  
AGENCY**  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Judith O. Ball  
Product Registration Manager  
North America  
Chemtura USA Corporation  
199 Benson Road  
Middlebury, CT 06749

SEP 26 2006

Subject: Procure 480SC  
EPA Registration No. 400-518  
Your Amendment dated May 8, 2006  
Decision No. 367641

Dear Ms. Ball:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following change is made:

1. Clarify the "Emergency Assistance" heading by adding the phrase "(Human health)" immediately after the heading.

One copy of the label stamped "Accepted with comments" is enclosed for your records. This label supercedes all labels previously accepted for this product. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

Sincerely,


A handwritten signature in black ink that reads "Mary L. Waller".

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505C)

Enclosure

HIGHLIGHTED COPY

# Procure® 480SC



## Agricultural Fungicide

Net Contents:  
**2.5**  
gallons

Active Ingredient: (% by weight)  
 Triflumizole [1-[1-(4-chloro-2-(trifluoromethyl) phenyl) imino)-2-propoxyethyl]-1H-imidazole] ..... 42.14%  
 Other Ingredients: ..... 57.86%  
 Total: ..... 100.0%

Contains 4 lbs. triflumizole per gallon.

### KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>EMERGENCY PHONE</b>	800-292-5898
<b>SAFETY DATA AND INFORMATION</b>	203-573-3303
<b>TRANSPORTATION EMERGENCY (CHEMTREC)</b>	800-424-9300

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

SEP 26 2006

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

400-518

EPA REG. NO. 400-518  
EPA EST. NO.  
006

Chemtura USA Corporation  
Middlebury CT 06749



www.chemtura.com

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and Other Handlers Must Wear:** A long-sleeved shirt & long pants; chemical-resistant gloves made of any waterproof material; shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

**PHYSICAL or CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label directions before using.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

coveralls; chemical-resistant gloves made of any waterproof material; and shoes plus socks.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry location.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:**

**Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**

PROCURE 480SC is a broad-spectrum foliar fungicide for control of certain diseases of apples, cherries, cucurbits, grapes, pears, pineapples and strawberries. PROCURE 480SC is active as a protectant and as an eradicant by preventing disease symptoms after infection has occurred. It's anti-sporulant activity reduces spores after lesions become visible.

Read the entire label and directions before using PROCURE 480SC.

**APPLICATION INSTRUCTIONS**

PROCURE 480SC can be applied with dilute or concentrate ground equipment. Aerial applications for tree fruits can be made in accordance with use directions noted for each crop. Equipment should be frequently checked for calibration. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, bloom and fruit.

Use higher recommended rates on more susceptible varieties and/or under higher disease pressure.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

**GENERAL PRECAUTIONS AND RESTRICTIONS**

Do not allow livestock to graze on treated orchard ground cover.

Aerial application of PROCURE 480SC is allowed unless prohibited under a specific crop use direction. Do not apply this product through any type of irrigation system.

**RESISTANCE MANAGEMENT:** Populations of fungal pathogens are known to develop resistance to fungicides with the same mode of action when used repeatedly. Thus, Chemtura encourages responsible resistance management to ensure an effective long-term control of diseases. Resistance management strategies include alternating and/or tank mixing with other labeled fungicides having different modes of action or limiting the total number of applications per season.

**RESTRICTIONS ON ROTATIONAL CROPS:** Do not plant leafy vegetables within 30 days after application. Do not plant root vegetables within 60 days after application. Rotation to all other crops within one year after application, unless PROCURE 480SC is registered for use on those crops, is prohibited.

**APPLES & PEARS**

On apples and pears, PROCURE 480SC should be tank mixed with standard protectant fungicides labeled for control of other diseases during the season for maximum fruit disease control. Make applications at 7 to 10 day intervals through the first cover and at approximately 10 to 14 day intervals thereafter as needed. PROCURE 480SC may be applied up to 14 days prior to harvest. PROCURE 480SC rates should be adjusted in response to disease pressure, weather conditions, and varietal susceptibility. If disease pressure is above normal, rates per acre must be increased as indicated in the "Comments" section for each disease. Do not use less than the recommended rates.

AERIAL APPLICATIONS: When making aerial applications use 12 to 16 fl. oz. PROCURE 480SC per acre\*

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Apples	<b>Powdery Mildew</b> ( <i>Podosphaera leucotricha</i> )	8 to 16	Initiate spraying at first sign of mildew, usually at the 1/2" green tip stage. Continue applications at 7 to 14 day intervals as needed through the terminal growth period.
	<b>Scab</b> ( <i>Venturia inaequalis</i> ) <b>Cedar Apple Rust</b> ( <i>Gymnosporangium juniperi-virginianae</i> )	8 to 16	<b>Standard Spray Schedule</b> Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre at 7 to 10 day intervals from 1/2" green tip stage through second cover or until primary scab sporulation is complete. Higher rates should be used for more susceptible varieties and heavy disease pressure. For improved scab control apply PROCURE 480SC in combination with a protectant fungicide. If changing to a protectant fungicide only after the end of the primary infection period, use 2 PROCURE 480SC applications, 7 days apart. <b>Eradicant Spray Schedule (Post Infection Schedule)</b> Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre within 96 hours of the start of an infection period. Under low to moderate pressure conditions, 8 fl. oz per acre can be applied within 48 hours after the start of an infection period. Higher rates should be used if disease pressure is high or the time interval after the start of infection is greater than 48 hours. If a second infection period occurs within 3 days of the first application, a second application should be made within 7 days of the first treatment. For improved scab control apply PROCURE 480SC in combination with a protectant fungicide. After first cover, PROCURE 480SC can be continued in a standard cover spray program in combination with protectant fungicides which control other diseases. Do not apply more than 64 fl. oz of PROCURE 480SC per acre of apples per season. Application of PROCURE 480SC for control of scab will also control cedar apple rust.
Pears	<b>Powdery Mildew</b> ( <i>Podosphaera leucotricha</i> )	8 to 16	Initiate spray at pink or white stage. Continue applications at 7 to 14 day intervals or as required. PROCURE 480SC may applied up to 14 days prior to harvest.
	<b>Scab</b> ( <i>Venturia pirina</i> )	8 to 16	<b>Standard Spray Schedule</b> Apply PROCURE 480SC at the rate of 8 to 16 fl. oz/A, at 7 to 10 day intervals from green tip stage through second cover. Higher rates should be used for more susceptible varieties and heavy disease pressure. Continue applications of PROCURE 480SC in a standard cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the primary infection period, use 2 PROCURE 480SC applications, 7 days apart. <b>Eradicant Spray Schedule (Post Infection Schedule)</b> Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre within 96 hours of the start of an infection period. Under low to moderate pressure conditions, 8 fl. oz per acre can be applied within 48 hours after the start of an infection period. Higher rates should be used if disease pressure is high or the time interval after the start of infection an infection period is greater than 48 hours. If a second infection period occurs within 3 days of the first application, a second application should be made within 7 days of the first treatment. For improved scab control, apply PROCURE 480SC in combination with a protectant fungicide. After first cover, PROCURE 480SC can be continued in a standard cover spray program of 8 to 12 fl. oz/A, in combination with protectant fungicides which control other diseases. Do not apply more than 64 fl. oz. of PROCURE 480SC per acre of pears per season.

\*Not registered for use by air in California.

**CHERRIES (SWEET & TART)**

PROCURE 480SC can be applied throughout the season from delayed dormant to within 1 day of harvest. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, flowers and fruit. Applications should be timed to maintain coverage on susceptible plant tissues throughout possible infection periods. Do not use less than the recommended rates.

CROP	DISEASE	Rate of Procure 480SC <sup>(1)</sup> fl. oz/acre	COMMENTS
Cherries (Sweet & Tart)	<b>Powdery Mildew</b> ( <i>Podosphaera clandestina</i> )	8 to 16	Initiate treatments at early popcorn and continue at 7 to 14 day intervals until terminal growth ceases.
	<b>Blossom Blight &amp; Fruit Rot</b> ( <i>Monilinia spp.</i> )	10 to 16	<b>Blossom Blight Control:</b> Treatments should be applied at early popcorn, full bloom and petal fall. Complete coverage must be maintained throughout the bloom period. <b>Fruit Rot Control:</b> Depending on disease pressure, continue the blossom blight schedule with applications at 7 to 10 day intervals throughout the cover spray period.
	<b>Leaf Spot</b> ( <i>Blumeriella jaapii</i> )	10 to 16	PROCURE 480SC applications made to control <i>Podosphaera</i> or <i>Monilinia</i> will suppress the development of leaf spot. To control leaf spot, PROCURE 480SC should be tank mixed with other products registered for leaf spot control.

(1) Application Rates:

When making aerial applications, use at least 12 fl. oz PROCURE 480SC per acre. Where feasible, ground application should be used because it provides better canopy penetration and coverage.

Always use higher recommended rates on more susceptible varieties or when higher disease pressure conditions exist.

Do not apply more than 96 fl. oz. of PROCURE 480SC per acre per season. Do not apply closer than 1 day before harvest.

**CUCURBITS**

PROCURE 480SC can be used to control powdery mildews caused by *Erysiphe cichoracearum* or *Podosphaera xanthii*. If other leaf diseases in addition to powdery mildew must be controlled, PROCURE 480SC can be tank mixed with other fungicides registered for those diseases. Thorough coverage of the foliage is necessary for maximum disease control. Repeated applications should be made if conditions of powdery mildew development persist.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Cucumbers Gherkins Gourds Muskmelons (includes cantaloupes, honeydews, Persian melons and other melons) Pumpkins Squash (summer and winter) Watermelons	<b>Powdery Mildew</b> ( <i>Erysiphe cichoracearum</i> , <i>Podosphaera xanthii</i> )	4 to 8	Apply PROCURE 480SC as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin at the start of vining or the first indication of disease. Repeated applications can be made at 7 to 14 day intervals. When conditions are favorable for severe disease pressure, use higher rates shorter intervals. Do not apply more than 40 fl. oz. of PROCURE 480SC per acre per season. Applications can be made up to the day of harvest.

6/7

**GRAPES**

PROCURE 480SC should be used as a protectant for control of powdery mildew. Start applications before bloom and continue on a 14 to 21 day interval at 4 to 8 fl. oz/acre. For the most susceptible varieties, use 6 to 8 fl. oz/acre and/or shorter spray intervals. When disease pressure is low, use 4 to 6 fl. oz/acre and/or the spray interval can be extended up to 21 days. Select spray volume, nozzle adjustments and ground speed to ensure complete coverage of fruit clusters and foliage. Use a minimum of 50 gallons of water per acre for early season applications. Use enough volume to ensure thorough coverage as the amount of foliage increases. Do not apply within 7 days of harvest. Do not exceed 32 fl. oz. PROCURE 480SC per acre per season. Do not apply by air. Sulfur at recommended use rates can be applied before, with, and/or between PROCURE 480SC applications.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Grapes	<b>Powdery Mildew</b> ( <i>Uncinula necator</i> )	4 to 8	Start applications before bloom and continue on a 14 to 21 day interval. Use higher rates and/or shorter intervals on susceptible varieties or under heavy disease pressure. PROCURE 480SC applied in a powdery mildew spray schedule, will enhance the activity of registered fungicides used for the control of Botrytis bunch rot. Applications must be made on a 14-day spray schedule for suppression of Botrytis bunch rot.

**HAZELNUTS (FILBERTS)**

Application may be made by ground or air. Minimum recommended spray volumes are 50 gallons per acre for aerial applications and 100 gallons per acre for ground applications. Use the highest labeled rate by air. Apply PROCURE 480SC in sufficient spray volume to ensure thorough coverage and mixing the lowest labeled rate of a spray schedule with PROCURE 480SC may improve spray coverage and penetration. Ground application is preferred.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Hazelnuts (Filberts)	<b>Eastern Filbert Blight</b> ( <i>Ascochyta arbuti</i> )	4 to 8	PROCURE 480SC should be applied at budbreak when the first green tissue is visible as a preventative spray. Continue applications of PROCURE 480SC at 7 to 14 day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during wet and cool spring weather. The spraying schedule may be adjusted, apply the highest rate to varieties that are highly susceptible to the fungal disease or when severe disease conditions exist. Do not apply more than 32 fl. oz. of PROCURE 480SC per acre per season. Do not apply within 7 days of harvest. PROCURE 480SC may be applied as a tank mix application with other fungicides as a pest management strategy.

**PINEAPPLES**

**Pre-Plant Seed Piece Material Treatment**

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	DISEASE PRESSURE	COMMENTS
Pineapples	<b>Pineapple Butt Rot</b> ( <i>Chalara paradoxa</i> )	4 - 8 8-12 12-16	Light Moderate Severe	Apply PROCURE 480SC at 4 - 16 fl. oz/100 gallons as a planting material dip treatment. Freshly removed seed material used for immediate planting should be immersed within 12 hours of removal for thorough wetting and coverage. Use a higher label rate under heavy disease pressure. Do not use treated green forage or green fodder for animal feed.

**STRAWBERRIES**

PROCURE 480SC can be applied for control of strawberry powdery mildew with dilute or concentrate ground equipment. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, flowers and fruit. Use higher recommended application rates and shorter spray intervals on more susceptible varieties and/or under higher disease pressure. PROCURE 480SC may be tank mixed with other materials registered for use on strawberries. Apply tank mixtures in accordance with all applicable label directions. Do not apply within 1 day of harvest. Do not exceed 32 fl. oz. PROCURE 480SC per acre per season.

CROP	DISEASE	Rate of Procure 480SC fl. oz./acre	COMMENTS
Strawberries	<b>Powdery Mildew</b> ( <i>Sphaerotheca macularis</i> <i>f. sp. fragariae</i> )	4 to 8	Protection of leaves before bloom will reduce in-season disease pressure and result in greater control of flower and fruit infections. Initiate applications at the first sign of powdery mildew. Continue applications at approximately 14 day intervals while conditions favor disease development.

**IMPORTANT NOTICE**—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

• PROCURE is a Registered Trademark of Chemtura Corporation

© Copyright 2006, Chemtura Corporation