

400-518

6/12/2009

Jacket
10/12

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Judith O. Ball
Product Registration Manager
North America
Chemtura USA Corporation
199 Benson Road
Middlebury, CT 06749

JUN 12 2009

Subject: Procure 480 SC
EPA Registration No. 400-518, PP#: 7E&183
Your Amendment dated January 25, 2007, Adding the new use- Brassica, leafy greens, subgroup 5B and turnip greens
PRIA Decision No. 374868

Procure 480 SC
EPA Registration No. 400-518, PP#: 7E7258
Your Amendment dated August 15, 2007, Adding the new use- Leafy greens, subgroup 4A, except spinach, cilantro, Swiss chard, pineapple, Papaya and related tropical fruits, and hops
PRIA Decision No. 383406

Procure 480 SC
EPA Registration No. 400-518, PP#: 7E7286
Your Amendment dated October 18, 2007, Adding the new use- Head and Stem Brassica, subgroup 5A
PRIA Decision No. 385791

Dear Ms. Ball:

The labeling referred to above, submitted in connection with registration under section under 3(c)(7)(b) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided you make the following changes:

1. Please remove the company website from the label. Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading.

In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency finds or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance. As an alternative, you may refer consumers to the company's phone number or email address.

2. Page 2, after the "Personal Protective Equipment" heading, add the acronym "(PPE)."
3. Page 1, add "User Safety Requirements" heading. Add a third bullet which states, "Remove PPE immediately after handling the product. Wash...into clean clothing."
4. Page 1, add the word, "exist" after the word "washables."
5. Pages 5, 6, and 9, delete the references to the phrase, "recommended rates" or "recommended spray volumes" and replace the phrase with "specified application" or "specified spray volumes."
6. Page 8, delete the phrase "(including field and greenhouse grown)" in the "Leafy Greens" heading. Also, delete the sentence "For greenhouse grown leafy greens apply 6 to 8 fl. oz. in 100 gallons of water." within the comments section of the leafy greens table.
7. Page 8, delete the crop listing for "Swiss chard" under the "Leafy Greens (except spinach)" crop group heading.
8. In the "Environmental Hazards" section, delete the "period" after the word "washwaters" and add the phrase, "or rinsate." The sentence shall read, "DO NOT contaminate water when cleaning equipment or disposing of equipment washwater or rinsate."
9. In the "User Safety Recommendations" section revise the second bullet to read, "Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing."
10. Please update the Warranty Statement in accordance with the label review manual <http://www.epa.gov/oppfead1/labeling/lrm/chap-12.htm#IX>.

Data Requirements:


11. Submit to the Agency an immunotoxicity study (OPPTS Guideline No. 870.7800) by April 2010.

30f12

Page 3
Procure 480SC
Decision Numbers: 374868, 383406, 385791

One copy of the label stamped "Accepted with comments" is enclosed for your records. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment. If you have any questions, please contact Tamue L. Gibson by phone at (703) 305-9096 or via email at gibson.tamue@epa.gov.

Sincerely,




Shaja Brothers Joyner
Product Manager (20)
Fungicide Branch
Registration Division (7505P)

Enclosure

40f12

Procure[®] 480SC



Agricultural Fungicide

Net Contents:

Active Ingredient: (% by weight)

Triflumizole [1-[1-(4-chloro-2-(trifluoromethyl) phenyl) imino]-2-propoxyethyl]-1H-imidazole.....	42.14%
Other Ingredients:	57.86%
Total:.....	100.00%

Contains 4 lbs. triflumizole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything to an unconscious person.
<p>EMERGENCY ASSISTANCE (Human health): Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
EMERGENCY PHONE	800-292-5898
SAFETY DATA AND INFORMATION	866-430-2775
TRANSPORTATION EMERGENCY (CHEMTREC)	800-424-9300

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JUN 12 2009
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-518

EPA REG. NO. 400-518
EPA EST. NO. 013

Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury CT 06749



www.chemtura.com

50412

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; chemical-resistant gloves made of any waterproof material; shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label directions before using.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

coveralls; chemical-resistant gloves made of any waterproof material; and shoes plus socks.

60912

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry, secure location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill this container.

For containers equal to or less than 5 gallons in size: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

If pressure rinsing: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

GENERAL INFORMATION

PROCURE 480SC is a broad-spectrum foliar fungicide for control of certain diseases of apples, cherries, cucurbits, grapes, pears, pineapples and strawberries. PROCURE 480SC is active as a protectant and as an eradicant by preventing disease symptoms after infection has occurred. It's anti-sporulant activity reduces spores after lesions become visible.

Read the entire label and directions before using PROCURE 480SC.

APPLICATION INSTRUCTIONS

PROCURE 480SC can be applied with dilute or concentrate ground equipment. Aerial applications for tree fruits can be made in accordance with use directions noted for each crop. Equipment should be frequently checked for calibration. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, bloom and fruit.

Use higher recommended rates on more susceptible varieties and/or under higher disease pressure.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not allow livestock to graze on treated orchard ground cover. Aerial application of PROCURE 480SC is allowed unless prohibited under a specific crop use direction. Do not apply this product through any type of irrigation system.

RESISTANCE MANAGEMENT: Populations of fungal pathogens are known to develop resistance to fungicides with the same mode of action when used repeatedly. Thus, Chemtura encourages responsible resistance management to ensure an effective long-term control of diseases. Resistance management strategies include alternating and/or tank mixing with other labeled fungicides having different modes of action or limiting the total number of applications per season.

RESTRICTIONS ON ROTATIONAL CROPS: Do not plant spinach or fruiting vegetables within 30 days after application. Do not plant bulb vegetables or root vegetables within 60 days after application. Do not plant cotton, small cereal grains and all other crops not registered on the label for PROCURE 480SC, within one year after application.

APPLES & PEARS

On apples and pears, PROCURE 480SC should be tank mixed with standard protectant fungicides labeled for control of other diseases during the season for maximum fruit disease control. Make applications at 7 to 10 day intervals through the first cover and at approximately 10 to 14 day intervals thereafter as needed. PROCURE 480SC may be applied up to 14 days prior to harvest. Do not apply more than 64 fl. oz. of PROCURE 480SC per acre of apples or pears per season.

PROCURE 480SC rates should be adjusted in response to disease pressure, weather conditions, and varietal susceptibility. If disease pressure is above normal, rates per acre must be increased as indicated in the "Comments" section for each disease. Do not use less than the recommended rates.

AERIAL APPLICATIONS: When making aerial applications use 12 to 16 fl. oz. PROCURE 480SC in a minimum of 10 gallons of water per acre*

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Apples	Powdery Mildew (<i>Podosphaera leucotricha</i>)	8 to 16	Initiate spraying at first sign of mildew, usually at the 1/2" green tip stage. Continue applications at 7 to 14 day intervals as needed through the terminal growth period.
	Scab (<i>Venturia inaequalis</i>) Cedar Apple Rust (<i>Gymnosporangium juniperi-virginianae</i>)	8 to 16	Standard Spray Schedule Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre at 7 to 10 day intervals from 1/2" green tip stage through second cover or until primary scab sporulation is complete. Higher rates should be used for more susceptible varieties and heavy disease pressure. For improved scab control apply PROCURE 480SC in combination with a protectant fungicide. If changing to a protectant fungicide only after the end of the primary infection period, use 2 PROCURE 480SC applications, 7 days apart. Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre within 96 hours of the start of an infection period. Under low to moderate pressure conditions, 8 fl. oz per acre can be applied within 48 hours after the start of an infection period. Higher rates should be used if disease pressure is high or the time interval after the start of infection is greater than 48 hours. If a second infection period occurs within 3 days of the first application, a second application should be made within 7 days of the first treatment. For improved scab control apply PROCURE 480SC in combination with a protectant fungicide. After first cover, PROCURE 480SC can be continued in a standard cover spray program in combination with protectant fungicides which control other diseases. Application of PROCURE 480SC for control of scab will also control cedar apple rust.
Pears	Powdery Mildew (<i>Podosphaera leucotricha</i>)	8 to 16	Initiate spray at pink or white stage. Continue applications at 7 to 14 day intervals or as required. PROCURE 480SC may applied up to 14 days prior to harvest.
	Scab (<i>Venturia pirina</i>)	8 to 16	Standard Spray Schedule Apply PROCURE 480SC at the rate of 8 to 16 fl. oz/A, at 7 to 10 day intervals from green tip stage through second cover. Higher rates should be used for more susceptible varieties and heavy disease pressure. Continue applications of PROCURE 480SC in a standard cover spray program with a protectant fungicide. If changing to a protectant fungicide only after the end of the primary infection period, use 2 PROCURE 480SC applications, 7 days apart. Eradicant Spray Schedule (Post Infection Schedule) Apply PROCURE 480SC at the rate of 8 to 16 fl. oz per acre within 96 hours of the start of an infection period. Under low to moderate pressure conditions, 8 fl. oz per acre can be applied within 48 hours after the start of an infection period. Higher rates should be used if disease pressure is high or the time interval after the start of infection an infection period is greater than 48 hours. If a second infection period occurs within 3 days of the first application, a second application should be made within 7 days of the first treatment. For improved scab control, apply PROCURE 480SC in combination with a protectant fungicide. After first cover, PROCURE 480SC can be continued in a standard cover spray program of 8 to 12 fl. oz/A, in combination with protectant fungicides which control other diseases.

*Not registered for use by air in California.

BRASSICA LEAFY GREENS

PROCURE 480SC can be used to control powdery mildew and black spot on the following crops. If other leaf diseases in addition to powdery mildew or black spot must be controlled, PROCURE 480SC can be tank mixed with other fungicides registered for those diseases. Repeated applications should be made if conditions of powdery mildew development persist.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens (Including all cultivars and/or hybrids of these)	Powdery Mildew <i>(Erysiphe polygoni)</i> Black Spot <i>(Alternaria spp.)</i>	6 to 8	Apply PROCURE 480SC as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin at the first indication of disease. Use higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Repeated applications should be made on a 14 day schedule. Do not apply more than 18 fl. oz. of Procure 480SC per acre per season. Use a minimum of 10 gallons per acre spray volume for ground applications. Use a minimum of 2 gallons per acre spray volume for aerial applications. Applications can be made one day before harvest.

CHERRIES (SWEET & TART)

PROCURE 480SC can be applied throughout the season from delayed dormant to within 1 day of harvest. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, flowers and fruit. Applications should be timed to maintain coverage on susceptible plant tissues throughout possible infection periods. Do not use less than the recommended rates.

CROP	DISEASE	Rate of Procure 480SC ⁽¹⁾ fl. oz/acre	COMMENTS
Cherries (Sweet & Tart)	Powdery Mildew <i>(Podosphaera clandestina)</i>	8 to 16	Initiate treatments at early popcorn and continue at 7 to 14 day intervals until terminal growth ceases.
	Blossom Blight & Fruit Rot <i>(Monilinia spp.)</i>	10 to 16	Blossom Blight Control: Treatments should be applied at early popcorn, full bloom and petal fall. Complete coverage must be maintained throughout the bloom period. Fruit Rot Control: Depending on disease pressure, continue the blossom blight schedule with applications at 7 to 10 day intervals throughout the cover spray period.
	Leaf Spot <i>(Blumeriella jaapii)</i>	10 to 16	PROCURE 480SC applications made to control <i>Podosphaera</i> or <i>Monilinia</i> will suppress the development of leaf spot. To control leaf spot, PROCURE 480SC should be tank mixed with other products registered for leaf spot control.

(1) Application Rates:
 When making aerial applications, use at least 12 fl. oz PROCURE 480SC in a minimum of 10 gallons of water per acre. Where feasible, ground application should be used because it provides better canopy penetration and coverage.
 Always use higher recommended rates on more susceptible varieties or when higher disease pressure conditions exist.
 Do not apply more than 96 fl. oz. of PROCURE 480SC per acre per season. Do not apply closer than 1 day before harvest.

90912

CUCURBITS (Including greenhouse transplants)

PROCURE 480SC can be used to control powdery mildews caused by *Erysiphe cichoracearum* or *Podosphaera xanthii*. If other leaf diseases in addition to powdery mildew must be controlled, PROCURE 480SC can be tank mixed with other fungicides registered for those diseases. Thorough coverage of the foliage is necessary for maximum disease control. Repeated applications should be made if conditions of powdery mildew development persist.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Cucumbers Gherkins Gourds Muskmelons (includes cantaloupes, honeydews, Persian melons and other melons) Pumpkins Squash (summer and winter) Watermelons	Powdery Mildew (<i>Erysiphe cichoracearum</i> , <i>Podosphaera xanthii</i>)	4 to 8	Apply PROCURE 480SC as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin at the start of vining or the first indication of disease. Repeated applications can be made at 7 to 14 day intervals. When conditions are favorable for severe disease pressure, use higher rates shorter intervals. Do not apply more than 40 fl. oz. of PROCURE 480SC per acre per season. Applications can be made up to the day of harvest. For greenhouse grown cucurbits apply 4 to 8 fl. oz. in 100 gallons of water.

GRAPES

PROCURE 480SC should be used as a protectant for control of powdery mildew. Start applications before bloom and continue on a 14 to 21 day interval at 4 to 8 fl. oz/acre. For the most susceptible varieties, use 6 to 8 fl. oz/acre and/or shorter spray intervals. When disease pressure is low, use 4 to 6 fl. oz/acre and/or the spray interval can be extended up to 21 days. Select spray volume, nozzle adjustments and ground speed to ensure complete coverage of fruit clusters and foliage. Use a minimum of 50 gallons of water per acre for early season applications. Use enough volume to ensure thorough coverage as the amount of foliage increases. Do not apply within 7 days of harvest. Do not exceed 32 fl. oz. PROCURE 480SC per acre per season. Do not apply by air. Sulfur at recommended use rates can be applied before, with, and/or between PROCURE 480SC applications.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Grapes	Powdery Mildew (<i>Uncinula necator</i>)	4 to 8	Start applications before bloom and continue on a 14 to 21 day interval. Use higher rates and/or shorter intervals on susceptible varieties or under heavy disease pressure. PROCURE 480SC applied in a powdery mildew spray schedule, will enhance the activity of registered fungicides used for the control of Botrytis bunch rot. Applications must be made on a 14-day sprayschedule for suppression of Botrytis bunch rot.

HAZELNUTS (FILBERTS)

Application may be made by ground or air. Minimum recommended spray volumes are 15 gallons per acre for aerial applications and 100 gallons per acre for ground applications. Use the highest labeled rate by air. Apply PROCURE 480SC in sufficient spray volume for thorough coverage. Tank mixing the lowest labeled rate of a spray surfactant with PROCURE 480SC may improve spray coverage and penetration. Ground application is preferred.

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Hazelnuts (Filberts)	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	4 to 6	PROCURE 480SC should be applied at budbreak (when the first green tissue is visible) as a preventative spray. Continue applications of PROCURE 480SC at 10 to 14 day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during bud break and rapid shoot elongation. The expanding shoot tip must be protected. Apply the high rate to varieties that are highly susceptible to the indicated disease or when severe disease conditions exist. Do not apply more than 24 fl. oz. of PROCURE 480SC per acre per season. Do not apply within 18 days of harvest. PROCURE 480SC may be applied as a tank mix or alternated with non-DMI fungicides as a resistance management strategy.

HEAD AND STEM BRASSICAS

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Head & Stem Brassicas: Broccoli Broccoli, Chinese (gai lon) Brussels sprouts Cabbage Cabbage, Chinese (napa) Cauliflower Cavalo broccolo Kohlrabi Mustard, Chinese (gai choy)	Powdery Mildew <i>(Erysiphe polygoni)</i> Black Spot / Alternaria leaf spot <i>(Alternaria spp.)</i>	6 to 8	Apply PROCURE 480SC as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin at the first indication of disease. Use higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Repeated applications should be made on a 14 day schedule. Do not apply more than 18 fl. oz. of Procure 480SC per acre per season. Use a minimum of 10 gallons per acre spray volume for ground applications. Use a minimum of 2 gallons per acre spray volume for aerial applications. Applications can be made one day before harvest.

HOPS

CROP	DISEASE	Rate of Procure 480SC fl. oz/acre	COMMENTS
Hops	Powdery Mildew <i>(Podosphaera macularis)</i>	12	Apply PROCURE 480SC as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin prior to disease development or at the first indication of the disease. Use higher rates for heavier infections and under conditions in which thorough coverage is more difficult. Repeated applications should be made on a 14 day schedule. Do not apply more than 36 fl. oz. of PROCURE 480SC per acre per season. Use a minimum of 50 gallons per acre spray volume for ground applications or a minimum of 10 gallons per acre spray volume for aerial applications. Applications can be made up to 7 days before harvest. Do not graze livestock in treated areas or harvest crops grown in treated areas for silage or for hay.

LEAFY GREENS (including field and greenhouse grown)

PROCURE 4805C can be used to control powdery mildew and black spot/Alternaria leaf spot on the following brassica and non-brassica leafy greens. If other leaf diseases in addition to powdery mildew or black spot/Alternaria leaf spot must be controlled, PROCURE 4805C can be tank mixed with other fungicides registered for those diseases. Repeated applications should be made if conditions of powdery mildew or black spot/Alternaria leaf spot development persist.

CROP	DISEASE	Rate of Procure 4805C fl. oz/acre	COMMENTS
Leafy Greens (except spinach) and Swiss chard Amaranth Arugula Chervil Chrysanthemum, edible-leaved & garland Cilantro Corn salad Cress, garden & upland Dandelion Dock Endive Lettuce (head and leaf) including Cos (Romaine) & Butterhead varieties (both field and greenhouse) Orach Parsley Purslane, garden and winter Radicchio Swiss chard (including all cultivars and/or hybrids of these)	Powdery Mildew (<i>Erysiphe</i> spp.) Black Spot/ Alternaria leaf spot (<i>Alternaria</i> spp.)	6 to 8	Apply PROCURE 4805C as a foliar spray in sufficient water to provide thorough coverage of the foliage. Applications should begin prior to disease development or at the first indication of the disease. Use higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Repeated applications should be made on a 14 day schedule. Do not apply more than 18 fl. oz. of PROCURE 4805C per acre per season. Application may be made by air. Use a minimum of 10 gallons per acre spray volume for ground applications and 2 gallons spray volume for aerial applications. Applications can be made up to the day of harvest (zero day PHI). For greenhouse grown leafy greens apply 6 to 8 fl. oz. in 100 gallons of water.

PAPAYAS

CROP	DISEASE	Rate of Procure 4805C fl. oz/acre	COMMENTS
Canistel Mango Papayas Sapodilla Sapote, black Sapote, mammey Star apple	Black Spot (<i>Asperisporium caricae</i>) Powdery Mildew (<i>Oidium caricae</i>)	8 to 10	Apply PROCURE 4805C as a foliar spray in sufficient water and use appropriate surfactant to provide thorough coverage of the foliage. Applications should begin prior to disease development or at the first indication of the disease. Use higher rates for heavier infections and under conditions in which thorough coverage is more difficult. Repeated applications should be made on a 14 day schedule. Do not apply more than 24 fl. oz. of PROCURE 4805C per acre per season. Use a minimum of 75 gallons per acre spray volume for ground applications. Application may be made by air. Use a minimum of 2 gallons of water per acre for aerial applications. Applications can be made up to the day of harvest (zero day PHI).

PINEAPPLES

PROCURE 480SC can be used as a pre-plant seed piece material treatment or as a post-harvest fruit treatment to control butt rot in pineapples.

CROP	DISEASE	Rate of Procure 480SC fl. oz./100gal.	DISEASE PRESSURE	COMMENTS
Pineapples (pre-plant seed piece material treatment)	Butt Rot (<i>Chalara paradoxa</i>)	4 to 8 8 to 12 12 to 16	Light Moderate Severe	Apply PROCURE 480SC at 4-16 fl. oz. in 100 gallons of water as a planting material dip treatment. Freshly removed seed material used for immediate planting should be immersed within 12 hours of removal for thorough wetting and coverage. Use a higher label rate under moderate to severe disease pressure. Do not use treated green forage or green fodder for animal feed.
Pineapples (post-harvest fruit treatment)		8 to 16		Apply specified dosage in 100 gallons of water with appropriate oil or wax after harvest. Use higher rates when disease history in the field has been known to be moderate to severe and when a susceptible variety is used. Fruit should be dipped or sprayed for thorough coverage with PROCURE 480SC + wax or PROCURE 480SC + oil and allowed to drain. Fruit discarded from fresh packing operation may be used for processing. Cannery waste may be fed.

STRAWBERRIES

PROCURE 480SC can be applied for control of strawberry powdery mildew with dilute or concentrate ground equipment. Apply PROCURE 480SC in sufficient water to ensure complete and thorough coverage of foliage, flowers and fruit. Use higher recommended application rates and shorter spray intervals on more susceptible varieties and/or under higher disease pressure. PROCURE 480SC may be tank mixed with other materials registered for use on strawberries. Apply tank mixtures in accordance with all applicable label directions. Do not apply within 1 day of harvest. Do not exceed 32 fl. oz. PROCURE 480SC per acre per season.

CROP	DISEASE	Rate of Procure 480SC fl. oz./acre	COMMENTS
Strawberries	Powdery Mildew (<i>Sphaerotheca macularis</i> f. sp. <i>fragariae</i>)	4 to 8	Protection of leaves before bloom will reduce in-season disease pressure and result in greater control of flower and fruit infections. Initiate applications at the first sign of powdery mildew. Continue applications at approximately 14 day intervals while conditions favor disease development.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer, to the extent allowed by law, assumes the risk of any such use.

© PROCURE is a Registered Trademark of Chemtura Corporation
 © Copyright 2009, Chemtura Corporation