

400-544

9/6/2012

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Allen R. Blem
Chemtura Corporation
199 Benson Rd
Middlebury, CT 06749

SEP 06 2012

Subject: Label Amendment to Rancona 3.8 FS
EPA Registration No. 400-544
Decision No. 468664
Submission Date: 8/6/12

Dear Mr. Blem:

The labeling referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended to add a pest (*Verticillium dahliae*) to the spinach directions for use and revise the application rate for that pest, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions, please contact Dominic Schuler at (703) 347-0260 or via email at schuler.dominic@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Tony Kish".


Tony Kish
Product Manager 22
Fungicide Branch
Registration Division (7504P)

RANCONA[®] 3.8 FS

SEED TREATMENT FUNGICIDE

Active Ingredient:		
ipconazole (CAS No. 125225-28-7).....	40.7%*	
Other Ingredients:	59.3%	
Total :.....	100.0%	

- *Contains 3.77 pounds of active ingredient per gallon.
- *Contains 452 grams of active ingredient per liter.



KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
MEDICAL EMERGENCY	800-292-5898
TRANSPORTATION EMERGENCY (CHEMTREC)	800-424-9300
PRODUCT SAFETY DATA (MSDS)	866-430-2775
For PRODUCT USE INFORMATION: Call 800-243-2850	

Net Contents:

EPA REG. NO. 400-544
EPA EST. NO.
013/

Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749



www.ChemturaAgroSolutions.com

NOTE: **This product is undyed.** The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

SEED TREATMENT EQUIPMENT


RANCONA 3.8 FS may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound and well-cured before treatment. Refer to the label rates below. Product requires dilution prior to application. Contact your local Chemtura representative or supplier for specific recommendations.

TREATED SEED LABELING

Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This bag contains seed treated with ipconazole. Do not use for Food, Feed, or Oil Purposes. Persons opening this bag or loading/pouring the treated seed must wear a long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves. After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. *Exception:* Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction."

"Do not graze or feed livestock on treated areas or hay grown from treated peanut seed."



SOYBEANS

For protection against seed rot, damping off and seedling blight including *Fusarium* (seed and soil borne), *Rhizoctonia solani*, and seed borne *Diaporthe* (Phomopsis). Also protects against general seed rots caused by saprophytic organisms such as *Penicillium* and *Aspergillus*, and protects against seed-borne *Botrytis* and *Sclerotinia*. Apply 0.085 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

PEANUTS

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.085 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

Do not graze or feed livestock on treated areas or on hay grown from treated seed.

ROOT VEGETABLES

Beet (sugar and garden), burdock (edible), carrot, celeriac, chervil (turnip-rooted), chicory (roots), chufa, dandelion (roots), ginseng, parsley (turnip-rooted), parsnip, radish, radish (oriental), rutabaga, salsify, salsify (black), salsify (Spanish), skirret, turnip and yam bean.

For protection against seedborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed or propagating root or tuber material.

SUNFLOWER

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.085 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

BRASSICA (COLE) LEAFY VEGETABLES

Broccoli, broccoli (Chinese), broccoli raab, Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa, and Chinese mustard), cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens.

For protection against seedborne and soilborne fungi that cause seed decay, damping off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

CUCURBIT VEGETABLES

Chayote, Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Momordica, all muskmelon, pumpkin, squash and watermelon.

For protection against seedborne and soilborne fungi that cause seed decay, damping off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

DRIED SHELLED PEAS AND BEANS

Dried cultivars of bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean; tepary bean; bean (*Vigna* spp.) (includes adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum* spp.) (includes field pea); pigeon pea.

For protection against seed rot, damping off and seedling blight including *Fusarium* (seed and soil borne) and *Rhizoctonia solani*. Also protects against general seed rots caused by saprophytic organisms such as *Penicillium*, and *Aspergillus*, and protects against seed-borne *Botrytis* and *Sclerotinia*. Provides suppression of seed borne Anthracnose of dry bean. Apply 0.085 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

LEAFY VEGETABLES (Except Brassica Vegetables)

Amaranth (leafy, Chinese spinach, tampala), arugula, cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (Florence, finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach, spinach (New Zealand), spinach (vine), and Swiss chard.

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

Spinach: For control of seed borne *Verticillium dahliae* to prevent the introduction of this disease into soils through infected seed, as well as for protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight. Apply 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

MUSTARD, RAPE, CANOLA

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, as well as for protection against *Phoma lingam* causing blackleg, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

ORNAMENTAL FLOWERS

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

CONIFERS

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

TURF GRASS

For protection against seedborne and soilborne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.085 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

COTTON

For protection against seed and soil borne fungi that cause seed decay, damping-off, and seedling blight, apply 0.051 - 0.34 fl. oz. of RANCONA 3.8 FS per 100 pounds of seed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate, dry and out of direct sunlight. Avoid excess heat. Do not freeze. ALWAYS SHAKE OR STIR WELL BEFORE USE.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Cover or incorporate spilled treated seeds.

EQUIPMENT CLEANING: Following applications, all seed treatment equipment should be thoroughly rinsed with water. Do not contaminate water by disposal of equipment washwater or rinsate.

CONTAINER DISPOSAL:

Nonrefillable Container

Plastic Containers: Do not reuse or refill this container. Triple rinse (or equivalent) container promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows:

For containers with capacity equal to or less than 5 gallons or 50 pounds: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

For containers with capacities greater than 5 gallons or 50 pounds: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Recycling

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or <http://www.acrecycle.org>.

IMPORTANT NOTICE - Chemtura Corporation warrants that this product conforms to the chemical description and is reasonably fit for the purposes stated on the label only when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond the control of Chemtura, Chemtura makes no claims or guarantees as to germination of carry-over seed.

RANCONA® is a registered trademark of Chemtura Corporation. Chemtura AgroSolutions™ and Belmont™ 2.7 FS are trademarks of Chemtura Corporation. ©Copyright 2012, Chemtura Corporation

