

400-571

7/6/2009

1087



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

400-571

Date of Issuance:

JUL 6 2009

NOTICE OF PESTICIDE:

- X Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Rancona Xtra

Name and Address of Registrant (include ZIP Code):

Chemtura Corporation
199 Benson Road, 2-5
Middlebury, CT 06749

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Signature of Approving Official:

Mary J. Waller

Mary Waller, Product Manager (21)
Fungicide Branch, Registration Division (7505P)

Date:

7/6/2009

JUL 6 2009

2 of 7

2. Submit the following conditional data before the due date of 12/31/2010:
 - a. Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies.
3. Make the following changes to the label:
 - a. Change the product registration number to "EPA Reg. No. 400-571"
 - b. If you wish you may delete the signal word CAUTION from page 1 and from the **HAZARDS TO HUMANS AND DOMESTIC ANIMALS** section
 - c. If you wish you may delete the First Aid statements
 - d. If you wish you may delete the following statements from the section **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**: "Causes moderate eye irritation. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Avoid breathing vapor." If you wish to continue to use these warning statements delete "Wear protective eyewear."
 - e. On page 2 in the **GENERAL** section add the following sentence to the last paragraph: "Do not tank mix with any pesticide that has a prohibition against tank mixing."
 - f. On page 3 in the Note below the table for **Tank Mix with a Metalaxyl Product**:
 1. Delete the sentence "The seed treatment products may be mixed together before application."
 2. Change the first sentence of the **Note** to read "MAXIMUM USAGE when applying multiple metalaxyl and/or mefenoxam containing products to the same crop...."
 - g. On page 4 in the **STORAGE AND DISPOSAL** box move the section **EQUIPMENT CLEANING** to the end of the **SEED TREATMENT EQUIPMENT** section on page 2.
4. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Note: Should you wish to retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website

30f7

JUL 6 2009

contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance. As an alternative, you may refer consumers to the company's phone number or e:mail address.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.




Mary Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure:

Label stamped "Accepted with Comments"
Product Chemistry Review DP361377 dated June 2, 2009
Acute Toxicity Review DP361378 dated June 2, 2009

4097

RANCONA™ Xxtra



SEED PROTECTANT

Net Contents:

Active Ingredients: (% by weight)	
ipconazole (CAS No. 125225-28-7).....	1.029%*
metalaxyl (CAS No. 57837-19-1).....	1.647%*
Other Ingredients:	97.324%
Total:.....	100.000%

*Contains 0.0916 lb. ipconazole and 0.147 lb. metalaxyl per gallon.

*Contains 11 grams ipconazole and 17.6 grams metalaxyl per liter.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. • Call a poison control center or doctor for further treatment advice.
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
EMERGENCY PHONE	800-292-5898
SAFETY DATA AND INFORMATION	866-430-2775
TRANSPORTATION EMERGENCY (CHEMTREC)	800-424-9300

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JUL 6 2009
**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No. 400-571**

Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749
EPA REG. NO. 400-571
EPA EST. NO.
001



www.chemtura.com

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Avoid breathing vapor. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistance category selection chart.

Applicators and Other Handlers Must Wear: Long-sleeved shirt and long pants; chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or Viton; and shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seeds. Do not contaminate water when disposing of equipment washwater or rinsate.

Ipconazole may persist in soil, and long-term use in the same field may result in an accumulation of active ingredient.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours.**

Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or Viton; shoes plus socks.

GENERAL

RANCONA™ Xxtra is a systemic and contact broad-spectrum fungicide for seed treatment that protects against a wide variety of diseases. RANCONA Xxtra is effective against *Fusarium*, *Diaporthe (Phomopsis)*, *Rhizoctonia*, *Sclerotinia* and *Pythium*. It is also effective against seed rot fungi such as *Aspergillus* and *Penicillium*. Seed should be sound and well cured before treatment.

RANCONA Xxtra may be used in combination with a metalaxyl containing product (such as MetaStar™ ST) for enhanced *Phytophthora* protection.

Do not use RANCONA Xxtra in combination with other seed treatment products unless their compatibility has been verified. Always read and follow carefully all label directions and precautions of each combination product. When using combinations of products, the most restrictive of label limitations and precautions must be followed.

SEED TREATMENT EQUIPMENT

RANCONA Xxtra may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. This product is for both commercial and on-farm application.

Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound and well cured before treatment. Refer to the label rates below.

In most cases, RANCONA Xxtra is ready to use and may be applied to seed as-is. However, to achieve more uniform product to seed coverage, dilution with water may be appropriate for some types of treaters and/or when treating under dry and/or hot conditions. Mix RANCONA Xxtra and water thoroughly into a slurry before treating seed. Recalibrate treating equipment to compensate for the addition of water. The amount of dilution water that is most suitable depends on the seed, the treater type and the temperature. Use a total slurry application volume of 3.5 fl. oz. (that is, as-is) to 15 fl. oz. per 100 lb. of seed. Contact your local Chemtura representative or supplier for specific recommendations.

APPLICATION INSTRUCTIONS

Application rate (all crops): 3.5 fl.oz./100 pounds of seed

(This equals 0.04 oz. ipconazole plus 0.064 oz. metalaxyl per 100 pounds)

Crop	Symptoms Controlled (Disease Organism)	Symptoms Suppressed (Disease Organism)
Soybeans	Seed rot, damping off and seedling blight, (including seed- and soil-borne <i>Fusarium</i> , and seed-borne <i>Diaporthe</i> (<i>Phomopsis</i>), <i>Rhizoctonia solani</i> and <i>Pythium</i>) General seed rots (including those caused by saprophytic organisms such as <i>Penicillium</i> and <i>Aspergillus</i>)	
Dried shelled beans and peas: Bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); Bean (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean; tepary bean); Bean (<i>Vigna</i> spp.) (includes adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; Pea (<i>Pisum</i> spp.) (includes field pea); pigeon pea.	Seed rot, damping off and seedling blight (seed- and soil-borne <i>Fusarium</i> , <i>Rhizoctonia solani</i> and <i>Pythium</i>) General seed rots (including those caused by saprophytic organisms such as <i>Penicillium</i> and <i>Aspergillus</i>) Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i>	Seed-borne <i>Anthracnose</i> of bean

Tank Mix with a Metalaxyl Product

RANCONA Xxtra provides control of *Pythium* and protection against *Phytophthora*. However, if the target field has a history of high *Phytophthora* pressure, RANCONA Xxtra may be applied in combination with a product containing metalaxyl (such as MetaStar™ ST) at the rate registered for your crop for increased protection. For example, use the following rates when mixing RANCONA Xxtra with MetaStar ST. Use a higher rate of MetaStar ST when the field has a history of higher *Phytophthora* pressure:

Product	Rate
RANCONA Xxtra + MetaStar ST	3.5 fl.oz/100 lbs. of seed + 0.17-0.53 fl.oz/100 lbs. of seed

NOTE: MAXIMUM USAGE when applying both metalaxyl and mefenoxam containing products to the same crop within the same season. DO NOT apply more than the maximum seasonal total for the active ingredient as stated on the label of the lowest seasonal total on that crop.

The seed treatment products may be mixed together before application. Agitate the slurry to maintain uniformity. Recalibrate your treating equipment to compensate for the addition of the metalaxyl product. Read and follow carefully all label directions and precautions of each combination product. When using combinations of products, the most restrictive of label limitations and precautions must be followed.

TREATED SEED LABELING

Seeds treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This package or bag contains seed has been treated with ipconazole and metalaxyl. Do not use for food, feed or oil purposes. Store away from feeds and foodstuffs. Persons opening this bag or package or loading/pouring the treated seed must wear a long-sleeved shirt, long pants, shoes and socks."

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. Care must be exercised in the handling of treated seed. Do not use augers used for handling treated seed to move seed for feed, food or oil processing. Do not re-use bags from treated seed to handle food or feed products.

IN CALIFORNIA: ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use.

ALWAYS SHAKE OR STIR PRODUCT WELL BEFORE USE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate, dry and out of direct sunlight. Avoid excess heat. Do not freeze.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Cover or incorporate spilled treated seeds.

EQUIPMENT CLEANING: Following applications, thoroughly rinse seed treatment equipment with water. Do not contaminate water by disposal of equipment washwater or rinsate.

CONTAINER DISPOSAL: Nonrefillable Container

Plastic containers: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Recycling

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

IMPORTANT NOTICE - Chemtura Corporation warrants that this product conforms to the chemical description and is reasonable fit for the purposes stated on the label only when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond the control of Chemtura, Chemtura makes no claims or guarantees as to germination of carry-over seed.

RANCONA™ is a trademark of Chemtura Corporation
MetaStar™ is a trademark of LG Life Sciences, Ltd.
© Copyright 2009, Chemtura Corporation