


400-576

0210412010

1/11

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	<p>EPA Reg. Number: 400-576</p>	<p>Date of Issuance: FEB 04 2010</p>
	<p>Term of Issuance: Conditional</p>	
	<p>Name of Pesticide Product: Attendant 480 FS</p>	
<p>Name and Address of Registrant (include ZIP Code): Chemtura Corporation 199 Benson Road, 2-5 Middlebury, CT 06749</p>		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p style="text-align: center;">This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data. 2. Make the following label changes before you release the product for shipment: <ul style="list-style-type: none"> • Revise the EPA Registration Number to read, "EPA Reg. No 400-576." • On page 4, under the crop table for Canola, revise the comment that reads: "In areas where flea beetles and foliar insects are in high numbers, the higher application rate is required" to read, "In areas where flea beetles and foliar insects are in high numbers, use the higher specified application rate" to make it consistent with other labeling. 		
<p>Signature of Approving Official:  Venus Eagle, Product Manager 01 Insecticide-Rodenticide Branch, Registration Division (7505P)</p>	<p>Date: FEB 04 2010</p>	

- On page 7, in the "Rotational Plant-Back Interval" box, move the bolded statement associated with the asterisk "For all other crops not listed on an imidacloprid label..." to the paragraph above the table, right before the last sentence "The following plant-back intervals must be observed for listed crops." Then remove the asterisks.
- On page 8, under the "Treated Seed Labeling" section, revise the second sentence to read: "Seed that has been treated with this product that is then packaged or bagged for future use must also contain..."
- On page 8, under the "Treated Seed Labeling" section, bullet each separate statement to make this section more clear and easy to read.
- On page 8, under the "Treated Seed Labeling" section, add the statement "Store seed away from food and feedstuff."

3. Storage stability (830.6317) and corrosion characteristics (830.6320) data must be submitted within 18 months from the date of this registration notice.

4. Oxidation/Reductions (830.6314) data must be submitted or cited within 18 months from the date of this registration notice.

5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure

ATTENDANT[®] 480 FS

SEED TREATMENT INSECTICIDE

Active Ingredient:
imidacloprid (CAS No. 138621-41-3).....40.23%*
Other Ingredients: 59.77%
Total: 100.00%

*Contains 4 lbs. active ingredient per gallon, or 480 grams active ingredient per liter

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
<p>EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-Hour Medical Emergency Assistance (Human or Animal), Call 1-800-292-5898. For Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call CHEMTREC at 1-800-424-9300. For Chemical Safety Data and Information, Call 1-866-430-2775 Toll Free, or Visit our Web Site at www.chemtura.com</p>	

Net Contents:

EPA File Symbol 400-LTA
EPA Est. No.
001/

ACCEPTED
With COMMENTS
In EPA Letter Dated:
FEB 04 2010

Manufactured for:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749

Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
pesticide Registered under EPA Reg. No:

(CHEMTURA logo)

400-376

www.chemtura.com

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately, from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This product is **highly toxic to birds and aquatic invertebrates**. Cover or incorporate spilled treated seeds. Treated seed exposed on soil surface may be hazardous to birds.

For On-Farm Use: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

For Commercial Seed Treatment Facilities: Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant discharge elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions and precautions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 12 hours**.

Exception: Once the seed is planted in soil or other planting media, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no contact with soil/subsurface.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves, shoes plus socks, protective eyewear.

ATTENDANT® 480 FS is a seed treatment insecticide that protects the labeled seeds and seedlings from injury from certain early-season insects. ATTENDANT 480 FS will also provide protection to seed in storage against damage from certain insects. Federal regulations have established official tolerances for pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use this product only at the rates contained on this label.

ATTENDANT 480 FS is formulated for both commercial and (for specific crops) agricultural establishment (on-farm) application. See below for Seed Treatment directions by crop, for application details. ATTENDANT 480 FS may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best insect protection and optimum performance. Only sound and well-cured seed are to be treated. Refer to the application instructions below for label rates and application details. ATTENDANT 480 FS will treat similarly to other concentrated imidacloprid seed treatment formulations. The specific treating process and equipment that is most suitable depend on factors such as the seed type, rate, treater type and temperature. Contact your local Chemtura representative or supplier for specific recommendations if assistance is required.

Seed Dye or Colorant: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 153.155.

ALWAYS MIX PRODUCT THOROUGHLY BEFORE USING.

Always pretest tank mixes to assure physical compatibility between formulations. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. Do not tank mix with any product that contains a label prohibition against tank mixing.

STORED SEED PROTECTION

Applied at labeled rates equal to or above 1.0 fluid ounces per hundredweight (fl. oz. / cwt.), ATTENDANT 480 FS will provide protection to seed against injury from the following insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*) and Lesser Grain Borer (*Rhizopertha dominica*) for all labeled crops. Fumigate seed that has an existing population of stored grain pests before treating with this product and bagging seed.

6/11

SEED TREATMENT FOR EARLY SEASON PROTECTION AGAINST CERTAIN SUCKING INSECTS

ATTENDANT 480 FS will aid in the early season protection of seeds and seedlings against injury by certain insects. In some cases, ATTENDANT 480 FS also provides protection against spread of certain viruses.

Canola:

For End-Use Application At Agricultural Establishments: Always mix this product thoroughly before using. Apply using equipment such as a HCBT or Eight-Bag Batch Treater. Apply 6.4 to 16 fl. oz. of ATTENDANT 480 FS per 50 pound bag of seed. This product may be diluted with an approved fungicide mixture for extended disease protection. When using a batch treater, treat one-half of seed with one-half of the slurry mix; then add the remainder of the seed and slurry, and continue mixing until all seed is thoroughly covered.

For Commercial Seed Treatment: Apply with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed.

Crop	Pest	Rate	Comments
Canola	Aphids, Flea beetles, Wireworms	12.8 – 32.0 fl. oz. per one hundred pounds (cwt.) of seed	Early season protection of seedlings. In areas where flea beetles and foliar insects are in high numbers, the higher application rate is required. Please consult your local agricultural extension service office for pest patterns, history and forecasts to assist in determining the appropriate rate for your region.
	Second generation Lygus bugs (including <i>Lygus</i> spp.); Cabbage seedpod weevil larvae (including <i>Ceutorhynchus</i> <i>assimilis</i>)	19.2 – 32.0 fl. oz./cwt.	For suppression only. Use the higher rate when insects are in high numbers. Please consult your local agricultural extension service office for pest patterns, history and forecasts to assist in determining the appropriate rate for your region.

Sorghum:

For End-Use Application At Agricultural Establishments: Always mix product thoroughly before using. Apply 4 fl. oz. of ATTENDANT 480 FS per 50 pounds of seed using equipment such as a HCBT or Eight-Bag Batch Treater. This product must be diluted for a slurry application rate of 8 to 10 fluid ounces of slurry per 50 pounds of seed. When using a batch treater, treat one-half of the seed with one-half of the slurry mix. Then add remainder of the seed and slurry, and continue mixing until all seed is thoroughly covered. To improve seed flow, add dry talc at the rate of 0.75 ounce per 50 pounds of seed after slurry treatment, and allow it to distribute evenly on the seed.

For Commercial Seed Treatment: Apply 8.0 fl. oz. of product per hundredweight (cwt.) seed using any mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed.

Crop	Pest	Rate	Comments
Sorghum	Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Fire ants ⁽¹⁾ , Wireworms	8.0 fl. oz./cwt.	Provides early season protection of seedlings against injury. Thorough coverage is necessary for optimal control. ⁽¹⁾ Note: Not to kill or control fire ants. Only for protection of seed. Restriction: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Wheat, Barley, Oats, Rye, Triticale:

For End-Use Application At Agricultural Establishments: Always mix product thoroughly before using. Apply using equipment such as a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.) or other on-farm seed treating equipment capable of accurately applying low rates of ATTENDANT 480 FS. Apply 1.0 – 3.0 fl. oz. per hundredweight (cwt.) of seed as a slurry treatment prior to planting. ATTENDANT 480 FS must be combined with a fungicide product for seed and seedling protection against fungal pathogens, as well as the listed insects. Depending on the fungicide product used, dilution with water may be necessary for optimum coverage. ATTENDANT 480 FS may also be applied on-farm to seed previously treated with a fungicide. In this case, dilution is necessary.

For Commercial Seed Treatment: ATTENDANT 480 FS may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed. For maximum effectiveness, seed must be treated uniformly and covered thoroughly.

Restriction: DO NOT feed or graze livestock on treated areas for 45 days after planting.

Crop	Pest	Rate	Comments
Wheat, Barley, Oats, Rye, Triticale	Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly, Wireworms	1.0 – 3.0 fl. oz./ cwt.	Provides early season protection of seedlings. Reduces possible spread of barley yellow dwarf virus disease from aphid vectoring. Use the higher rate to increase time of protection and/or when insect pressure is heavy.
	Wireworms	0.16 – 0.32 fl. oz./ cwt.	Provides suppression of activity on seed and young seedlings only. The 0.32 fl. oz./cwt. rate increases the time of protection and/or improves protection when insect pressure is high, compared with lower rates of imidacloprid.
	Grasshoppers	1.5 – 3.0 fl. oz./ cwt.	Treated seed may be planted as a 50- to 60-foot border around the perimeter of the field to help minimize early season damage from grasshoppers. Consult your local agricultural extension service office for grasshopper control information specific to your area. Use the higher rate to increase time of protection and/or when insect pressure is heavy.

Cotton (Delinted Seed Only):

For End-Use Application At Agricultural Establishments: Always mix product thoroughly before using. Apply using equipment such as a HCBT or Eight-Bag Batch Treater at rates listed in the table below. This product may be diluted/slurried with water or an approved fungicide for disease control. Mix slurry so that rate is 8 to 10 fluid ounces of diluted slurry per 50 pounds of seed. When using a batch treater, treat one-half of the seed with one-half of the slurry; then add the remainder of seed and slurry, and continue mixing until all seed is thoroughly covered.

For Commercial Seed Treatment: Apply at 0.375 mg. a.i. per seed or 16.0 fluid ounces per hundredweight of seed (16 fl. oz./cwt.), whichever is less. Apply using any mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed.

Crop	Pest	Rate	Comments
Cotton (delinted seed only)	Early season Thrips and Aphids	16.0 fl. oz./ cwt. or 0.375 milligram a.i./seed	Provides protection of seedlings against injury. Restriction: DO NOT apply more than a total of 0.5 lb. of imidacloprid per acre per cropping cycle, regardless of the type of application (seed treatment, soil, or foliar).

Sugar Beets:

For Commercial Seed Treatment Only: Apply at rates specified below with any mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed. **IMPORTANT:** if rate exceeds 3.0 fl. oz. per unit, seed must be pelleted with at least 2 parts of pelleting mixture per part of raw seed.

Crop	Pest	Rate ⁽¹⁾	Comments
Sugar Beets (commercial seed treatment)	Whiteflies, Aphids ⁽³⁾ , Leafhoppers ⁽³⁾ , Root aphids, Thrips, Wireworms	Pelleted seeded: 3.0 - 6.3 fl. oz. per seed unit ⁽²⁾ Non-pelleted raw seed: 3.0 fl. oz. per seed unit ⁽²⁾	Provides early season protection of seedlings. ⁽¹⁾ 3.0 to 6.3 fl. oz./seed unit may be applied if applied in or on a unit of pelleted sugar beet seed with a weight ratio of at least 2:1 pelleting mixture to raw seed . If applying to non-pelleted raw seed, apply in a film coat and do not exceed a rate of 3.0 fl. oz. per unit of seed. ⁽²⁾ 1 seed unit = 100,000 seeds, approximately 1 kilogram (2.2 lbs.) ⁽³⁾ Including reducing risk of spread of Curly top and Yellow mosaic virus by insect vectors.

Other Crops

For Commercial Seed Treatment Only: Soybeans, carrots and the peas, beans and pod vegetables specified below are to be treated only by commercial treaters. ATTENDANT 480 FS may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment is calibrated to accurately and uniformly apply the product to seed.

Crop	Pest	Rate	Comments
Soybean	Seed corn maggots, Soybean aphids, over-wintering Bean leaf beetles, suppress spread of certain viruses	2.0 – 4.0 fl. oz./cwt.	Protects planted seed from seed corn maggot. Reduces feeding damage caused by soybean aphids and over-wintering bean leaf beetles. Use higher rate to provide increased length of protection and for heavy insect pressure. ATTENDANT 480 FS can be used as an over-treatment. Restriction: DO NOT graze or feed livestock on soybean forage or hay grown from treated seed.

Adzuki bean, Asparagus bean, Broad bean (succulent or dry), Catjang bean, Chinese long-bean, Field bean, Guar Bean, Jackbean, Kidney bean, Lablab Bean, Lima bean (succulent or dry), Moth bean (succulent or dry), Mung bean, Navy bean, Pinto bean, Rice bean, Runner bean, Snap bean, Sword bean, Tepary bean, Urd bean, Wax bean, Yardlong bean, Blackeyed bean (succulent or dry), Chickpea, Cowpea (succulent or dry), Crowder pea, Dwarf pea, Edible-Pod pea, English pea, Field pea, Garden pea, Green pea, Pigeon pea (succulent or dry), Snow pea, Southern pea (succulent or dry), Sugar snap pea, Grain lupine, Sweet lupine, White lupine, White sweet lupine, Lentil*	Wireworms, Bean leaf beetles, Imported fire ants ⁽¹⁾ , Aphids	2.0 – 4.0 fl. oz./cwt.	Provides early season protection of seedlings against injury. *Restriction: Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California. ⁽¹⁾ Note: Not to kill or control fire ants. Only for protection of seed.
Carrot	Seed corn maggots, Wireworms	8.0 fl. oz./cwt.	Provides early season protection of seedlings against injury.

ALWAYS MIX THIS PRODUCT THOROUGHLY BEFORE USING

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Rotational Plant-Back Intervals*				
Immediate Plant-Back				30-Day Plant-Back
Artichoke Barley Borage Brassica (cole) Leafy Vegetables Canola Carrot Cilantro Corn, Field Corn, Sweet Cotton Cranberry	Crambe Cucurbits Eggplant Flax Ground cherry Leaf Petiole Vegetables Leafy Vegetables Legume Vegetables (succulent or dried, except soybean)	Millet Mustard Seed Oats Okra Pepinos Pepper Popcorn Potato Rapeseed Root and Tuber Vegetables	Rye Safflower Sorghum Soybean Strawberry Sugar beet Sunflower Tomatillo Tomato Triticale Watercress Wheat	Buckwheat Rice

*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

*For all other crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient has been established, a 12-month plant-back interval must be observed.

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

IMPORTANT – USE RESTRICTIONS

- Treated seed must not be used for or mixed with food or animal feed or processed for oil.
- **Wheat, Barley, Oats, Rye, Triticale and Sorghum:** Do not graze or feed livestock on treated areas for 45 days after planting.
- **Cotton (delinted seed only):** Regardless of the type of application (seed treatment, soil, or foliar), do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle.
- **Seed-and-pod vegetable seed treated in California** must be destined for planting in states other than California and is not to be planted in California.

TREATED SEED LABELING

Seed commercially treated with Attendant 480 FS Seed Treatment must be labeled in compliance with all the requirements of the Federal Seed Act. Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This bag contains seed treated with imidacloprid. Treated seed must not be used for or mixed with food or animal feed or processed for oil. Persons opening this bag or loading/pouring the treated seed must wear a long-sleeved shirt, long pants, socks and shoes, and chemical resistant gloves."

"Dispose of seed packaging in accordance with local requirements."

"Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 12 hours.**"

"*Exception:* Once the seed is planted in soil or other planting media, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no contact with soil/subsurface."

"PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves, shoes plus socks, protective eyewear."

"This product is **highly toxic to birds and aquatic invertebrates.** Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds."

"Wheat, Barley, Oats, Rye, Triticale, and Sorghum: Do not graze or feed livestock on treated areas for 45 days after planting."

"Cotton (delinted seed only): Do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle, regardless of the type of application (seed treatment, soil, or foliar)."

"Soybean: Do not graze or feed livestock on soybean forage or hay grown from treated seed."

"Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California."

ROTATIONAL PLANT-BACK INTERVALS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Immediate Plant Back: Artichoke; Barley; Borage; Brassica (cole) Leafy Vegetables; Canola; Cilantro; Corn, Field; Corn, Sweet; Cotton; Cranberry; Crambe; Cucurbits; Eggplant; Flax; Ground cherry; Leaf Petiole Vegetables; Leafy Vegetables; Legume Vegetables (succulent or dried, except soybean); Millet; Mustard Seed; Oats; Okra; Pepino; Pepper; Popcorn; Potato; Rapeseed; Root and Tuber Vegetables; Rye; Safflower; Sorghum ; Soybean; Strawberry; Sugarbeet; Sunflower; Tomatillo; Tomato; Triticale; Watercress; Wheat .

30-Day Plant-Back: Buckwheat; Rice

12-Month Plant Back: For all other crops not listed on this product label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed. Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed."

11/11

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate, dry and out of direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container

Plastic Containers: Do not reuse or refill this container. Triple rinse (or equivalent) container promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows:

For containers with capacity equal to or less than 5 gallons or 50 pounds: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

For containers with capacities greater than 5 gallons or 50 pounds: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Recycling

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

IMPORTANT NOTICE - Seller warrants that this product conforms to the chemical description and is reasonably fit for the purposes stated on the label only when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

Treatment of highly mechanically damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond the control of Chemtura Corporation, Chemtura Corporation makes no claims or guarantees as to germination of carry-over seed.

ATTENDANT® is a registered trademark of Chemtura Corporation
Copyright 2010, Chemtura Corporation
001 ver 020110