400-582 Term of Issuance: Conditional Name of Pesticide Proc Prevamite TM O ation must be submitted to ar t always refer to the above E ered/reregistered under the Fe the Agency. In order to proto of a pesticide in accordance s giving the registrant a right	nd accepted by the PA registration number. ederal Insecticide, Fungio tect health and the with the Act. The accept
Conditional Name of Pesticide Proc Prevamite TM O ation must be submitted to art t always refer to the above E ered/reregistered under the Fe the Agency. In order to proto of a pesticide in accordance	nd accepted by the PA registration number. ederal Insecticide, Fungio tect health and the with the Act. The accept
Prevamite TM O ation must be submitted to an t always refer to the above E cred/reregistered under the Fe the Agency. In order to proto of a pesticide in accordance	nd accepted by the PA registration number. ederal Insecticide, Fungio tect health and the with the Act. The accept
t always refer to the above E ered/reregistered under the Fe the Agency. In order to prot of a pesticide in accordance	PA registration number. ederal Insecticide, Fungio tect health and the with the Act. The accept
t always refer to the above E ered/reregistered under the Fe the Agency. In order to prot of a pesticide in accordance	PA registration number. ederal Insecticide, Fungio tect health and the with the Act. The accept
the Agency. In order to proto of a pesticide in accordance	tect health and the with the Act. The accept
of a pesticide in accordance	with the Act. The accept
a FIFRA section 3(o	c)(7)(A) provided
gistration/registrati gistrants of similar j	
e product for shipn	nent:
Reg. No. 400-582.	"
Date:	
SEP	1 6 2010
	gistration/registrati gistrants of similar e product for shipn Reg. No. 400-582. Date:

b. A's stated in 40CFR156.140(a)(4), a batch code must be included on the label or package for container and containment for nonrefillables.

3. Storage stability (830.6317) and corrosion characteristics (830.6320) data must be submitted within 18 months of the registration notice for Prevamite, EPA Reg. No. 400-572, dated May 14, 2010.

4. Note: Should you wish to retain a reference to the company's website on your label, please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

5. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions regarding this notice, please contact Jennifer Gaines at (703) 305-5967 or via e-mail, <u>gaines.jennifer@epa.gov</u>.

A stamped copy of the label is enclosed for your records.

John Hebert Product Manager (07) Insecticide-Rodenticide Branch Registration Division (7505P)

Enclosure

bofle

with CC...IMENTS In EPA Letter Dated SEP 1 6 2010 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-582

TED

ACO

GROUP	6	ACARICIDE
GROUP	un	ACARICIDE

MITICIDE/INSECTICIDE

Prevamite

Active Ingredient: (% by weight)	
bifenazate: hydrazine carboxylic acid, 2-(4-methoxy-	
[1,1'-biphenyl]-3-yl) 1-methylethyl ester (CAS No. 149877-41-8)43.2%*	
Abamectin (CAS No. 71751-41-2)	
Other Ingredients:	
Total	2
*Contains 4.0 lbs. bifenazate per gallon.	4

τм

**Contains 0.20 lbs. abamectin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

	FIRST AID	
IF SWALLOWE	 Call a poison control center or doctor treatment advice. 	•
	 Have person sip a glass of water if ab 	
	 Do not induce vomiting unless told to center or doctor. Do not give anything to an unconscionation 	
IF INHALED	 Move person to fresh air. If person is not breathing, call 911 or artificial respiration, preferably by m Call a poison control center or doctor 	outh-to-mouth, if possible
	SSISTANCE: Have the product container or location of the second s	abel with you when calling
	ENCY PHONE	800-292-5898
EMERG		
	DATA/INFORMATION	866-430-2775

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure. gallons This box contains 2x2.5 gallon containers

Net Contents: 5

or

4

gallons This box contains 4x1 gallon containers

> or 8

quarts This box contains 8x1 quart

containers

or 64

ounces This box contains 8x8 ounce containers

EPA REG. NO. 400-EPA EST. NO. 002 Manufactured for: Chemtura Corporation 199 Benson Road Middlebury CT 06749



www.chemtura.com

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING/AVISO

May be fatal if swallowed. Harmful if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves (made of any waterproof material for mixing/loading activities)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, estuarine/marine invertebrates, fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short pants and short-sleeved shirt
- Chemical-resistant gloves made of any waterproof material
- · Chemical-resistant shoes and socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONARY STATEMENTS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR MITE/INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

USE INFORMATION

Prevamite O is a suspension concentrate (flowable) that will control the pests on the crops listed on this label when it is applied according to the DIRECTIONS FOR USE. Thorough coverage of foliage is essential for good mite and insect control.

NOTE:

- Adjuvant: Residual control of pests may be enhanced with the addition of a horticultural spray oil at 0.5 to 1.0% of the spray volume on field-grown woody ornamentals, landscape plants and Christmas trees. Some plants are sensitive to oils and so without prior experience the user should spray a small number of plants and observe plants for 2 weeks before spraying the remaining plants. Excessive cold or warm temperatures may increase the chance of plant damage following application with oils. Carefully read and follow directions on the oil label and do not exceed maximum rates listed on either label.
- **Phytotoxicity:** The user should pre-test any proposed mixtures with **Prevamite O** to ensure the physical compatibility and lack of phytotoxic effects.

Resistance Management

Prevamite O contains the active ingredients abamectin and bifenazate. Because of the inherent risks of pests developing resistance to any product, it is strongly advised that **Prevamite O** be used in a sound resistance management program. Treatment may not be effective against labeled pests if insect or mite-tolerant strains develop. When applying **Prevamite O** to plants that are hosts of labeled pests with multiple generations per crop per year, use resistance management practices.

Resistance management practices include, but are not limited to:

- rotating **Prevamite O** with other products with different modes of action,
- not treating successive generations with products based on abamectin or other Group 6 acaricides, or the bifenazate group,
- always use labeled rates,
- · using non-chemical alternatives such as beneficial arthropods,

• ensure thorough spray coverage to all foliage For additional information regarding the implementation of these or other resistance management practices, consult your local agricultural advisor or company representative.

USE DIRECTIONS

ORNAMENTALS

ø

Prevamite O provides quick knockdown and long residual control of a wide variety of mite pests listed below on the following plants:

All ornamental plants, including bedding plants, flowering plants, foliage plants, bulb crops, perennials and trees and shrubs.
 All non-bearing fruit trees which will not bear fruit for a minimum of 12 months.

Prevamite O can be used in the following areas where these plants are grown and/or maintained in containers or in the ground: • Greenhouses and shadehouses

- Nurseries, including Christmas tree / conifer plantations
- Landscapes
- Interiorscapes

NOT INTENDED FOR HOMEOWNER USE

Target Pests	Application Rates Fl. Ozs. Per 100 Gals. Water	Application Instructions
<u>Mites:</u> Bamboo spider mite Broad mite Bud mite Carmine spider mite Citrus red mite Clover mite Cyclamen mite European red mite Lewis mite Pacific spider mite Rust mite Southern red mite Spruce spider mite Strawberry mite Two-spotted spider mite	3 - 4	Apply in sufficient volume of water to ensure thorough coverage of the foliage. Actual spray volume will vary depending on the size of the plants being treated. Do not exceed spray volumes of 300 gals./A. Begin applications as soon as mites appear. One additional application can be made after 7 days or as necessary to maintain control. Do not make more than two (2) applications per crop per year.
<u>Leafminers:</u> Boxwood Liriomyza	6	Apply in sufficient volume of water to ensure thorough coverage of the foliage. Actual spray volume will vary depending on the size of the plants being treated. Do not exceed spray volumes of 200 gals./A. For control of mining larvae, apply when adults are beginning to lay eggs on the new foliage.
Suppression only: Aphids Thrips Whiteflies		For suppression of pest populations, young immatures must be contacted by the spray. One additional application can be made after 7 days or as neces- sary to maintain control. Do not make more than two (2) applications per crop per year.

USE RESTRICTIONS

• Do not use in citrus nurseries.

• Apply by ground equipment including compressed air sprayers, hydrayulic sprayers, ground boom sprayers, air blast sprayers, hose-end sprayers. Do not apply aerially.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dryer formulations above.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING: Nonrefillable containers: 2.5 gallons, 1 gallon, 1 quart or 8 ounces. Do not reuse or refill this container. Triple rinse (or equivalent) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

If pressure rinsing: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration. Do not burn, unless allowed by state and local ordinances.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

IMPORTANT NOTICE—Sellers warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and to the extent consistent with applicable law, the buyer assumes the risk of any such use.

™PREVAMITE and Chemtura AgroSolutions™ are Trademarks of Chemtura Corporation ©Copyright 2010, Chemtura Corporation