


First Aid Treatment

SYMPTOMS OF POISONING

IF SWALLOWED: ...
IF IN EYES: ...
IF ON SKIN: ...



KEEP OUT OF REACH OF CHILDREN
DO NOT GET INTO EYES, NOSE, OR MOUTH
IF SWALLOWED, DO NOT INDUCE VOMITING
IF IN EYES, FLUSH WITH WATER
IF ON SKIN, WASH WITH SOAP AND WATER

ABSORBED THROUGH THE SKIN.
MAY BE FATAL IF SWALLOWED.

DYANAP (H.W.)
KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

NO ACCEPTANCE

ACC. No 400-6530

DYANAP D NITRO

ONE LIQUID GALLON

CAUTION

STOPE ABOVE

SHAKE WELL BEFORE USE
DO NOT USE, POUR, SPILL,
OR STORE NEAR HEAT OR
OPEN FLAME

06530

UNIROYAL

FOR MORE INFORMATION

GENERAL INFORMATION

PEANUTS

PEANUTS—PREEMERGENCE TO CRACKING-TIME
TANK MIXTURE
DYANAP H.W. PLUS LASSO
EMULSIFIABLE CONCENTRATE

SOYBEANS



DYANAP (H.W.)



Kills annual weeds and grasses in peanuts and soybeans.

Do not get in eyes, on skin, on clothing, avoid breathing spray drift. Do not take internally. Do not wear contaminated clothing or shoes. Keep away from heat and open flame. Absorbed through the skin—may be fatal if swallowed.

[See Antidote Statement]

DANGER! KEEP OUT OF REACH OF CHILDREN.

SOYBEANS — DYANAP (H.W.) OVERTOP APPLICATION

STATE LABEL FOR OKLAHOMA ONLY

(The following section follows AERIAL APPLICATION up to the crook stage on soybeans.)

SOYBEANS— DYANAP (H.W.) OVERTOP APPLICATION

An application of **DYANAP** (H.W.) by ground or air over the tops of soybeans after the second trifoliate leaf stage of growth may also be made. For best results spray when the weeds are small and succulent. In the southern soybean areas with determinate type soybeans **DYANAP** (H.W.) should not be applied after flowering begins. In the midwest with indeterminate type soybeans **DYANAP** (H.W.) should not be applied after the soybeans reach twenty inches in height.

DIRECTIONS FOR USE

A. **GROUND APPLICATION** — Apply 2 to 4 quarts of **DYANAP** (H.W.) in 8 to 10 gallons of water per acre (broadcast rate). The spray boom should be positioned 3 feet above the tops of the soybean plants and be equipped with hollow cone or flat fan-type nozzles. Maintain a high spray pressure (i.e. 60 to 70 psi) during application. This insures a fine spray mist for better overall coverage. Use 2 to 3 quarts of **DYANAP** (H.W.) on broadleaf weeds

up to 3 inches tall, such as jimsonweed, morningglory, pigweed and smartweed. This application will control velvetleaf up to one inch tall. If the majority of the weeds are 3 to 6 inches tall, 4 quarts of **DYANAP** (H.W.) should be used. **DYANAP** (H.W.) is very effective for the control of cocklebur from the time it emerges until it is 6 inches tall. **DYANAP** (H.W.) will suppress the growth of cocklebur and morningglory taller than 6 inches, reducing their competition with the soybean crop.

B. **AERIAL APPLICATION** — Apply 2 to 4 quarts of **DYANAP** (H.W.) in a minimum of 5 gallons of water per acre and follow instructions given for ground application above. Fly plane 10 to 15 feet above the tops of the soybean plants (DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREA TREATED)

NOTE — Applications should not be made to wet soybean foliage. Avoid drift onto other crops. Do not use surfactants with **DYANAP** (H.W.) in overtop applications. Delay spray application if rain is expected within 6 hours of spraying.

THIS LABEL IS IN ADDITION TO THE
STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER.