

**HILL FLY BAIT . . .**  
**Attracts More Flies!**  
**Kills More Flies!**

**DIRECTIONS**


**As a Dry Bait:** Sprinkle HILL FLY BAIT on floors, window sills, walks and other areas where flies gather. Use 4 ounces per 1,000 square feet. **IMPORTANT**—a faster kill may be obtained if applied to a moistened surface. Repeat application as necessary. Apply liberally as a dry bait to manure piles, garbage dumps and other fly breeding places.

**As a Liquid Bait:** Dissolve one (1) pound HILL FLY BAIT in one-half gallon of water. This solution can then be sprayed (as a coarse, wet spray) on walls, painted on sills and surfaces where flies gather. Where suitable surfaces are not available, treat burlap sacks or strips of cardboard, placed where flies gather.

**THE NEW FLY KILLER  
 THAT FLIES FEAST ON AND  
 DIE IN SECONDS**

Recommended for use in: Dairy Barns, Stock Yards, Stables, Loading Docks, Drive-Ins, Poultry Houses, Rendering Plants, and Hog Farms.

**Not Recommended For Use in Homes**



**ACTIVE INGREDIENTS**

\*(2,2-dichlorovinyl dimethyl phosphate) . . . . . 0.5%

**INERT INGREDIENTS** . . . . . 99.5%

**100.0%**

\*(2,2-dichlorovinyl dimethyl phosphate) 0.46% and related compound, 0.04%

Formulated with 0.8% DDVP, but declared lower to allow for possible loss during storage.

**KILLS-RESISTANT AND NON-RESISTANT FLIES  
 FLIES ARE ATTRACTED TO FLY BAIT—FEAST ON IT  
 AND DIE IN SECONDS.**

**ECONOMICAL TO USE**

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE BACK PANEL FOR ADDITIONAL CAUTION.  
 DO NOT CONTAMINATE FOOD OR FEED.  
 USDA REG. NO. 402-73

**CAUTION:**

**Keep Out of Reach of Children.**

**Avoid Breathing Spray Mist.**

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after using. Do not scatter by hand—use a container of some kind. Avoid contamination of food and animal feeds. Do not apply to animals. Do not use in human dwellings. Keep out of reach of children, pets and poultry. Do not treat portions of buildings where animals lick. Do not use treated garbage as animal feed. Do not use where milk is stored or processed. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

If poisoning occurs GET PROMPT MEDICAL ATTENTION. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

**Net Weight One Pound**

**HILL MANUFACTURING CO. ATLANTA, GA.**