

Copy

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

SWIMMING POOL WATER DISINFECTION - For a new pool or spring start-up, superchlorinate with 77 to 155 ounces of Super Chlor 84 for each 10,000 gallons of water to yield 5 to 10 ppm free-chlorine by weight. Check the level of available chlorine with a test kit. Adjust and maintain pool water pH to between 7.2 to 7.6. Adjust and maintain the

stability of the pool to between 80 to 100 ppm. To maintain the pool, add manually or by a feeder device 10 ounces of Super Chlor 84 for each 10,000 gal. of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Do not reenter pool until the chlorine residual is between 1.0 to 3.0 ppm.

Every 7 days or as necessary, superchlorinate the pool with 77 to 155 oz. of Super Chlor 84 for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Do not reenter pool until the chlorine residual is between 1.0 to 3.0 ppm.

SPAS / HOT TUBS - Apply 7.5 ounces of Super Chlor 84 per 1000 gallons of water to obtain a free available chlorine concentration of 5 ppm, as determined by a suitable chlorine test kit. Adjust and maintain pool water pH to between 7.2 and 7.6. Some oils, lotions, fragrances, cleaners, etc. may cause foaming or cloudy water as well as reduce the efficiency of the product.

To maintain water, apply 7.5 ounces of product per 1000 gallons of water over the surface to maintain a chlorine concentration of 5 ppm.

SANITATION OF NON-POROUS FOOD CONTACT SURFACES

RINSE METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that available chlorine does not drop below 50 ppm.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not seek equipment overnight.

SANITIZATION OF POROUS FOOD CONTACT SURFACES

RINSE METHOD - Prepare a 800 ppm solution by thoroughly mixing 9 oz. of Super Chlor 84 with 10 gallons of water. Chlor surfaces in the normal manner. Rinse all surfaces thoroughly with the 800 ppm solution, maintaining contact at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 2 oz. of Super Chlor 84 with 10 gallons of water. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Do not rinse and do not seek equipment overnight.

with COMMENTS

ACCEPTED

JUL 29 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

402-78

HILL #84

SUPER CHLOR 84

ACTIVE INGREDIENT: Sodium Hypochlorite 8.4%
INERT INGREDIENTS 51.6%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

STATEMENT OF PRACTICAL TREATMENT

If contact with eyes occur - Flush with water for at least 15 min. Get prompt medical attention.

If contact with skin occur - Wash with plenty of soap and water. If ingested - Drink large quantities of milk or gelatin solution, if these are not available, drink large quantities of water. Do not give vinegar or other acids. DO NOT INDUCE VOMITING. Get prompt medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER: Corrosive. May cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear safety glasses or goggles and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not discharge into lakes, streams, ponds or public waterways unless in accordance with a NPDES permit. For guidance the regional office of the U.S. Environmental Protection Agency.

PHYSICAL OR CHEMICAL HAZARDS
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release chlorine gas which is irritating to eyes, lungs and mucous membranes.

STORAGE AND DISPOSAL

Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water. Product or residues that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not reuse container but place in trash collection. Do not contaminate food or feed by storage, disposal or cleaning of equipment.



HILL
MANUFACTURING COMPANY, INC.
1500 JONESBORO ROAD SE ATLANTA, GA 30313 404-522-8364

SANITIZATION OF NON-POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 9 oz. of Super Chlor 84 with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not seek equipment overnight.

DISINFECTION OF NON-POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 9 oz. of Super Chlor 84 with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 10 minutes. Do not rinse equipment with water after treatment and do not seek equipment overnight.

SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 9 ounces of Super Chlor 84 with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not seek equipment overnight.

SEWAGE AND WASTEWATER TREATMENT

EFFLUENT SLIME CONTROL - Apply a 100 to 1000 ppm available chlorine solution at a location which will allow complete mixing. Prepare this solution by mixing 15 to 150 ounces of Super Chlor 84 with 100 gallons of water. Once control is evident, apply a 15 ppm available chlorine solution. Prepare this solution by mixing 4.5 ounces of Super Chlor 84 with 100 gallons of water.

FILTER BEDS SLIME CONTROL - Remove filter from service, drain to a depth of 1 foot above filter sand, and add 50 ounces of Super Chlor 84 per 20 square feet evenly over the surface. Wait 30 minutes before draining water to a level that is even with the top of the filter. Wait for 4 to 6 hours before completely draining and backwashing filter.

MILK AND PAPER MILK PROCESS WATER SYSTEMS

SLUG FEED METHOD - Initial Dose: When system is normally loaded, apply 77 to 155 ounces of Super Chlor 84 per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

AGRICULTURAL USES

FOOD EGG SANITIZATION - Thoroughly clean all eggs. Thoroughly mix 9 ounces of Super Chlor 84 with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before eating or brooding. Do not apply a potable water rinse. The sanitizer should not be re-used to sanitize eggs.

Net Contents _____ gallons
Recommended for Commercial / Industrial use only.
EPA Reg. No. 408-78 EPA Est. 462-GA-1

7/8