

Center Panel

1143)

HILCO #74
ROACH & PEST SPRAY

Contains [®] Baygon

For Industrial and Institutional Use

Residual Spray is a highly effective ready-to-use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and long residual activity.

ACTIVE INGREDIENTS:

o-Isopropoxyphenyl methylcarbamate *	1.00 %
2,2-Dichlorovinyl dimethyl phosphate	0.23 %
Related Compounds	0.02 %
Pyrethrins	0.05 %
Piperonyl Butoxide, Technical **	0.10 %
N-Octyl Bicycloheptene dicarboximide	0.166 %
Petroleum Distillate	83.95 %

INERT INGREDIENTS:

14.484%

* U. S. Patent No. 3,111,539

This material used under license from Chemagro Corp.

[®] Baygon is the registered trademark of Farbenfabriken Bayer A.G. Chemagro Corp. licensee.

** Consists of .08% (butylcarbityl) (6-propylpiperonyl) ether and .02% related compounds.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Millipedes, Mosquitoes, Paperlice, Punkies, Sandflies, Scorpions, Silverfish, Sowbugs and Spiders. Exposed stages of Cadelles, Cigarette Beetles, Dermestids, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

NET CONTENTS _____ GALLONS

USDA REG. NO. _____

HILL MANUFACTURING CO.
1500 Jonesboro Rd., S.E.
Atlanta, Ga. 30315

ACCEPTED

402-975

SIDE PANEL

2/0/3)

DIRECTIONS FOR USE

GENERAL:

HILCO #74 ROACH & PEST SPRAY is especially formulated for use in Flour Mills, Dairies, Canneries, Bottling Plants, Warehouses, Super Markets, Restaurants and Schools.

Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Treat exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches - Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

Ants - Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

Caddises, Cigarette Beetles, Dermestids, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils (exposed stages) - Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

Silverfish, Firebrats and Paperlice - Apply to localized areas infested by these insects.

Brown Dog Ticks - Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

Clover Mites, Crickets, Earwigs, Millipedes, Centipedes, Sowbugs, Scorpions and Spiders - Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention to window sills.

Fleas, Flies, Gnats, Mosquitoes, Punkies and Sandflies (outdoors only) - Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

For Use in Meat Packing and Food Processing Plants: Use only in the inedible product areas of these plants, including those that are federally inspected. Any exposed food products, equipment, or packaging materials must be removed or covered before spraying. Treat Food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants. Avoid breathing spray mist.

(continued)

2 (of 3)
CAUTION

May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Should illness occur, get prompt medical aid.
To physician - Atropine sulfate is antidotal.