## DIRECTIONS FOR USE

This disinfectant has certain advantages in that it is relatively non-corrosive and does not injure cloth or metals. HILL LEMON-CIDE is not affected appreciably by organic matter and has a pleasant odor.

FOR GENERAL DISINFECTING: Remove all filth and dirt from the premises and use a solution of one part disinfectant to 30 parts of water. This mixture acts as a disinfectant against S'. typhosa.

In the typhoid sick room, in the disinfecting of bed clothes, beds, and utensils, disinfectant should be used in the proportion of 4 ounces to each gallon of water. Use one-half cup of the disinfactant to two gallons of water for mopping, scrubbing, or cleaning purposes.

GARBAGE: To deodorize, sprinkle with solution one-half cup to a gallon of water. Wash the empty can with some of the solution and leave the wash in can.

FOR THE SICK ROOM: All utensils, such as bedpans, urinals, etc., wash with a solution of three ounces of disinfectant to  $\frac{1}{2}$  gallon of water.

WATER CLOSETS AND URINALS: Use four ounces of disinfectant to a gallon of water. Apply freely to all surfaces to be disinfected and scrub the exposed areas.

## No. 159 LEMON-CIDE

## DISINFECTANT ---- GERMICIDE ---- BACTERICIDE FUNGICIDE — DEODORANT **PHENOL COEFFICIENT 5** Against Salmonella Typhosa and **Staphyloccus Aureus.**

ACTIVE INGREDIENTS Isopropanol, Dodecyl Benzene Sodium Sulfonate, o-benzyl p-chlorophenol, Orthophenylphenol, **P-Tert.-Amylphenol**, Essential Oils INERT INGRREDIENTS (Water)

**CAUTION:** KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

HILL MANUFACTURING CO., ATLANTA, GA.

34.5%

61.5%

FOR ANIMAL QUARTERS: Before applying the disinfectant, remove the gross filth and heavy soil. Keep the barns, kennels, pens and sties thoroughly clean by scrubbing and spraying freely with a solution of disinfectant, in the proportion of 4 oz. to a gallon of water. Scrub all floorings with the same solution.

MATCHERIES: Wash floors and trays with a solution of one part disinfectant to 30 parts of water and repeat as often as necessary. In all contagious diseases, separate the healthy fowls from the affected ones, giving them fresh, clean quarters.

FOR USE AS A FUNGICIDAL AGENT AGAINST ATHLETE'S FOOT FUNGI ON ENVIRONMENTAL SURFACES: As a fungicidal agent against Athlete's Foot Fungi in locker rooms, swimming pool surfaces, dilute by dissolving four ounces of disinfectant in one gallon of water. Scrub walkways, shower floors, foot bath surfaces, rubber mats, and other surfaces contacted by the feet periodcally.

CAUTION: Harmful if swallowed. Avoid getting in eyes or on skin. Solutions of this disinfectant may cause skin irritation in some persons who might be allergic to this type of compound. Such persons who might be allergic to this compound should avoid getting solution of this disinfectant on the skin, and if spilled thereon, remove by washing with water.

Do not reuse empty drum or container. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DU NOT CUNTAMINATE FOOD OR FEED FOR INDUSTRIAL USE ONLY EPA REG. NO. 402-113

> NET CONTENTS GAL.

ACCEPTED C [ ] ] 1971 The second second second second second F S & MARLEN AND BASE TO THE ALT FOR TO MOVIC PUISON RED STER. ED UND & NU. 402-