

USE DIRECTIONS:

Use at 2 ounces per gallon. This product will be effective against *Pseudomonas aeruginosa* PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use SAN-O-15 with a cloth or mop and apply to walls, floors and other hard surfaces.

If used for sanitization of food equipment or food contact items, limit the level of active quaternary to 200 ppm (2.8 ounces per 5 gallons).

NO RINSE REQUIRED.

For heavy soil or organic matter, a precleaning step is recommended.

PHENOL COEFFICIENTS:

Guaranteed	Minimum
S. aureus	42
S. typhosa	27

Rinse empty container thoroughly with water and discard.

ACCEPTED
2-8-73
402-116
FOR UNDER NO. TO ATTACHED COPIES



SAN-O-15

CLEANER

DISINFECTANT

DEODORIZER

FUNGICIDE

Designed for Hospital and Institutional use.

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	2.25%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	2.25%
Sodium Carbonate	3.00%
Tetrasodium ethylenediamine tetraacetate	1.00%
INERT INGREDIENTS:	91.50%
	100.00%

NET CONTENTS _____ GAL. EPA REG. NO. 402-116

WARNING KEEP OUT OF REACH OF CHILDREN

See right side panel for additional precautionary statements.

HILL MANUFACTURING CO. - ATLANTA, GA.

SAN-O-15 is a cleaner, disinfectant and sanitizer, designed for institutional use such as Hospitals, Nursing Homes and Institutions where housekeeping will be of prime importance in controlling the hazard of cross-infection on environmental surfaces. SAN-O-15 can provide the assurance you need in this area, when used as directed. For use on floors, walls, sink tops, garbage pails, telephones, rest rooms, bed pans and operating rooms. Effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. Also effective against pathogenic fungi, *Trichophyton interdigitale*. When used on hard inanimate surfaces at 2 ounces per gallon of water is effective against (1) Adenovirus type 1, (2) Herpes Simplex virus, (3) Influenza, A2 virus and (4) Vaccinia virus.

When used for sanitization of food processing equipment and food contact items, limit active quaternary to 200 ppm. At this level, No Potable Water Rinse is required.

WARNING

Keep Out of Reach of Children. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.