

Then adjust sprayer to deliver aerosol size particles. Apply as a space spray at the rate of 3 to 6 fl. oz. per 1,000 cu. ft. of room space. Repeat dosage as necessary for control. Leave the room closed for 1 hour. Do not remain in treated area. Ventilate the area thoroughly before re-entry.

FOR WASPS AND SPIDERS: Spray insects directly and apply heavy spray to nests and webs. The most effective wasp control is obtained by spraying after dark when all insects have returned to the nest.

FOR CONTROL OF CARPET BEETLES: Clean all rugs, carpets and fabrics to be treated. Spray both surfaces of carpets and floor space beneath evenly with a fine mist spray. Particular attention should be paid to cracks and crevices around the baseboards.

ANIMAL SPRAY FOR HORSES AND DAIRY AND BEEF CATTLE: For protection against Stableflies, Hornflies and Houseflies attacking animals, spray the animal against the grain of the hair, directing a fine mist around the base of the horns, on the legs and along the sides and belly and over the back. Do not spray udders. Apply just enough to wet the ends of the hair but not enough to wet the hide. One or two ounces of spray is sufficient to treat an adult cow. Apply once daily or as necessary.

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting. Wash face, hands and arms with soap and water after handling and before eating or smoking. Do not remain in treated area. Ventilate the area thoroughly before re-entry. Do not apply directly on very young animals. Do not use in poultry houses. Remove pets, birds and cover fish aquariums before spraying. Do not apply in a food processing plant while plant is in operation. When applied in a food processing plant, remove or protect food during treatment. After treatment, wash all equipment, benching, shelving, etc. with potable water where exposed food will be handled.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate any body of water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty containers. Destroy by perforating or crushing and burying in a safe place away from water supplies. Return drums to reconditioners.

PHYSICAL HAZARD

KEEP CONTAINER CLOSED. DO NOT USE. POUR SPILL OR STORE NEAR HEAT OR OPEN FLAME.


BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH THE DIRECTIONS GIVEN HEREWITH.



B. G.



SINCE 1904



PYRETHRUM FOGGING SPRAY

Containing

PYRETHRUM WITH ACTIVATOR FOR HOME, FARM AND INDUSTRY

A contact spray for control of listed flying insects in home, restaurants, hotels, food storage warehouses, dairies, meat packing plants and similar establishments with electric mist spraying or fogging equipment. Also outdoor adult mosquito control.

ACTIVE INGREDIENTS	BY WEIGHT
Pyrethrins	0.10%
Piperonyl Butoxide, Technical*	0.75%
Isopar Hydrocarbons	99.15%
TOTAL	100.00%

*Consists of 96.9% (butylcarbitol) (6-propylpiperonyl) ether and 0.15% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

See side panel for additional cautions.

B. G. PRATT DIVISION
GABRIEL CHEMICALS LTD.
204 21st Avenue
Paterson, N. J. 07509

22

NET CONTENTS: 1 5 30 55 GALLONS

ACCEPTED
JUL 11 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 704-162

EPA Reg. No. 904 162
EPA Est 984 NJ 1
Prod No 3400

TO REDUC
TOES, GNA
swimming p
playground
penetrate
mentals an
theaters. T
patio, por
peated ap
heavy. Ap
1,000 sq. f
Do not spr
only the t
Direct spra

FOR CONT
SMALL FL
buildings
ventilators
birds and
fine mist c
be affected
toward the
area is fil
10 to 15 m
area thoro
insects.

CONTACT
COARSE
ROACHES
SECTS: G
FLOUR B
MOTH LA
AND SAW
posed foc
the base
cabinets,
trucks us
in small
ment cap
equipment
spaces in
houses.
de-energ
for oil to
in "on"
remain in
re-entry,
treated b
intended f

CONTACT
SILVERFI
SITUATIC
spray Ap
woodwork
in all pla
insects as

B. G.

PRATT

SINCE 1904

PYRETHRUM FOGGING SPRAY

Containing

PYRETHRUM WITH ACTIVATOR FOR HOME, FARM AND INDUSTRY

A contact spray for control of listed flying insects in home, restaurants, hotels, food storage warehouses, dairies, meat packing plants and similar establishments with electric mist spraying or fogging equipment. Also outdoor adult mosquito control.

ACTIVE INGREDIENTS	BY WEIGHT
Pyrethrins	0.10%
Piperonyl Butoxide, Technical	0.75%
Isoparaffinic Hydrocarbons	99.15%
TOTAL	100.00%

*Consists of 0.60% (butylcarbityl) (6-propylpiperonyl) ether and 0.15% related compounds.

CAUTION

ACCEPTED KEEP OUT OF REACH OF CHILDREN.

See side panel for additional cautions.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 904-162

B. G. PRATT DIVISION
GABRIEL CHEMICALS LTD.

204 21st Avenue
Paterson, N. J. 07505

EPA Reg. No. 904-162
EPA EST. 984 NA-1
Prod. No. 5400

NET CONTENTS: 1 5 30 55 GALLONS

DIRECTIONS

OUTDOOR SPRAY

TO REDUCE THE ANNOYANCE OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS around patios, swimming pools, home yards, and outdoor areas in parks and playgrounds: When the air is still, spray as a fine mist or fog to penetrate hiding places such as grasses, shrubs and ornamentals around homes, picnic areas, stadiums and open air theaters. Treat as large an area as possible surrounding the patio pool or grounds where insect protection is desired. Repeated applications may be necessary when infestation is heavy. Apply at the rate of 1.5 gallons per acre (4.4 fl. oz. per 1,000 sq. ft.).

Do not spray flowers and ornamentals at close range. Allow only the fine mist or "dry" fog to contact desirable plants. Direct spray away from the swimming pool.

INDOOR SPRAY

FOR CONTROL OF FLIES, ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS in homes, food establishments, farm buildings and warehouses: Close all windows and stop all ventilators. Pilot lights should be extinguished. Remove pets, birds and cover fish aquariums before spraying. Allow only fine mist or "dry" fog to contact fabric or surfaces that might be affected by the oil base. Direct fine mist or fog upward toward the center of the room and around the sides until the area is filled with fog or mist particles. Keep area closed for 10 to 15 minutes. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Clean up and destroy fallen insects.

CONTACT SPRAY — FOR USE WITH SPRAYERS THAT GIVE COARSE WETTING SPRAY PARTICLES TO KILL COCKROACHES AND ACCESSIBLE STAGES OF GRANARY INSECTS: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RED FLOUR BEETLES, INDIAN MEAL MOTH LARVAE AND ADULTS, CADELLES, GRAIN MITES, AND SAW-TOOTHED GRAIN BEETLES: Cover or remove exposed food. Spray outside walls of grain bins, floorings, about the base of milling machinery, walls, baseboards, areas behind cabinets, around pipes and inside surfaces of boxcars and trucks using 1 gallon of spray per 1,000 sq. ft. of space. Apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream in points between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, conduits, motor housings, and junction or switch boxes. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in "on" position. Repeat applications as necessary. Do not remain in treated areas. Ventilate the area thoroughly before re-entry. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

CONTACT AND SPACE SPRAY — TO KILL COCKROACHES, SILVERFISH EARWIGS, ANTS AND SPIDERS IN SHELTERED SITUATIONS. Adjust spraying device to deliver a coarse wet spray. Apply as a contact spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food

904-162