Rats And Mice

vith PROLIN

improved Warfarin product

5 POUNDS





FOR REQUIRED PRECAUTIONS


yl)-4-hydroxycoumarin _____ 0.025% (Sulfaquinoxaline) _____ 0.025% 99.950% OUT OF REACH OF CHILDREN EPA No. 404-118

DIRECTIONS

RATS: Place generous amounts (minimum 4 ounces) of RAT-O-MICE in places where rats feed or frequent. Protect bait from children, dogs, cats, livestock and wildlife by means of a bait box, cage or suitable covering. (See Illustration.)







Watch bait stations closely and refill with RAT-O-MICE before bait is entirely consumed. Baiting should be continued as long as evidence of feeding is observed. (at least 10 days).

Where a continuous source of infestation is present from nearby dumps or fields, a supply of RAT-O-MICE should be maintained in permanent bait stations at all times.

MICE: Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8 to 12 foot intervals. Bait should be placed in locations not accessible to children, pets, domestic animals and wildlife or in tamper-proof bait boxes. An uninterrupted supply of bait should be maintained (generally up to fifteen days).

NOTE: Bury or destroy rodents found on premises. Bait which has deteriorated due to becoming sour or moldy should be promptly replaced.

MANUFACTURED BY



ACCEPTED

JUN 22 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER. ED UNDER NO. 404-18 SUBJECT TO ATTACHED COMMENTS.