## McConnon INSECTICIDAL DUST

## **ACTIVE INGREDIENTS:**

Rotenone	DEN
Other Cube' Extractives	75%
INERT INGREDIENTS.	. 1.50%
INERT INGREDIENTS:	97.75%
Total	100 000

EPA Reg. No. 404-325 EPA Est. 404-MN-1

CAUTION: KEEP OUT OF REACH OF CHILDREN.

ACCEPTED

Manufactured by

DADER 144 POUNDS

FOR ECONOMIC POISON REGISSION

WEIGHT 4 POUNDS

LOW SUBJECT

WINDER 14 POUNDS

## **DIRECTIONS FOR USE**

WHEN TO APPLY—Whenever possible, apply dust at a time when plants are wet with dew, or after a rain. This helps the dust to adhere and give a better control. Repeat application when eggs hatch or insects reappear. Discontinue use within one day of harvest.

HOW TO APPLY—Make a thorough application, directing the dust against both the upper and lower leaf surfaces.

PLANT USES — McConnon Insecticidal Dust may be used in the control of the following:

IMPORTED CABBAGE WORM ON CABBAGES CABBAGE LOOPER ON CABBAGES THRIPS ON CUCUMBERS STRIPED CUCUMBER BEETLE ON CUCUMBERS (Apply at the rate of I pound per 1,000 sq. ft. Repeat if needed.) FLEA BEETLES ON BROCCOLI, CAULIFLOWER, SWISS CHARD, TURNIPS VEGETABLE WEEVILS ON BROCCOLI, CABBAGE, CARROTS, CAULIFLOWERS, COLLARD, KALE, TURNIPS GREENHOUSE WHITE FLY ON GREENHOUSE LETTUCE -CHERRY FRUIT FLIES ON CHERRIES MEXICAN BEAN BEETLE ON BEANS THRIPS ON TOMATOES VEGETABLE WEEVILS ON MUSTARD GREENS ASPARAGUS BEETLE ON ASPARAGUS GREENHOUSE WHITE FLY ON GREENHOUSE TOMATOES PEA APHIDS ON PEAS JAPANESE BEETLES ON APPLES TOBACCO FLEA BEETLES ON TOMATOES, POTATOES

CAUTION: Avoid inhalation of dust.

"Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes."

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

