

McConnon Potato and Tomato Dust is a convenient, ready-to-use garden dust. When used as directed, it is an effective combination of insecticide and fungicide to control the insects and diseases on the crops listed below.

**CROPS PESTS AND DISEASES CONTROLLED**

**POTATO  
TOMATO**

PESTS: Colorado potato beetle, leafhoppers, flea beetles, lace bugs, tomato fruitworm, tomato hornworm, stink bugs and tarnished plant bugs.  
DISEASES: Early and late blight, Anthracnose, and grey leaf mold.

McConnon Potato and Tomato Dust may be applied up to 5 days of harvest.

**SWEET CORN**

PESTS: Corn rootworm adults, fall armyworm, flea beetles, Japanese beetles and leafhoppers. Do not feed treated forage to livestock.

McConnon Potato and Tomato Dust may be applied up to 7 days before harvest.

**CARROTS**

PESTS: Fleabeetles, leafhoppers, tarnished plant bugs and stink bugs.  
DISEASES: Alternaria leaf spot and Cercospora leaf spot.  
Do not use tops for food or feed purposes.

McConnon Potato and Tomato Dust may be applied up to 7 days before harvest.

**CUCUMBERS  
SUMMER SQUASH  
MELONS**

PESTS: Cucumber beetles, squashbug, melonworm, pickleworm, flea beetles and leafhoppers.  
DISEASES: Scab, Anthracnose, downy mildew, Alternaria leaf spot, Cercospora leaf spot and gummy stem blight.

McConnon Potato and Tomato Dust may be applied up to 5 days of harvest.

**DIRECTIONS FOR USE:** McConnon's Potato and Tomato Dust may be used in conventional hand duster or sprinkled from small holes made in the can. Before applying please read label carefully and use strictly in accordance with label cautions, warnings and directions. Apply McConnon Potato and

Tomato Dust as soon as foliage appears. McConnon's Potato and Tomato Dust should be applied to all exposed parts of the plant including the lower leaf surface. To control the insects and diseases listed on the chart that can affect these crops, repeat dusting at 5 to 10 day intervals.

**BEE CAUTION: MAY KILL HONEY BEES IN SUBSTANTIAL NUMBERS**

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond the flight range until 1 week after application or to take other equally effective precautions.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions herewith.

**NOTE:** This container is sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping.

ACCEPTED  
MAY 24 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 404-326

# Mc Connon

## POTATO and TOMATO DUST

**INSECTICIDE - FUNGICIDE**  
**Contains "Carbaryl"**

For Control of Garden Insects and Diseases listed on the back panel

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate) .....	5.0%
A Coordination product of zinc ion and manganese ethylene bisdithiocarbamate .....	4.8%
(Metallic manganese content .96%, metallic zinc content .12%)	
<b>INERT INGREDIENTS: .....</b>	<b>90.2%</b>
<b>TOTAL</b>	<b>100.0%</b>

EPA Registration No. 404-326

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS HARMFUL IF INHALED OR SWALLOWED.**


Do not breathe dust. Do not take internally. Avoid contact with skin. Skin contact may be harmful. In case of skin contact, wash with water; for eyes, wash thoroughly with water and get medical attention. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS. Contamination of fish-bearing waters with this product may harm fish. Food utensils such as tablespoons should not

be used for food purposes after use with pesticides. DO NOT REUSE CONTAINER. DESTROY BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs. (See back panel for additional precautions) Important: Store in a cool, dry place. Do not allow to become wet or over-heated in storage. Keep container closed when not in use.

**NET WEIGHT ONE POUND**

Manufactured by  
**McCONNON & COMPANY**  
Winona, MN 55987

# McConnon and Company



**Manufacturing Chemists**

WINONA, MINNESOTA 55987

Telephone (507) 452-2911

May 14, 1973

U. S. Environmental Protection Agency  
Pesticides Office  
Registration Division  
Washington, D. C. 20250

Attention: Mr. James M. Rea  
Acting Section Head  
Insecticide-Rodenticide Branch

Subject: McCONNON POTATO and TOMATO DUST  
EPA Reg. No. 404-226  
Your letter of May 7, 1973

C 899803

404-GEA-PA-326

Dear Mr. Rea:

Attached, please find five (5) copies of the finished label for McCONNON POTATO and TOMATO DUST, incorporating the changes as suggested in your letter dated May 7, 1973 a copy of which is enclosed.

Sincerely,

McCONNON and COMPANY

*Mrs. Shirley Snider*

Mrs. Shirley Snider  
Registration Department