

[404-327]
1082



MALATHION
DRY INSECTICIDE

A POWDERED INSECTICIDE
FOR USE ON LIVESTOCK AND
POULTRY. WITH DEODORANT.

ACTIVE INGREDIENT:

Malathion*..... 5%

INERT INGREDIENTS: 95%

*O, O-dimethyl phosphorodithioate
of diethyl mercaptosuccinate

EPA Reg. No. 404-327

CAUTION: KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Avoid breathing of
dust. Avoid contact with skin. Wash thoroughly
after handling. Avoid contamination of feed and
foodstuffs, feeding troughs and water fountains.
Apply this product only as specified on the label.
Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes,
streams or ponds.

NET WEIGHT 25 POUNDS

Manufactured by

McConnon & Company

WINONA MINNESOTA

404-327

[404-327] 2 of 2

DIRECTIONS



HOGS: (Lice) Dust animals and pens thoroughly. Repeat thereafter as needed.

CATTLE (Beef and Non-Milking): (Lice) Treat animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 3 tbsps. per animal. Apply on back and neck of the animal and repeat applications at 10-14 day intervals.

Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: (Horn Fly) Use 3 tbsps. per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 5 hours before milking or during milking. Repeat applications at 10-14 day intervals.

POULTRY: (Chicken Red Mite, Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary. Dust bath box: (Northern Fowl Mite, Poultry Lice) Use 1 pound dust per box (18 x 12 x 3") per 30 hens. Remove boxes when dust has been used. Dust bath boxes or natural soil wallows 15 pounds per box (2" x 3' x 1') per 100 birds or apply in natural soil wallows at 7½ pounds per 50 birds. Treatment of poultry for Northern Fowl Mite and Chicken Red Mite should be a supplemental treatment to the treatment of the premises. Premise Treatment - Apply 1 lb. per 50 to 60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas.

DOGS AND CATS: (Fleas) Direct application: Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, puff duster, or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid necessary applications, animals should be closely examined for fleas between treatments. If the animal is confined to an area free of fleas, one treatment may be sufficient.

Do not treat animals under one month of age.
When applying dusts avoid contamination of feed,
food containers and watering troughs.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.