

DIRECTIONS FOR USE

To mix this material, slowly sift the desired amount into the partially filled tank of water with the agitator running and when thoroughly mixed apply at once with continuous agitation. Where desired this material may first be made to a thick paste by adding a little water. This should then be diluted and added to the tank while it is filling. Where this material is used with Lead Arsenate, no Lime need be used as a safener, since this material is itself a good safener for Lead Arsenate.

APPLES: For early-season control of scab, cedar rust and blotch, use 1 to 1½ pounds of this material, per 100 gallons of water in pre-bloom, calyx and early cover sprays. For rust control make an additional bloom application using ½ pound per 100 gallons of water. For late-season control of scab, black rot, bitter rot, sooty blotch, fly speck and Brooks' spot use ¾ pound per 100 gallons of water. Where bitter rot and Brooks' spot are severe problems increase dosage to 1½ to 2 pounds per 100 gallons of water. Do not apply within 7 days of harvest.

On Golden Delicious, Jonathan, Baldwin and other varieties of apples which are especially apt to russet some injury may result from the use of this type of material if used in the early season or bloom applications. The best time for use is in the cover sprays.

When used in combination with sulfur, do not apply within 10 days to 2 weeks of an oil spray or just prior to or during periods of excessively high temperatures.

PEARS: For control of scab and sooty blotch, use 1½ pounds per 100 gallons of water in pink, calyx, first and second cover sprays, and 1 pound in summer sprays. Do not apply within 7 days of harvest.

GRAPES: For control of black rot, use 2 pounds per 100 gallons of water, applying just before bloom, just after bloom and two weeks later. Do not apply within 7 days of harvest.

CHERRIES: For control of brown rot and leaf spot, use 1½ pounds per 100 gallons of water in petal-fall and cover sprays.

PEACHES: For leaf curl, use 1½ pounds per 100 gallons of water. Apply in the fall after leaves drop or in the spring before buds begin to swell. Do not apply within 21 days of harvest.

CABBAGE: For control of downy mildew in plant beds, use 1½ pounds per 100 gallons of water, making 2 to 3 applications each week.

TOMATOES: Gray Mold (Botrytis) - Use 3 pounds per 100 gallons of water. Spray for complete coverage. Make seed bed application just before seedlings break the soil surface and repeat at 7 day intervals. Make field application when disease threatens and repeat at 7 to 10 day intervals as necessary to maintain control.

TOBACCO: For control of blue mold in plant beds, use 2 to 3 pounds per 100 gallons of water. Apply at the rate of 3 gallons per 100 square yards when plants are small, increasing to 6 gallons when plants are ready for transplanting. Begin applications when plants are the size of a dime, or when blue mold is reported in the area, and repeat twice weekly until plants are transplanted.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 407-318

L-9811A-C

Imperial

FERBAM FUNGICIDE

ACTIVE INGREDIENT:

Ferbam (Ferric Dimethyl dithiocarbamate) 76.0%

INERT INGREDIENTS: 24.0%

Total 100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation of eyes, nose, throat and skin! Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water, for eyes, get medical attention.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

Keep packages or Dust Away From Sparks, Flames or Hot Surfaces.

(DO NOT USE IN UNDILUTED FORM)

NET WEIGHT 1 POUND

Manufactured by

IMPERIAL INC.

SHENANDOAH, IOWA 51601

