

ACTIVE INGREDIENTS	
Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid)*	6.82%
Dimethylamine Salts of related acids	1.20%
Dimethylamine Salt of 2,4- dichlorophenoxyacetic acid**	13.34%
INERT INGREDIENTS	78.64%
Total	100.00%

*Equivalent to 5.64% 3,6-dichloro-o-anisic acid (dicamba) or .5 pounds per gallon.

**Equivalent to 11.09% 2,4-dichlorophenoxyacetic acid or .1 pound per gallon.

DIRECTIONS

Time of Application: Best results are obtained when Lawn Weed Killer is used on weeds that are young and in a rapid growing condition.

Lawn and ornamental turf use: Dilute with water and apply as a spray. For best results, apply when air temperatures are 50° to 85°F. To minimize drift, use coarse spray. One gallon of mixed spray should cover about 250 square feet.

For early season use against broadleaf plantain, buckhorn plantain and dandelions, mix 1 tablespoon Lawn Weed Killer per gallon of water. Apply to thoroughly wet the weeds.

For late season use against mature broadleaf plantain, buckhorn plantain, dandelions, and for control of wild onion, garlic, henbit, chickweed, white clover, poison ivy and poison oak, mix 1½ tablespoons Lawn Weed Killer per gallon of water. Apply to thoroughly wet all plant foliage. On wild onion and garlic, repeat treatment Spring and Fall.

Imperial

LAWN WEED KILLER

(Contains BANVEL D and 2,4-D)

CAUTION

KEEP OUT OF THE REACH
OF CHILDREN

See Back Panel for Additional Cautions

NET CONTENTS 1 PINT

Manufactured by

IMPERIAL INC.

SHENANDOAN, IOWA 51601

25



CAUTION

Avoid inhalation of spray mist. Avoid skin, eyes and clothing. In case of contact with water; for eyes, get medical attention immediately. Keep out of the reach of children.


IMPORTANT

Keep out of any body of water. Avoid coarse sprays are less likely to drift onto desirable crop plants. Do not apply near line of small trees or shrubs. Do not use in domestic or irrigation waters. Do not use to clean sprayer near desirable trees where it may contaminate water. Do not use near fertilizer, seeds, insecticides or other chemicals. Do not use on St. Augustine grass or other lawns or turf where desirable bentgrass are present. Do not use on newly seeded turf. When container is empty immediately and thoroughly clean. Never re-use. Use proper method for applying 2,4-D combinations.

WARRANTY

Seller warrants that this product conforms to the chemical description and is reasonably safe under the conditions of use stated on the label when used in accordance with the directions under normal conditions of use. This warranty nor any other warranty, express or implied, extends to the use of this product under abnormal conditions, or under abnormal conditions not reasonably foreseeable, and buyer assumes the risk of any such use.

EPA Reg. No. 107-261



Imperial

LAWN WEED KILLER

(Contains BANVEL D and 2,4-D)

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See Back Panel for Additional Cautions

NET CONTENTS 1 PINT

Manufactured by

IMPERIAL INC.

SHENANDOAN, IOWA 51601

25



CAUTION

Avoid inhalation of spray mist. Avoid contact with skin, eyes and clothing. In case of contact flush with water; for eyes, get medical attention. Harmful if swallowed. Keep out of the reach of children.

IMPORTANT

Keep out of any body of water. Avoid drift of spray (coarse sprays are less likely to drift) on to vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants. Do not apply within the drip line of small trees or shrubs. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer near desirable trees or shrubs or where it may contaminate water. Do not store or use near fertilizer, seeds, insecticides or fungicides. Do not use on St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never re-use. Use spray equipment for applying 2,4-D combinations only to turf.

WARRANTY

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-261

L-9688A-C

WTS	
wt of dicamba (2-anisic acid)*	0.82%
wt of related acids	1.20%
wt of 2,4-dichlorophenoxyacetic acid**	12.34%
WTS	78.64%
	100.00%

3,6-dichloro-2-anisic acid
pounds per gallon.


2,4-dichlorophenoxyacetic acid
pound

DIRECTIONS
Best results are obtained when used on weeds that are young and in good condition.

Turf use: Dilute with water and for best results, apply when air is between 80° and 95°F. To minimize drift, use a combination of mixed spray should cover the area.

Against broadleaf plantain, buckhorn, and dandelions, mix 1 tablespoon Lawn Weed Killer per gallon of water. Apply to thoroughly wet weeds.

Against mature broadleaf plantain, dandelions, and for control of wild radish, chickweed, white clover, poison ivy, and ground ivy, mix 1½ tablespoons Lawn Weed Killer per gallon of water. Apply to thoroughly wet weeds. Repeat in late fall.



Imperial

ACTIVE INGREDIENTS	
Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid)*	6.82%
Dimethylamine Salts of related acids	1.20%
Dimethylamine Salt of 2,4- dichlorophenoxyacetic acid**	13.34%
INERT INGREDIENTS	78.64%
Total	100.00%

*Equivalent to 5.66% 3,6-dichloro-o-anisic acid (dicamba) or .5 pounds per gallon.
 **Equivalent to 11.09% 2,4-dichlorophenoxyacetic acid or 1 pound per gallon.

DIRECTIONS

Time of Application: Best results are obtained when Lawn Weed Killer is used on weeds that are young and in a rapid growing condition.

Lawn and ornamental turf use: Dilute with water and apply as a spray. For best results, apply when air temperatures are 50 to 95 F. To minimize drift, use coarse spray. One gallon of mixed spray should cover about 250 square feet.

For early season use against broadleaf plantain, buckhorn plantain and dandelions, mix 1 tablespoon Lawn Weed Killer per gallon of water. Apply to thoroughly wet the weeds.

For late season use against mature broadleaf plantain, buckhorn plantain, dandelions, and for control of wild onion, garlic, henbit, chickweed, white clover, poison ivy and poison oak, mix 1 1/2 tablespoons Lawn Weed Killer per gallon of water. Apply to thoroughly wet all plant foliage. On wild onion and garlic, repeat treatment Spring and Fall.

LAWN WEED KILLER

(Contains BANVEL)

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel for Additional Information

NET CONTENTS

Manufactured by

IMPERIAL CHEMICAL COMPANY

SHENANDOAH, VA.