

Directions for Use

GENERAL INFORMATION Important! Make sure you read the entire label before using this product. This label contains important information which must be read and understood before using this product. See **Caution** for important information concerning the use of this product.

Statement: When used as directed, will give control of many annual and perennial broadleafed weeds in corn, wheat, sorghum, grain, and roadside, pasture, and lawn areas. It is most effective when used on mature, growing weeds. Temperature best for application from 70 to 90 degrees. Do not apply if rain is expected within the hour. Use low pressure equipment (40 pounds) on crop and field. Garden type pressure sprayers work well on spot spraying. Do not let the spray particles "wind drift" to sensitive nearby plants. The amount of water required for low volume applications may vary from 1 to 2 gallons per acre. In any case, use the same amount of 2,4-D recommended per acre regardless of the amount of water used per acre.

CAUTION

CAUTION—Harmful if swallowed. Do not get in eyes, on, or in clothing. In case of contact, wash with soap and water.

DON'T OVER DOSE! Follow directions carefully. Do not let spray drift to sensitive plants such as vegetables, legumes, grapes, flowers or other 2,4-D susceptible plants. Use correct spray to reduce "wind drift." Do not spray near feathers, seeds, insects, or other birds. Local spray conditions will vary. Check local recommendations first. Use separate sprayers for insecticides and fungicides. Do not use on a large scale on cotton or sensitive plants. Do not use in high temperature conditions. Vaporize the product with more susceptible plants. Do not use in irrigation ditches or water courses for any purpose.

Do not use on St. Augustine grass, carpet grass, or any other lawn or newly seeded lawns.


Do not use on any other crop. Destroy empty containers in accordance with local, state, and federal laws.

CORN—Post-Emergence After crop is up. May be applied at several stages of growth. Use 1/2 to 1 pint per acre when corn is 10 inches and over. Do not use on corn less than 10 inches high. For post-emergence control of weeds such as barnyard, milo, and other weeds, use 1/2 to 1 pint per acre in water and wet weeds.

Early Spraying: When corn is 10 inches high, use 1/2 to 1 pint per acre in water and wet weeds. Do not use on corn less than 10 inches high. For post-emergence control of weeds such as barnyard, milo, and other weeds, use 1/2 to 1 pint per acre in water and wet weeds.

Lay-By Spraying: When corn is 10 inches high, use 1/2 to 1 pint per acre in water and wet weeds. Do not use on corn less than 10 inches high. For post-emergence control of weeds such as barnyard, milo, and other weeds, use 1/2 to 1 pint per acre in water and wet weeds.

SORGHUM—Post-Emergence Only: Employ drop nozzles. Do not use on sorghum less than 10 inches high. For post-emergence control of weeds such as barnyard, milo, and other weeds, use 1/2 to 1 pint per acre in water and wet weeds.



CAUTION

KEEP OUT OF REACH OF CHILDREN

See other cautions on side panel

ACTIVE INGREDIENTS:

Isocetyl ester of
2,4-Dichlorophenoxyacetic acid*

INERT INGREDIENTS:

Total

*Equivalent to 2,4-Dichlorophenoxyacetic acid 47.0%

72.00%

28.00%

100.00%

NET CONTENTS

Manufactured by

IMPERIAL INC.

SHENANDOAH, IOWA

Directions for Use

HARD TO KILL WEEDS IN NON-CROP LAND—For perennial weeds such as Sow Thistle and Russian knapweed (see list) in drainage ditch banks, fence rows and farm roadside, use 2 to 3 pints in 20 gallon water with 1 to 2 oz. liquid detergent and thorough wetting. Do not let spray drift. Treat in spring and again at bloom time before weeds can seed. Repeat the following year. Do not contaminate water used for domestic and irrigation purposes.

APPLICATION CHART

(Do not apply to light sandy soil)

Crop	Amount Per Acre	App. Acres 1 Gallon Will Treat
CORN	Pre-Emergence	2 to 4
	Post-Emergence	16
SMALL GRAIN	Spring Wheat	13 to 16
	Oats, Barley	
	Winter Wheat	
	Rye	
PASTURES AND TURE	1 quart	1
NON-CROP AREAS AND SPOT TREATMENT	2 to 3 pints in 20 gals. water	over all spray

SUSCEPTIBLE WEEDS

(Use lower dosage rate)

Arrowhead	Lambquarter	Sunflower
Buckhorn	Mustard	Wild Carrot
Cocklebur	Morning Glory	Wild Radish
Dandelion	Pigweed	Wild Sweet
Dock	Plantain	Potato
Knottweed	Ragweed	

WEEDS MORE DIFFICULT TO KILL

(Use higher dosage rate)

Bindweed	Horse Nettle	Hoary Cress
Buckbrush	Lambsweed	Wild Onion
Buckwheat	Milkweed	Willow
Canada Thistle	Poison Ivy	Sheep Sorrel

SMALL GRAIN—(Wheat, Barley, Rye, Oats): For weeds in WINTER WHEAT and RYE, treat only in the spring. Use 1/2 to 1 pint in sufficient water to cover one acre. Apply from the tillered to the jointing stage. In SPRING WHEAT, OATS and BARLEY, use 1/2 to 1 pint per acre. Use higher dosage for hard to kill weeds. Treat when crop is fully tillered 1 1/2 to 2 inches high up until early boot stage. Do not spray during seedling stage or boot to dough stage. Grain interplanted with legumes should not be sprayed unless injury to legumes can be tolerated. Do not spray on areas treated grain field within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.

PASTURES: For broadleafed weeds (see list) use 1 quart per acre in sufficient water to thoroughly cover. Apply early in summer when weeds are actively growing and near boot stage. Do not spray on recently seeded pastures or areas with new establishment. **Note:** Do not spray pastures for 2 to 3 weeks within 100 feet of cattle or other animals. Do not spray when grass is in boot to milk stage or after heading begins. Milk legumes, including white clover, should not be damaged by water sprays. Do not spray pastures with 2,4-D if the grass is in the seedling stage. Do not spray pastures with 2,4-D if the grass is in the seedling stage.

NOTICE

Some manufacturers of the product may have a claim for patent on the product. The purpose stated on the label when used in accordance with directions under normal conditions of use, but not to be construed as a warranty, or a representation of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. The product is sold as a general use product and is not intended for use in any other manner. The product is sold as a general use product and is not intended for use in any other manner.