

IMPERIAL SEVIN* DUST

DIRECTIONS

GENERAL DIRECTIONS

Application: SEVIN. Dust when used for insect damage to plants. Make up dust in the morning or evening when the temperature is between 60° and 80° F. Dust upper and lower surface of leaves with fine dust for complete coverage. Repeat applications at 7-10 day intervals or as necessary until control is obtained. Interval is specified below for the major insect pests.

Do not reuse empty containers. Dispose of them properly with waste or burning. Stay away from smoke or fumes.

VEGETABLE INSECT CONTROL

May, be used up to day of harvest on the following: BEANS: green beans, lima beans, snap beans, cowpeas, black-eyed Mexican bean beetle, leafhopper, cucumber beetle, Japanese beetle. CORN: corn rootworm, adults, European corn borer, fall arm worm, corn earworm. For corn borers apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 7-day intervals starting when first silks appear and continuing until silk begins to dry. Three or more applications may be required, depending on severity of infestation. Timing and good coverage are essential to effective control. Application of DDT is optional. In the case of corn, if control is not achieved, a short period will seriously reduce the yield of the crop. CARROTS: flea beetle, leafhopper, European bug. CUCUMBER, SUMMER and WINTER SQUASH, MELONS, PUMPKINS: pickleworm, melonworm, cucumber beetle, squash bug, flea beetle, leafhopper. Avoid excess of mulch around plants. Some leaf beetles, and other insects, are made when mulch is used. In the case of the present crop, DDT may be used on the following: OKRA, cowpeas, snap beans. POTATO: apply as for corn and cut at 100 lbs. TOMATO, EGGPLANT, PEPPER: Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall arm worm, lace bugs, tomato fruit worm, tobacco hornworm, tunneling plant borer, stink bugs.

Do not apply within 3 days of harvest on the following: GARDEN BEETS, TURNIPS, SALSIFY. Do not apply within 14 days of harvest of tops are to be used as food crops: BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER, KOHLRABBI. Do not apply within 14 days of harvest of roots are to be used as food crops: green beetle, harlequin bug, armyworm, clubmole, cutworm, corn earworm. PARSNIPS, RADISHES, RUTABAGAS, flea beetle, leafhopper.

Do not apply within 14 days of harvest on the following: LETTUCE, HEAD-LEAFY SPINACH, SWISS CHARD, CHINESE CABBAGE, COLE SLAWS, BROCCOLI, CAULIFLOWER, CABBAGE, CARROTS, CELERY, CUCUMBERS, ENDIVE, FENNEL, GARLIC, KALE, KIDNEY BEANS, LEEKS, MINT, PEAS, PARSNIPS, PARSNIP TOPS, PASTINAZ, POTATOES, RUTABAGA, SPINACH, SWEET POTATOS, TURNIPS, TURNIP TOPS, YARROW.

GRASSHOPPERS ON THE ABOVE CROPS

Apply as often as necessary to control grasshoppers on the above crops. See chapter 10 for crop limitations.

SMALL FRUIT INSECT CONTROL

Use sufficient dust to obtain full coverage. Apply when insects are then damage appears. Repeat at 7- to 10-day intervals, or as necessary.

GRAPES: Grape leaf folder, grape leafhopper, and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge (June 10) and a second for leafhoppers. May be applied up to day of harvest.

STRAWBERRIES.—meadow spittlebugs, strawberry leaf miner, and strawberry weevil. May be used up to 7 days before picking.

BLUEBERRIES—Use the "maggot" (larva) and "crabapple" (adult worms) of Japanese beetles. Apply 2 weeks before harvest and repeat 10 days later for a second crop. May be applied up to day of harvest.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES AND RASPBERRIES. Japanese beetle, leaf miner, leafhopper and raspberry aphid. Do not apply within 7 days of harvest.

PET USE

On DOGS and CAES: For packs and their pups, there is no other animal, avoiding the crowd and the struggle into the pack, if only our dead ticks, a few bears, and the hunter. Repeat an already intimate, and a little bit of a sentiment of, between us, just a few more, and more.

in or around dog and cat quarters: In
a dog and cat quarters, the handler should use a
water hose, mop, broom, and/or other means
to remove all dirt, debris, hair, and fecal matter
from the quarters. The dog and cat quarters should
be clean and dry.

NOTICE

[illegible]

5-21-71

Imperial SEVIN DUST

FOR THE CONTROL OF CERTAIN INSECTS ON VEGETABLES, SMALL FRUIT AND PETS

ACTIVE INGREDIENTS:	By Wt.
Carbaryl (1-naphthyl <u>N</u> -methylcarbamate)	5%
INERT INGREDIENTS:	<u>95%</u>
Total	100%

SEVIN is a registered trade mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Skin Contact. Wash Hands Thoroughly After Using. Avoid Contamination of Feed And Foodstuffs. Note for Physician: Carbaryl is a moderately reversible, Cholinesterase inhibitor. Atropine is antidotal.

NET CONTENTS 1 POUND

Manufactured by

IMPERIAL INC.

SHE SANDOVAL HOWA