IMPERIAL SEVIN 50W GARDEN SPRAY

DIRECTIONS

SEEDS [illegible]

VEGETABLES Mix 2 [illegible] gallon of water. Apply [illegible] Apply when insects [illegible] Repeat at 7 to 10-day interval [illegible] longer interval is specified.

May be used up to day of harvest on the following: BEANS, [illegible] CORN [illegible] beetles, leafhoppers, [illegible] CARROTS [illegible] CUCUMBER, SQUASH [illegible] PEPPERS [illegible] TOMATO, EGGPLANT, PEPPERS [illegible]

Do not apply within 3 days of harvest on the following: BEETS, TURNIPS [illegible] BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI [illegible] RADISHES [illegible]

Do not apply within 14 days of harvest on the following: LETTUCE, [illegible]

LAWN INSECT CONTROL [illegible] earwigs, fall armyworm, fleas, leafhoppers, mole cricket, sod webworms, Lawn moths), millipeds, [illegible] Use gallon hose sprayer or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

ORNAMENTAL INSECT CONTROL Ornamental plants, flowers, shrubs and shade trees including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arborvitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston ivy.

Use 2 level tablespoonfuls per gallon of spray (2 pounds per 100 gallons) when insects or their damage appear. Repeat weekly or as needed to control: apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf beetle, elm leaf aphid, flea beetle, gypsy moth, Japanese beetle, lace bugs, leafhoppers, leaf rollers, mealybugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

SMALL FRUIT INSECT CONTROL Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10-day interval or as necessary.

GRAPES Use 2 tablespoonfuls per 1 gallon of water for European fruit lecanium, grape leaf folder, grape leafhopper and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolled leaves needed for leafhopper. May be applied up to day of harvest.

STRAWBERRIES For meadow spittlebug, leaf roller, strawberry weevil, use 2 tablespoonfuls per gallon of water. Make 2 [illegible] bloom applications as [illegible] May be used up to 1 day before picking.

NOTICE Seller warrants that the product conforms to the chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions, under normal conditions of use. [illegible] EXPRESS OR IMPLIED, including any warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, except as [illegible] to the extent that [illegible] contrary to label instructions or [illegible] abnormal conditions or under conditions not [illegible] to other [illegible]

EPA Reg. No. [illegible]

Imperial

SEVIN* 50W Garden Spray

(WETTABLE POWDER CONCENTRATE)

	By Wt.
ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	50%
INERT INGREDIENTS:	50%
Total	100%

*SEVIN is a registered trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust. **Skin Contact May Be Harmful.** Wash Skin Contact With Soap and Water Promptly After Using. **Avoid Contamination of Feed and Foodstuffs.** [illegible]

NET CONTENTS 6 OUNCES

Manufactured By
IMPERIAL INC.
SHENANDOAH, IOWA 51601

ACCEPTED