

IMPERIAL CAPTAN 50W

DIRECTIONS

To control certain plant diseases use this product at the rate of 2½ tablespoons per gallon of water or 2 lbs. per 100 gallons.

APPLES and PEARS — Apple scab, fruit rots, and blotch — make applications at 7 to 10 day intervals as needed to maintain cover in pre-bloom, bloom, petal fall and successive cover sprays. Continue applications to within 30 days of harvest to control blotch, up to start of harvest to control fruit rots.

CHERRIES, PEACH, PLUM, PRUNES AND GRAPE — Brown rot, cherry leaf spot, blossom blight, peach scab, grape black rot — apply at pink bud, full bloom and petal fall stages. Repeat at 10 to 14 day intervals until harvest.

RASPBERRIES, BLACKBERRIES, DEWBERRIES — Anthracnose and botrytis fruit rots — begin application at first bloom. Make 2 repeat applications at 10 to 15 day intervals.

STRAWBERRIES — Botrytis fruit rot and leaf spot — make application when new growth starts and repeat at 7 day intervals through harvest.

TOMATOES — Anthracnose, early and late blight, gray leaf spot, septoria leaf spot — Apply at first sign of disease. Repeat at 5 to 7 day intervals as needed to maintain control. Use 3 tablespoons per gallon of water except where blight conditions are severe. Under these conditions increase dosage to 5 tablespoons per gallon of water. Start application 30 days after first bloom.

POTATOES — Early and late blight — Timing and rates of application same as tomatoes above.

LAWNS & TURF — Brown patch, damping

off, leaf spot, root rot, melting out — Use 2 lbs. per 100 gallons or 2½ tablespoons per gallon and apply to surface at the rate of 10 gallons per 1,000 sq. ft. of turf. Repeat at 7 to 10 day intervals as needed. Apply at first signs of disease. For melting out, increase dosage to 3 lbs. per 100 gallons, when grass resumes growth in the spring. Repeat at 7 to 10 day intervals as long as cool, wet weather persists.

ROSES — Black Spot — Apply at first signs of disease. Repeat at 7 to 10 day intervals.

PREPLANT DIP FOR GLADIOLUS, TUBEROUS BEGONIAS, SEED POTATOES — Seed rot, damping off — Mix 6 to 8 tablespoons per gallon of water. Immerse bulbs in mixture for 20 to 30 minutes. Drain and plant.

CARNATIONS, CHRYSANTHEMUMS — Rust, leaf spot, flower blight — apply at first signs of disease. Repeat applications as needed at 7 to 10 day intervals.

NOTE: Always follow good sanitation practices. Destroy or burn all diseased vegetation.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 407-290

K-8707A-

Imperial

CAPTAN 50W

A FUNGICIDE FOR PLANT DISEASE CONTROL

ACTIVE INGREDIENTS:

Captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide) 50%

INERT INGREDIENTS: 50%

Total 100%

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

Avoid Inhalation of Dust or Spray Mist. Avoid Contact With Skin. Avoid Contamination of Feed or Foodstuffs.

This product is toxic to fish. Keep out of any body of water.

Containers should be disposed of by burying with wastes, or by burning (Keep out of smoke.) Do not contaminate water by cleaning of equipment, or disposal of wastes.

NET CONTENTS 8 OZ.

Manufactured By
IMPERIAL INC.

SHENANDOAH, IOWA 51601

