

DIRECTIONS FOR USE

FIELD CORN

To protect germinating seed and seedlings against seed corn beetles and seed corn maggot, apply 3 to 4 $\frac{1}{2}$ ounces of IMPERIAL SEED TREATER per bushel of seed corn at planting time. (Contents of this package treats 4 to 6 bushels of seed corn.) Thoroughly mix in a box or tub or by dusting it over the seed as it is dumped into the planter box. Thoroughly stir the seed with a paddle or lath so that each seed is coated with the seed protectant. DO NOT USE BARE HANDS.

IMPERIAL SEED TREATER should be applied in conjunction with or over an approved fungicidal seed treatment, but do not use on seed already treated with an insecticide. Plant within two weeks of treatment.

WARNING

May be fatal if swallowed. May be absorbed through the skin. Do not breathe dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of feed and food-stuffs. Keep out of reach of children and domestic animals. Do not use on livestock, household pets, or humans. Do not use treated seed for food, feed or oil purposes.

Toxic to fish and wildlife. Do not contaminate any body of water by cleaning of equipment or disposal of wastes and containers. Treated seed exposed on soil surfaces will be hazardous to birds and wildlife.

Destroy Used Containers: Do not re-use for any purpose. Destroy containers by burning or burying in an isolated area. If burned keep out of smoke.

**TREATED SEED SHOULD BE LABELED AS
FOLLOWS IF IT IS FOR RESALE:**

**TREATED SEED
DO NOT USE FOR FOOD, FEED OR OIL.**

WARRANTY: Imperial Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Imperial Inc., and buyer assumes the risk of any such use.

E

eed corn beetles
PERIAL SEED
ne. (Contents of
oroughly mix in
dumped into the
or lath so that
NOT USE BARE

conjunction with
t do not use on
in two weeks of

gh the skin. Do
clothing. Wash
feed and food-
animals. Do not
not use treated

ody of water by
ainers. Treated
irds and wildlife.
urpose. Destroy
If burned keep

ED AS
:

OIL.

luct conforms to
e purpose stated
as under normal
her warranty of
, express or im-
abel instructions.
t reasonably for-
of any such use.

4-631A-B

ACCEPTED
3-31-70
407-294
Imperial*

DIAZINON* and GRAPHITE

TREATS 4 TO 6 BUSHELS OF SEED CORN

Protects Germinating Seeds and Seedlings
Against

Seed Corn Beetles Seed Corn Maggots

ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl -6-methyl-4-pyrimidinyl) phosphorothioate	33.33%*
INERT INGREDIENTS:	66.67%
Total	100.00%

*Diazinon — Reg. T. M. Geigy Chemical Company

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE OTHER WARNINGS ON BACK PANEL

NET CONTENTS 1 LB. 2 OZS. (Total 18 Ounces)

Manufactured by
Imperial Inc.
SHENANDOAH, IOWA