

407-293

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid inhaling dust. Avoid contamination of foodstuffs and feed. Avoid contact with skin and eyes. May be absorbed through skin, and dust may irritate nasal membrane. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment or disposal of waste. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

S1 AGE: Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

Applicators must wear the following protective clothing during the application process; long-sleeved shirt, long pants. Wear a disposable dust mask covering 1/3 of face or more. GLADIOLUS--

Dust twice a week from early stages until spikes begin to open for control of the following insects and diseases: Thrips, Botrytis, Curvularia, Stemphylium. Repeat as necessary. Continue dusting every 7 to 10 days as long as insects are present.

SPECIAL STORAGE CARE FOR GLADIOLUS

Apply dust to bulbs after they are cleaned (as you put them in storage).

ALSO RECOMMENDED FOR USE ON THESE:

Azaleas, Camellias: To control Petal Blight, apply as least twice a week while flowers are opening. Also apply to soil and litter around the plants while the flowers are in bloom. Repeat every 7 to 10 days for insect control.

Carnations, Chrysanthemums, Hollyhocks, Snapdragons: To control ust, Leaf Spot, and Anthracnose, apply as soon as disease threatens. Repeat very 7 to 10 days or as often as necessary to keep new foliage protected.

Roses: For control of Black Spot, dust at 7 to 10 day intervals and after every rain. For best results, apply before plants become affected.

NOTICE OF WARRANTY

MANUFACTURER AND SELLER MAKE NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extends beyond the statement on this label.

BEST DOCUMENT AVAILABLE