

ACCEPTED
 JUN 21 1971
 U.S. DEPT. OF AGRICULTURE
 THE FEDERAL INSECTICIDE AND RODENTICIDE AGENCY
 ECONOMIC POISON REGISTER
 E.P. NO. SUBJECT
 ATTACHED COMMENTS

Imperial



WARNING
KEEP OUT OF REACH
OF CHILDREN.

See other warnings on side panel.

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene Containing 67% to 69% Chlorine)	58.98%
Xylene Range Aromatic Solvent	38.02%
INERT INGREDIENTS:	3.00%
Total	100.00%

Contains 6 pounds technical toxaphene per gallon
DO NOT STORE NEAR FIRE OR FLAME

NET CONTENTS

 Manufactured by
IMPERIAL INC.
 SHENANDOAH, IOWA 51601

IMPERIAL NO. 6 TOXAPHENE E. C.

WARNING

WHEAT, OATS, BARLEY AND RYE: DO NOT feed treated forage to dairy animals, poultry or animals being finished for slaughter. DO NOT use on forage to be sold commercially or shipped interstate.

SOYBEANS: DO NOT apply within 21 days of harvest. DO NOT feed treated forage to poultry, dairy animals or animals being finished for slaughter. DO NOT feed soybean mill trash to poultry or livestock.

ALFALFA: (GROWN FOR SEED ONLY) Apply in early spring only or after cutting, before new growth is 4 inches tall.

SORGHUM: DO NOT apply at 1 qt. rate per acre within 28 days of harvest. DO NOT apply at 1/2 gallon rate per acre within 40 days of harvest. DO NOT apply at either 1 qt. or 1/2 gallon rate more than once after heads start to form.

CORN: DO NOT feed treated forage to poultry, dairy animals or animals being finished for slaughter.

COTTON: DO NOT graze dairy animals or animals being finished for slaughter on fields receiving late applications.

NOTE: Make these applications on cotton in accordance with time schedule established by State Agricultural Extension Service.

CAUTION

DO NOT take internally. May be fatal if swallowed. **ANTI-DOTE:** Give 1 tablespoon of table salt in glass of warm water and repeat until vomit is clear. CALL PHYSICIAN. DO NOT get in eyes or on skin or clothing. DO NOT breathe vapors or spray mist. In case of contact wash with soap and water.

DO NOT use on household pets, dairy animals, animals being finished for slaughter or humans. DO NOT use in dairy barns or milk houses.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating and burying with waste or by burning. Stay out of smoke or fumes.

Imperial

10% TOXAPHENE

Granular Insecticide

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED FOR EUROPEAN CORN BORER CONTROL

ACTIVE INGREDIENTS:	% by Wt.
Toxaphene (Technical Chlorinated Camphene containing 67-69% chlorine)	10.00%
Aromatic Petroleum Hydrocarbons	1.50%
INERT INGREDIENTS	88.50%
	<u>100.00%</u>

WARNING

KEEP OUT OF REACH OF CHILDREN

Read Entire Label for Additional Handling Precautions.

WARNING: May be fatal if swallowed. Do not get on skin, in eyes or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe dust.

Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams and ponds. Apply only as specified on the label. Dispose of wastes by burying on non-crop lands away from water supplies.

Destroy containers by burning. Keep out of smoke.

ANTIDOTE: Internal — Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately. **External** — Wash with soap and water.

DIRECTIONS FOR USE

European Corn Borer Control — Apply 20 pounds per acre by either ground equipment or by airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second brood borers. Consult State Extension Service Specialists for most effective application dates.

Do not feed ensilage made from treated corn plants to poultry, dairy animals, or animals being finished for slaughter. Do not allow dairy animals or animals being finished for slaughter to graze or feed on treated stover.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-301

K-8500A-I

NET WEIGHT 40 POUNDS

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

ACCEPTED