

DIRECTIONS FOR USE

Dilute with water and use either as follows:

Stock - Tox Two	Amount of Water
1 gallon	100
1 quart	25
1 pint	12 1/2
1 cup	6 1/4

WARNING

DO NOT use in any form on dairy animals under 6 months of age. DO NOT use on new born animals. DO NOT spray meat animals within 30 days of slaughter. DO NOT use on animals within 60 days of slaughter. DO NOT use on animals being treated in cold, stormy weather. Avoid treating over heated animals. Animals should be sprayed on a warm sunny day during winter months to give time to dry before nightfall.

For control of horn flies and ticks on beef cattle, sheep, and hogs, and psoroptic mange on beef cattle and sheep, and to aid in the reduction of lice on beef cattle, sheep and hogs.

SPRAYERS: High pressure sprayers give better results if large numbers of animals are to be treated. Low pressure and hand sprayers are satisfactory for small groups of animals if the spray solution is kept thoroughly mixed.

DIPPING VATS: Dipping vats are effective for treating large groups of animals in one location. Dipping is most effective against mange or scabies where complete coverage of the animal is essential. DO NOT dip meat animals within 60 days of slaughter.

BACKRUBBER USE

1 gallon	10 gallons oil
1 pint	1 1/4 gallons oil

WARNING

DO NOT allow dairy animals or calves under 6 months of age or swine under 3 months of age to have access to backrubbers. DO NOT permit meat animals access to the backrubber within 30 days of slaughter.

BEEF CATTLE AND SWINE: For louse control and to aid in the reduction of lice, dilute 1 gallon with 10 gallons of fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the backrubber every 2 weeks or as often as necessary for control. Locate backrubber to give beef cattle or hogs free access to it, such as at watering places, salubaks, or feed racks. For Fly Control: To aid in the reduction of face flies on beef cattle, one end of backrubber must be no more than 15 or 20 inches above ground for proper face and head treatment. Dilute as for backrubber use.

HOGS: Lice and sarcoptic mange: A single application will usually give control. If more than one treatment appears necessary repeat at 14 day intervals as needed. DO NOT treat sows within 30 days of farrowing nor while they are suckling young. DO NOT spray within 30 days of slaughter.

NET CONTENTS

Imperial

STOCK-TOX TWO
(Livestock Spray Concentrate)


For Use as a Dip or Spray to Aid in the Control of Horn Flies, Lice, Sarcoptic and Psoroptic Mange, and Ticks
NOT FOR USE OR STORAGE IN OR AROUND THE HOME

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene containing 67 to 69% Chlorine)	15.00%
Gamma Isomer of Benzene Hexachloride from Lindane	1.00%
Petroleum Hydrocarbons	39.00%
Xylene Range Solvent	5.00%
INERT INGREDIENTS:	10.00%
Total	100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Precautions
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME



DIRECTIONS FOR USE

WARNING

ANTIDOTE

DO NOT use on dairy animals under 6 months of age. DO NOT use on new born animals. DO NOT use on animals within 30 days of slaughter. DO NOT use on animals within 60 days of slaughter. DO NOT use on animals being treated in cold, stormy weather. Avoid treating over heated animals. Animals should be sprayed on a warm sunny day during winter months to give time to dry before nightfall.

DO NOT reuse this container. Destroy by crushing or perforating and burying with waste in non-crop lands away from water supplies.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-309

K-8616A-J

Manufactured by
IMPERIAL INC.
SHENANDOAH, IOWA 51601